



GMP PE/Cyanine7 anti-human CD11b Antibody

Catalog# / Size 260002 / 100 tests

Clone ICRF44
Workshop IV M047

Other Names Integrin αM chain, C3biR, CR3, Mac-1, Mo1, ITGAM

Isotype Mouse IgG1, κ

Description CD11b is a 165-170 kD type I transmembrane glycoprotein also known as α_M

integrin, Mac-1, CR3, and C3biR. CD11b non-covalently associates with integrin β_2 (CD18) and is expressed on granulocytes, monocytes/macrophages, dendritic cells, NK cells, and subsets of T and B cells. CD11b/CD18 is critical for the transendothelial migration of monocytes and neutrophils. It is also involved in granulocyte adhesion, phagocytosis, and neutrophil activation. CD11b/CD18 interacts with ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4, CD14, CD23, heparin,

iC3b, fibrinogen, and factor X.

Product Details

Reactivity Human

Antibody Type Monoclonal

Host Species Mouse

Formulation Phosphate-buffered solution, pH 7.2, containing True-Stain Monocyte Blocker™,

0.09% sodium azide and 0.2% (w/v) BSA (origin USA), and a stabilizer.

Preparation The antibody was purified by affinity chromatography and conjugated with

PE/Cyanine7 under optimal conditions.

Concentration 200 µg/mL

Storage & Handling The antibody solution should be stored undiluted between 2°C and 8°C, and

protected from prolonged exposure to light. Do not freeze.

Application FC - Quality tested

Recommended Usage Each lot of this antibody is quality control tested by immunofluorescent staining with

flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal

performance for each application.

Excitation Laser Blue Laser (488 nm)

Green Laser (532 nm)/Yellow-Green Laser (561 nm)

Application Notes The ICRF44 antibody inhibits heterotypic adhesion of granulocytes in response to

fMLP. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections, immunofluorescence microscopy⁵, stimulation of monocytes³, blocking of heterotypic PMN aggregation⁸, and blocking of granulocyte activation¹². This clone

was tested in-house and does not work on formalin fixed paraffin-embedded (FFPE) tissue.

The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm

filtered) is recommended for functional assays (Cat. Nos. 301361 & 301362).

Application References

- 1. Knapp W. 1989. Leucocyte Typing IV. Oxford University Press New York.
- (PubMed link indicates BioLegend citation)
- Barclay N, et al. 1997. The Leucocyte Antigen Facts Book. Academic Press Inc. San Diego.
- 3. Rezzonico R, et al. 2001. Blood 97:2932. (Stim)
- 4. Marsik C, et al. 2003. Shock 20:493. (FC)
- 5. David A, et al. 2003. J. Leukoc. Biol. 74:551. (IF)
- 6. Charles N, et al. 2010. Nat. Med. 16:701. (FC) PubMed

- 7. Thurlow LR, et al. 2010. Infect. Immun. 128:1128. (FC) PubMed
- 8. Jadhav S, et al. 2001. J. Immunol. 167:5986. (Block)
- 9. Yoshino N, et al. 2000. Exp. Anim. (Tokyo) 49:97. (FC)
- 10. Sestak K, et al. 2007. Vet. Immunol. Immunopathol. 119:21. (FC)
- 11. Wen T, et al. 2014. J Immunol. 192:5481. (FC) PubMed
- 12. Sprong T, et al. 2003. Blood 102:3702. (Block)

Disclaimer

GMP RUO Flow Cytometry Antibodies. BioLegend GMP RUO fluorophore conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or therapeutic procedures. Our processes include:

- · Batch-to-batch consistency
- · Material traceability
- · Documented procedures
- · Documented employee training
- Equipment maintenance and monitoring records
- · Lot-specific certificates of analysis
- Quality audits per ISO 13485:2016
- · QA review of released products

Antigen Details

Structure Integrin, type I transmembrane glycoprotein, associates with integrin β_2 (CD18),

165-170 kD

Distribution Granulocytes, monocytes/macrophages, dendritic cells, NK cells, subset of T cells,

subset of B cells

Function Adhesion, phagocytosis, chemotaxis, neutrophil activation

Ligand/Receptor ICAM-1(CD54), ICAM-2 (CD102), ICAM-4, CD14, CD23, heparin, iC3b, fibrinogen,

factor X

Cell Type B cells, Dendritic cells, Granulocytes, Macrophages, Monocytes, Neutrophils, NK

cells, T cells, Tregs

Biology Area Cell Adhesion, Cell Biology, Costimulatory Molecules, Immunology, Innate Immunity,

Neuroscience, Neuroscience Cell Markers

Molecular Family Adhesion Molecules, CD Molecules

Antigen References 1. Stewart M, et al. 1995. Curr Opin Cell Biol. 7:690.


Gene ID <u>3684</u>

Related Protocols

Cell Surface Flow Cytometry Staining Protocol

Other Formats

APC anti-human CD11b, Biotin anti-human CD11b, PE anti-human CD11b, PE/Cyanine5 anti-human CD11b, Purified anti-human CD11b, Pacific Blue™ anti-human CD11b, Alexa Fluor® 488 anti-human CD11b, Pe/Cyanine7 anti-human CD11b, PerCP/Cyanine5.5 anti-human CD11b, Brilliant Violet 421™ anti-human CD11b, Brilliant Violet 570™ anti-human CD11b, FITC anti-human CD11b, Brilliant Violet 605™ anti-human CD11b, Brilliant Violet 510™ anti-human CD11b, Brilliant Violet 650™ anti-human CD11b, Brilliant Violet 650™ anti-human CD11b, Purified anti-human CD11b (Maxpar® Ready), Alexa Fluor® 594 anti-human CD11b, APC/Cyanine7 anti-human CD11b, Brilliant Violet 711™ anti-human CD11b, Brilliant Violet 785™ anti-human CD11b, Pe/Dazzle™ 594 anti-human CD11b, APC/Fire™ 750 anti-human CD11b, TotalSeq™-A0161 anti-human CD11b, TotalSeq™-B0161 anti-human CD11b, TotalSeq™-C0161 anti-human CD11b, Ultra-LEAF™ Purified anti-human CD11b, TotalSeq™-D0161 anti-human CD11b



Typical results from human peripheral blood granulocytes stained either with ICRF44 PE/Cyanine7 used at 5 μ L/test (filled histogram) or with an isotype control (open histogram).

Symbols Glossary*

Symbol	Meaning	Symbol Title	Symbol No.	Symbol	Meaning	Symbol Title	Symbol No.
REF	Catalog number	Catalogue number	5.1.6	<u> </u>	Indicates the need for the user to consult the instructions for use.	Consult instructions for use	5.4.3
1	Indicates the temperature limits to which the medical device can be safely exposed.	Temperature limit	5.3.7	淡	Indicates a medical device that needs protection from light sources.	Keep away from sunlight	5.3.2
K	Indicates the upper limit of temperature to which the medical device can be safely exposed.	temperature	5.3.6	Ω	Indicates the date after which the medical device is not to be used.	Use-by date	5.1.4
•••	Indicates the medical device manufacturer.	Manufacturer	5.1.1	EC REP	Indicates the authorized representative in the European Community.	Authorized representative in the European Community	5.1.2
LOT	Indicates the manufacturer's batch code so that the batch or lot can be identified.	Batch code	5.1.5	IVD	Indicates a medical device that is intended to be used as an in vitro diagnostic medical device.	In vitro diagnostic medical device	5.5.1

* Symbol information is from EN ISO 15223-1:2016 Medical devices – Symbols to be used with medical device labels, labelling and information to be supplied – Part 1: General requirements

For research use only. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587