



GMP APC anti-human CD10 Antibody

Catalog# / Size 260022 / 100 tests

Clone HI10a

Workshop V CD10.7

Other Names Common acute lymphoblastic leukemia antigen (CALLA), Enkephalinase, Neutral

endopeptidase, Neprilysin

Isotype Mouse IgG1, κ

Description CD10 is a 100 kD neutral endopeptidase and a member of the metalloprotease

family. It is a type II transmembrane protein also known as common acute lymphoblastic leukemia antigen (CALLA), enkephalinase, and neprilysin. CD10 is expressed on B cell precursors, T cell precursors, and neutrophils. CD10 is involved in B cell development and has been shown to bind opioid enkephalins,

bradykinin, angiotensins I and II, and other biologically active peptides.

Product Details

Reactivity Human

Antibody Type Monoclonal

Host Species Mouse

Formulation Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide, 0.2% (w/v)

BSA (origin USA) and a stabilizer.

Preparation The antibody was purified by affinity chromatography and conjugated with APC

under optimal conditions.

Concentration 100 μg/mL

Storage & Handling The antibody solution should be stored undiluted between 2°C and 8°C, and

protected from prolonged exposure to light. Do not freeze.

Application FC - Quality tested

Recommended Usage Each lot of this antibody is quality control tested by immunofluorescent staining with

flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μL per million cells in 100 μL staining volume or 5 μL per 100 μL of whole blood. It is recommended that the reagent be titrated for optimal

performance for each application.

Excitation Laser Red Laser (633 nm)

Disclaimer GMP RUO Flow Cytometry Antibodies. BioLegend GMP RUO fluorophore

conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or

therapeutic procedures. Our processes include:

Batch-to-batch consistency

· Material traceability

Documented procedures

• Documented employee training

Equipment maintenance and monitoring records

Lot-specific certificates of analysis

Quality audits per ISO 13485:2016

QA review of released products

Structure Type II transmembrane metalloprotease family, 100 kD

Distribution B cell precursors, T cell precursors, neutrophils

Function Zinc-binding metalloproteinase, B cell development

Ligand/Receptor Biologically active peptides including opioid enkephalins, bradykinin, angiotensions

& II

Cell Type B cells, Neutrophils

Biology Area Immunology

Molecular Family CD Molecules

Antigen References 1. Shipp M, et al. 1993. Blood 82:1052.

2. Lu B, et al. 1995. J. Exp. Med. 181:2271.


Gene ID <u>4311</u>

Related Protocols

Cell Surface Flow Cytometry Staining Protocol

Other Formats

Purified anti-human CD10, PE anti-human CD10, PE/Cyanine5 anti-human CD10, FITC anti-human CD10, APC anti-human CD10, APC/cyanine7 anti-human CD10, PE/Cyanine7 anti-human CD10, PerCP/Cyanine5.5 anti-human CD10, Brilliant Violet 421™ anti-human CD10, Brilliant Violet 510™ anti-human CD10, Brilliant Violet 605™ anti-human CD10, Purified anti-human CD10 (Maxpar® Ready), Brilliant Violet 711™ anti-human CD10, PE/Dazzle™ 594 anti-human CD10, APC/Fire™ 750 anti-human CD10, TotalSeq™-A0062 anti-human CD10, TotalSeq™-C0062 anti-human CD10, TotalSeq™-B0062 anti-human CD10, Brilliant Violet 785™ anti-human CD10, TotalSeq™-D0062 anti-human CD10



Typical results from human peripheral blood granulocytes stained either with Hl10a APC used at 5 µL/test (filled histogram) or with an isotype control (open histogram).

Symbols Glossary*

Symbol	Meaning	Symbol Title	Symbol No.	Symbol	Meaning	Symbol Title	Symbol No.
REF	Catalog number	Catalogue number	5.1.6	i	Indicates the need for the user to consult the instructions for use.	Consult instructions for use	5.4.3
1	Indicates the temperature limits to which the medical device can be safely exposed.	Temperature limit	5.3.7	淡	Indicates a medical device that needs protection from light sources.	Keep away from sunlight	5.3.2
1	Indicates the upper limit of temperature to which the medical device can be safely exposed.	temperature	5.3.6	Ω	Indicates the date after which the medical device is not to be used.	Use-by date	5.1.4
•••	Indicates the medical device manufacturer.	Manufacturer	5.1.1	EC REP	Indicates the authorized representative in the European Community.	Authorized representative in the European Community	5.1.2
LOT	Indicates the manufacturer's batch code so that the batch or lot can be identified.	Batch code	5.1.5	IVD	Indicates a medical device that is intended to be used as an in vitro diagnostic medical device.	In vitro diagnostic medical device	5.5.1

* Symbol information is from EN ISO 15223-1:2016 Medical devices – Symbols to be used with medical device labels, labelling and information to be supplied – Part 1: General requirements

For research use only. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587