



GMP APC anti-human CD4 Antibody

Catalog# / Size 260024 / 100 tests

Clone SK3

Other Names T4, Leu3a

Isotype Mouse IgG1, κ

Description CD4, also known as T4, is a 55 kD single-chain type I transmembrane

glycoprotein expressed on most thymocytes, a subset of T cells, and monocytes/macrophages. CD4, a member of the lg superfamily, recognizes antigens associated with MHC class II molecules and participates in cell-cell interactions, thymic differentiation, and signal transduction. CD4 acts as a primary receptor for HIV, binding to HIV gp120. CD4 has also been shown to

interact with IL-16.

Product Details

Reactivity Human

Antibody Type Monoclonal

Host Species Mouse

Formulation Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2%

(w/v) BSA (origin USA).

Preparation The antibody was purified by affinity chromatography and conjugated with APC

under optimal conditions.

Concentration 50 μg/mL

Storage & Handling The antibody solution should be stored undiluted between 2°C and 8°C, and

protected from prolonged exposure to light. Do not freeze.

Application FC - Quality tested

Recommended Usage Each lot of this antibody is quality control tested by immunofluorescent staining with

flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal

performance for each application.

Excitation Laser Red Laser (633 nm)

Application References

1. Evans RL, et al. 1981. Immunol. 78:544

(PubMed link indicates BioLegend citation)

2. Arno A et al. 1999. J. Infect. Dis. 180:56

3. Muech M, et al. 1997. Blood 89:1364

4. Wang L, et al. 2012. Cytometry A. 81:567. PubMed

Disclaimer

GMP RUO Flow Cytometry Antibodies. BioLegend GMP RUO fluorophore conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or therapeutic procedures. Our processes include:

- Batch-to-batch consistency
- · Material traceability
- Documented procedures
- · Documented employee training
- · Equipment maintenance and monitoring records
- · Lot-specific certificates of analysis
- Quality audits per ISO 13485:2016
- · QA review of released products

Antigen Details

Structure Ig superfamily, type I transmembrane glycoprotein, 55 kD

Distribution T cell subset, majority of thymocytes, monocytes/macrophages

Function MHC class II co-receptor, lymphocyte adhesion, thymic differentiation, HIV receptor

Ligand/Receptor MHC class II molecules, HIV gp120, IL-16

Cell Type Macrophages, Monocytes, T cells, Thymocytes, Tregs

Biology Area Immunology

Molecular Family CD Molecules

Antigen References 1. Center D et al. 1996. Immunol. Today 17:476.

2. Gaubin M et al. 1996. Eur. J. Clin. Chem. Clin. Biochem. 34:723.


Gene ID <u>920</u>

Related Protocols

Cell Surface Flow Cytometry Staining Protocol

Other Formats

PerCP anti-human CD4, Purified anti-human CD4, FITC anti-human CD4, PE anti-human CD4, PerCP/Cyanine5.5 anti-human CD4, Biotin anti-human CD4, PE/Cyanine7 anti-human CD4, APC anti-human CD4, APC/Cyanine7 anti-human CD4, Alexa Fluor® 488 anti-human CD4, Pacific Blue™ anti-human CD4, Alexa Fluor® 700 anti-human CD4, Purified anti-human CD4 (Maxpar® Ready), Brilliant Violet 421™ anti-human CD4, Brilliant Violet 510™ anti-human CD4, Alexa Fluor® 647 anti-human CD4, APC/Fire™ 750 anti-human CD4, PE/Dazzle™ 594 anti-human CD4, Brilliant Violet 785™ anti-human CD4, Brilliant Violet 750™ anti-human CD4, Brilliant Violet 605™ anti-human CD4, Brilliant Violet 711™ anti-human CD4, TotalSeq™-A0045 anti-human CD4, TotalSeq™-C0045 anti-human CD4, PE/Cyanine5 anti-human CD4, Spark Blue™ 550 anti-human CD4, Spark NIR™ 685 anti-human CD4, KIRAVIA Blue 520™ anti-human CD4, APC/Fire™ 810 anti-human CD4, PE/Fire™ 640 anti-human CD4, PE/Fire™ 700 anti-human CD4, Spark Violet™ 538 anti-human CD4 Antibody, Spark YG™ 581 anti-human CD4, Alexa Fluor® 660 anti-human CD4, Spark YG™ 593 anti-human CD4, PE/Fire™ 810 anti-human CD4, PE/Fire™ 750 anti-human CD4



Typical results from human peripheral blood lymphocytes stained either with SK3 APC used at 5 µL/test (red histogram) or with an isotype control (green histogram).

Symbols Glossary*

Symbol	Meaning	Symbol Title	Symbol No.	Symbol	Meaning	Symbol Title	Symbol No.
REF	Catalog number	Catalogue number	5.1.6	i	Indicates the need for the user to consult the instructions for use.	Consult instructions for use	5.4.3
1	Indicates the temperature limits to which the medical device can be safely exposed.	Temperature limit	5.3.7	淡	Indicates a medical device that needs protection from light sources.	Keep away from sunlight	5.3.2
1	Indicates the upper limit of temperature to which the medical device can be safely exposed.	temperature	5.3.6	Ω	Indicates the date after which the medical device is not to be used.	Use-by date	5.1.4
	Indicates the medical device manufacturer.	Manufacturer	5.1.1	EC REP	Indicates the authorized representative in the European Community.	Authorized representative in the European Community	5.1.2
LOT	Indicates the manufacturer's batch code so that the batch or lot can be identified.	Batch code	5.1.5	IVD	Indicates a medical device that is intended to be used as an in vitro diagnostic medical device.	In vitro diagnostic medical device	5.5.1

* Symbol information is from EN ISO 15223-1:2016 Medical devices – Symbols to be used with medical device labels, labelling and information to be supplied – Part 1: General requirements

For research use only. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587