

# GMP FITC anti-human CD45 Antibody

Catalog# / Size 260032 / 100 tests

Clone HI30

Other Names LCA, T200

Isotype Mouse IgG1, κ

**Description** CD45 is a 180-240 kD single chain type I membrane glycoprotein also known as leukocyte

common antigen (LCA) and T200. It is a tyrosine phosphatase expressed on the plasma membrane of all hematopoietic cells, except erythrocytes and platelets. CD45 is a signaling molecule that regulates a variety of cellular processes including cell growth, differentiation, cell cycle, and oncogenic transformation. CD45 plays a critical role in T and B cell antigen receptor-mediated activation by dephosphorylating substrates including p56Lck, p59Fyn, and other Src family kinases. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to bind galectin-1 and to be associated with several other cell surface antigens including CD1, CD2,

CD3, and CD4.

#### **Product Details**

Reactivity Human

Antibody Type Monoclonal

Host Species Mouse

Formulation Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin

USA).

Preparation The antibody was purified by affinity chromatography and conjugated with FITC under optimal

conditions.

Concentration 200 µg/mL

Storage & Handling The antibody solution should be stored undiluted between 2°C and 8°C, and protected from

prolonged exposure to light. Do not freeze.

Application FC - Quality tested

Recommended Usage Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric

<u>analysis</u>. For flow cytometric staining, the suggested use of this reagent is 5  $\mu$ L per million cells in 100  $\mu$ L staining volume or 5  $\mu$ L per 100  $\mu$ L of whole blood. It is recommended that the reagent be

titrated for optimal performance for each application.

Excitation Laser Blue Laser (488 nm)

Application Notes Additional reported applications (for the relevant formats) include: immunohistochemical staining of

acetone-fixed frozen tissue sections and formalin-fixed paraffin-embedded tissue sections<sup>9</sup>, inhibition of CD45 functions<sup>4</sup>, immunofluorescence<sup>11</sup>, Western blotting<sup>3</sup>, and spatial biology

(IBEX)<sup>16,17</sup>.

It was found that the HI30 clone and the 2D1 clone can cross block each other's binding.

Application References
(PubMed link indicates

BioLegend citation)

1. Knapp W, et al. 1989. Leucocyte Typing IV. Oxford University Press. New York.

2. Kishihara K, et al. 1993. Cell 74:143.

3. Esser M, et al. 2001. J. Virol. 75:6173. (WB)

4. Yamada T, et al. 2002. J. Biol. Chem. 277:28830.

5. Nagano M, et al. 2007. Blood 110:151.

6. Jiang Q, et al. 2008. Blood 112:2858. PubMed

7. Morozov A, et al. 2010. Clin Cancer Res. 16:5630. PubMed

8. Yoshino N, et al. 2000. Exp. Anim. (Tokyo) 49:97. (FC) 9. Friedman T, et al. 1999. J. Immunol. 162:5256. (IHC)

10. Oeztuerk-Winder F, et al. 2012. EMBO J. 31:3431. (FC) PubMed

11. Rees LE, et al. 2003. Clin. Exp. Immunol. 134:497. (IF)

- 12. Lee J, et al. 2015. J Exp Med. 212:385. PubMed
- 13. Breton G, et al. 2015. J Exp Med. 212:401. PubMed
- 14. Marquardt N, et al. 2015. J Immunol. 6:2467. PubMed
- 15. Bushway ME, et al. 2014. Biol Reprod. 90(5): 110. (IF) PubMed
- 16. Radtke AJ, et al. 2020. Proc Natl Acad Sci USA. 117:33455-33465. (SB) PubMed
- 17. Radtke AJ, et al. 2022. Nat Protoc. 17:378-401. (SB) PubMed

#### Disclaimer

**GMP RUO Flow Cytometry Antibodies.** BioLegend GMP RUO fluorophore conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or therapeutic procedures. Our processes include:

- · Batch-to-batch consistency
- · Material traceability
- Documented procedures
- · Documented employee training
- · Equipment maintenance and monitoring records
- · Lot-specific certificates of analysis
- Quality audits per ISO 13485:2016
- · QA review of released products

### **Antigen Details**

Structure Tyrosine phosphatases, type I transmembrane protein, 180-240 kD (multiple isoforms)

**Distribution** Hematopoietic cells, not expressed in circulating erythrocytes or platelets

Function TCR and BCR mediated activation

Ligand/Receptor Galectin-1, CD2, CD3, CD4

Cell Type Hematopoietic stem and progenitors

Biology Area Cell Biology, Immunology, Inhibitory Molecules, Innate Immunity, Neuroscience, Neuroscience Cell

Markers, Stem Cells

Molecular Family CD Molecules

Antigen References

- 1. Thomas M. 1989. Annu. Rev. Immunol. 7:339.
- 2. Trowbridge I, et al. 1994. Annu. Rev. Immunol.12:85.


Gene ID <u>5788</u>

### **Related Protocols**

Cell Surface Flow Cytometry Staining Protocol

## **Other Formats**

APC anti-human CD45, Biotin anti-human CD45, FITC anti-human CD45, PE anti-human CD45, PE/Cyanine5 anti-human CD45, Purified anti-human CD45, APC/Cyanine7 anti-human CD45, Alexa Fluor® 488 anti-human CD45, Alexa Fluor® 647 anti-human CD45, Pacific Blue™ anti-human CD45, Alexa Fluor® 700 anti-human CD45, PerCP anti-human CD45, PerCP/Cyanine5.5 anti-human CD45, Brilliant Violet 421™ anti-human CD45, Brilliant Violet 570™ anti-human CD45, Brilliant Violet 510™ anti-human CD45, Brilliant Violet 605™ anti-human CD45, Brilliant Violet 650™ anti-human CD45, Purified anti-human CD45 (Maxpar® Ready), Brilliant Violet 785™ anti-human CD45, Brilliant Violet 711™ anti-human CD45, PE/Dazzle™ 594 anti-human CD45, Alexa Fluor® 594 anti-human CD45, APC/Fire™ 750 anti-human CD45, TotalSeq™-A0391 anti-human CD45, TotalSeq™-B0391 anti-human CD45, TotalSeq™-B0391 anti-human CD45, APC/Fire™ 810 anti-human CD45, Spark YG™ 570 anti-human CD45, PE/Fire™ 700 anti-human CD45, Alexa Fluor® 660 anti-human CD45 Antibody, Spark Violet™ 538 anti-human CD45, Spark YG™ 593 anti-human CD45, GMP APC/Fire™ 750 anti-human CD45, Spark UV™ 387 anti-human CD45



Typical results from human peripheral blood lymphocytes stained either with HI30 FITC used at 5 µL/test (filled histogram) or with an isotype control (open histogram).

Symbols Glossary\*

Symbols Glossary							
Symbol	Meaning	Symbol Title	Symbol No.	Symbol	Meaning	Symbol Title	Symbol No.
REF	Catalog number	Catalogue number	5.1.6	$\bigcap_{\mathbf{i}}$	Indicates the need for the user to consult the instructions for use.	Consult instructions for use	5.4.3
1	Indicates the temperature limits to which the medical device can be safely exposed.	Temperature limit	5.3.7	类	Indicates a medical device that needs protection from light sources.	Keep away from sunlight	5.3.2
K	Indicates the upper limit of temperature to which the medical device can be safely exposed.	temperature	5.3.6	Ω	Indicates the date after which the medical device is not to be used.	Use-by date	5.1.4
•••	Indicates the medical device manufacturer.	Manufacturer	5.1.1	EC REP	Indicates the authorized representative in the European Community.	Authorized representative in the European Community	5.1.2
	Indicates the manufacturer's	Batch code	5.1.5		Indicates a medical device that is	In vitro diagnostic	5.5.1
LOT	batch code so that the batch or lot can be identified.			IVD	intended to be used as an in vitro diagnostic medical device.	medical device	

<sup>\*</sup> Symbol information is from EN ISO 15223-1:2016 Medical devices – Symbols to be used with medical device labels, labelling and information to be supplied – Part 1: General requirements

For research use only. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587