

GMP APC anti-human CD28 Antibody

Catalog# / Size	260250 / 100 tests
Clone	CD28.2
Workshop	V-CD28.05
Other Names	T44, Tp44
Isotype	Mouse IgG1, κ
Description	CD28 is a 44 kD disulfide-linked homodimeric type I glycoprotein. It is a member of the immunoglobulin superfamily and is also known as T44 or Tp44. CD28 is expressed on most T lineage cells, NK cell subsets, and plasma cells. CD28 binds both CD80 and CD86 using a highly conserved motif MYPPY in the CDR3-like loop. CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death. <i>In vitro</i> studies indicate that ligation of CD28 on T cells by CD80 and CD86 on antigen presenting cells provides a costimulatory signal required for T cell activation and proliferation.

Product Details

Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Preparation	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
Concentration	100 µg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Red Laser (633 nm)
Application Notes	The Ultra-LEAF™ Purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for highly sensitive assays.

Application References

(PubMed link indicates BioLegend citation)

- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
- Nunes J, *et al.* 1993. *Biochem. J.* 293:835.
- Calea-Lauri J, *et al.* 1999. *J. Immunol.* 163:62.
- Tazi A, *et al.* 1999. *J. Immunol.* 163:3511. (IHC)
- Marti F, *et al.* 2001. *J. Immunol.* 166:197. (Costim)
- Jeong SH, *et al.* 2004. *J. Virol.* 78:6995. (Costim)
- Rivollier A, *et al.* 2004. *Blood* 104:4029. (Costim)
- Scharschmidt E, *et al.* 2004. *Mol. Cell Biol.* 24:3860. (Costim)
- Sheng W, *et al.* 2007. *Elsevier* 580:6819. [PubMed](#)
- Mitsuhashi M. 2007. *Clin Chem.* 53:148. [PubMed](#)
- Ye Z, *et al.* 2008. *Infect. Immun.* 76:2541. [PubMed](#)
- Magatti M, *et al.* 2008. *Stem Cells* 26:182. (FA) [PubMed](#)
- Yoshino N, *et al.* 2008. *Exp. Anim. (Tokyo)* 49:97. (FC)
- Berg M, *et al.* 2008. *J Leukoc Biol.* 83:853. (IP) [PubMed](#)
- Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
- Leonard JA, *et al.* 2011. *J. Virol.* 85:6867. [PubMed](#)

Disclaimer

GMP RUO Flow Cytometry Antibodies. BioLegend GMP RUO fluorophore conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or therapeutic procedures. Our processes include:

- Batch-to-batch consistency
- Material traceability
- Documented procedures
- Documented employee training
- Equipment maintenance and monitoring records
- Lot-specific certificates of analysis
- Quality audits per ISO 13485:2016
- QA review of released products

Antigen Details

Structure	Ig superfamily, type I transmembrane glycoprotein, homodimer, 44 kD
Distribution	Mature T cells, thymocytes, NK cell subsets, plasma cells, EBV-positive B cells
Function	T cell costimulation
Ligand/Receptor	CD80, CD86
Cell Type	B cells, NK cells, Plasma cells, T cells, Thymocytes, Tregs
Biology Area	Costimulatory Molecules, Immunology
Molecular Family	CD Molecules
Antigen References	1. Schlossman S, <i>et al.</i> Eds. 1995. <i>Leucocyte Typing V</i> . Oxford University Press. New York. 2. June CH, <i>et al.</i> 1994. <i>Immunol. Today</i> 15:321. 3. Linskey PS, <i>et al.</i> 1993. <i>Annu. Rev. Immunol.</i> 11:191.
Gene ID	940


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-human CD28, Biotin anti-human CD28, FITC anti-human CD28, PE anti-human CD28, PE/Cyanine5 anti-human CD28, Purified anti-human CD28, Alexa Fluor® 488 anti-human CD28, Alexa Fluor® 700 anti-human CD28, PerCP/Cyanine5.5 anti-human CD28, Pacific Blue™ anti-human CD28, PE/Cyanine7 anti-human CD28, Ultra-LEAF™ Purified anti-human CD28, Brilliant Violet 421™ anti-human CD28, Brilliant Violet 510™ anti-human CD28, Purified anti-human CD28 (Maxpar® Ready), PE/Dazzle™ 594 anti-human CD28, Brilliant Violet 785™ anti-human CD28, Brilliant Violet 650™ anti-human CD28, Brilliant Violet 711™ anti-human CD28, APC/Fire™ 750 anti-human CD28, Alexa Fluor® 647 anti-human CD28, TotalSeq™-A0386 anti-human CD28, TotalSeq™-B0386 anti-human CD28, TotalSeq™-C0386 anti-human CD28, Brilliant Violet 605™ anti-human CD28, APC/Cyanine7 anti-human CD28, Brilliant Violet 750™ anti-human CD28, PE/Fire™ 810 anti-human CD28, GMP PE anti-human CD28, TotalSeq™-D0386 anti-human CD28, Spark Violet™ 423 anti-human CD28

Product Data



Typical results from human peripheral blood lymphocytes stained either with CD28.2 APC used at 5 μ L/test (filled histogram) or with an isotype control (open histogram).

For research use only. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587