

Brilliant Violet 570™ anti-mouse TCR β chain Antibody

Catalog# / Size	109231 / 125 µL
Clone	H57-597
Regulatory Status	RUO
Other Names	TCR-β chain, TCR-β, β-TCR
Isotype	Armenian Hamster IgG
Description	T cell receptor (TCR) is a heterodimer consisting of an α and a β chain (TCR α/β) or a γ and a δ chain (TCR γ/δ). TCR-β is a member of the immunoglobulin superfamily and a component of the CD3/TCR complex (along with TCR-α). It is expressed on α/β TCR-bearing T cells and thymocytes. The CD3/TCR complex plays a key role in antigen recognition, signal transduction, and T cell activation.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Armenian Hamster
Immunogen	Affinity purified TCR from mouse DO-11.10 cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
Preparation	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 570™ under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.</p> <p>Brilliant Violet 570™ excites at 405 nm and emits at 570 nm. The bandpass filter 585/42 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 570™ is a trademark of Sirigen Group Ltd.</p> <p>Learn more about Brilliant Violet™.</p> <p>This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.</p>
Excitation Laser	Violet Laser (405 nm)
Application Notes	H57-597 is a hamster mAb directed to an epitope of the C region of TCR β chain ¹² . The H57-597 antibody does not cross-react with γ/δ TCR-bearing T cells. Immobilized or soluble H57-597 antibody can activate α/β TCR-bearing T cells. Additional reported applications (for the relevant formats) for this antibody include: immunoprecipitation ² , <i>in vitro</i> stimulation ^{2,3} , <i>in vivo</i> depletion ⁴⁻⁶ , and immunohistochemical staining of acetone-fixed frozen sections ^{7,8,9} . The Ultra-LEAF™ purified

antibody (Endotoxin <0.01 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 109253-109258).

Application References

(PubMed link indicates BioLegend citation)

1. Gascoigne NJ. 1990. *J. Biol. Chem.* 265:9296.
2. Kruisbeek A, *et al.* 1991. *In Current Protocols in Immunology.* pp. 3.12.1. (Costim IP)
3. Davenport C, *et al.* 1995. *J. Immunol.* 155:3742. (Costim)
4. Drobyski W, *et al.* 1996. *Blood* 87:5355. (Deplete)
5. Kummer U, *et al.* 2001. *Immunol. Lett.* 75:153. (Deplete)
6. van der Heyde HC, *et al.* 1995. *J. Immunol.* 154:3985. (Deplete)
7. Tomita K, *et al.* 1999. *Genes Dev.* 13:1203. (IHC)
8. Podd BS, *et al.* 2006. *J. Immunol.* 176:6532. (IHC)
9. Ponomarev ED, *et al.* 2007. *J. Immunol.* 178:39. (IHC)
10. Chappaz S, *et al.* 2007. *Blood* doi:10.1182/blood-2007-02-074245. (FC) [PubMed](#)
11. Tsukumo S, *et al.* 2006. *J. Immunol.* 177:8365. (FC) [PubMed](#)
12. Grégoire C, *et al.* 1991. *Proc. Natl. Acad. Sci USA* 88:8077.

RRID

AB_10900242 (BioLegend Cat. No. 109231)

Antigen Details

Structure	Ig superfamily, CD3/TCR complex with CD3 and TCR α subunit
Distribution	Majority of T cells and thymocytes (correlated to differentiation)
Function	Antigen recognition, T cell activation
Ligand/Receptor	Peptide bound-MHC class I and II
Antigen References	<ol style="list-style-type: none">1. Davis MM, <i>et al.</i> 1998. <i>Ann. Rev. Immunol.</i> 16:523.2. Huppa JB, <i>et al.</i> 2003. <i>Nat. Immunol.</i> 4:749.3. Kubo R, <i>et al.</i> 1989. <i>J. Immunol.</i> 142:2736.
Gene ID	21577


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-mouse TCR β chain, Biotin anti-mouse TCR β chain, FITC anti-mouse TCR β chain, PE anti-mouse TCR β chain, PE/Cyanine5 anti-mouse TCR β chain, Purified anti-mouse TCR β chain, Alexa Fluor® 488 anti-mouse TCR β chain, Alexa Fluor® 647 anti-mouse TCR β chain, APC/Cyanine7 anti-mouse TCR β chain, PE/Cyanine7 anti-mouse TCR β chain, Alexa Fluor® 700 anti-mouse TCR β chain, Pacific Blue™ anti-mouse TCR β chain, Brilliant Violet 421™ anti-mouse TCR β chain, PerCP/Cyanine5.5 anti-mouse TCR β chain, Brilliant Violet 570™ anti-mouse TCR β chain, Brilliant Violet 510™ anti-mouse TCR β chain, Purified anti-mouse TCR β chain (Maxpar® Ready), Alexa Fluor® 594 anti-mouse TCR β chain, PE/Dazzle™ 594 anti-mouse TCR β chain, Brilliant Violet 605™ anti-mouse TCR β chain, Brilliant Violet 711™ anti-mouse TCR β chain, APC/Fire™ 750 anti-mouse TCR β chain, TotalSeq™-A0120 anti-mouse TCR β chain, Brilliant Violet 785™ anti-mouse TCR β chain, Brilliant Violet 650™ anti-mouse TCR β chain, Ultra-LEAF™ Purified anti-mouse TCR β chain, TotalSeq™-C0120 anti-mouse TCR β chain, TotalSeq™-B0120 anti-mouse TCR β chain

Product Data



C57BL/6 mouse splenocytes were stained with TCR β (clone H57-597) Brilliant Violet 570™ (filled histogram) or Armenian hamster IgG Brilliant Violet 570™ isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587