

PerCP/Cyanine5.5 anti-H2A.X-Phosphorylated (Ser139) Antibody

Catalog# / Size	613413 / 25 tests 613414 / 100 tests
Clone	2F3
Regulatory Status	RUO
Other Names	H2A.x, H2a/x, Histone 2A, Histone 2A.X, Gamma-H2AX
Isotype	Mouse IgG1, κ
Description	H2A.X is a 14 kD basal histone and a member of the H2 histone family. This nuclear protein is synthesized in the G1 and S phase of the cell cycle and is known to be important for DNA repair and maintaining genomic stability and for recombination between immunoglobulin switch regions. H2A.X becomes phosphorylated on serine 139 after double-stranded DNA breaks. Phosphorylated H2A.X promotes DNA repair and maintains genomic stability. The 2F3 monoclonal antibody reacts with phosphorylated human H2A.X (Ser139) and has been shown to be useful for Western blotting, immunofluorescence and flow cytometry.

Product Details

Verified Reactivity	Human, Mouse
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Modified peptide
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood. * PerCP/Cyanine5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.
Application Notes	Additional reported applications (for the relevant formats of this clone) include: immunohistochemistry on paraffin embedded sections ² , immunofluorescence microscopy ³⁻⁹ , Western blotting ¹⁰⁻¹² , and flow cytometry ^{1,13} . Clone 2F3 cross-reacts with mouse ⁴ . Intracellular staining protocol for Anti-H2A.X-Phosphorylated (Ser139) Antibody for Flow Cytometry 1. Prepare 70% absolute ethanol. Chill solution by storing at -20°C. 2. Prepare cells of interest. 3. Wash 1X with PBS, centrifuge at 350g for 5 min. 4. Discard the supernatant and vortex to loosen cell pellet. 5. Add pre-cooled 70% ethanol drop by drop, while vortexing. 6. Incubate at -20°C for 60 minutes. 7. Wash 3X with BioLegend Cell Staining Buffer and resuspend the cells at 0.5-1 X 10 ⁷ cells/ml in the cell staining buffer. 8. Perform immunofluorescent staining for flow cytometry.
Additional Product Notes	BioLegend is in the process of converting the name PerCP/Cy5.5 to PerCP/Cyanine5.5. The dye

molecule remains the same, so you should expect the same quality and performance from our PerCP/Cyanine5.5 products. Contact [Technical Service](#) if you have any questions.

Application References

(PubMed link indicates BioLegend citation)

1. Jha JC, *et al.* 2013. *J. Virol.* 87:5255. (FC) [PubMed](#)
2. Akbay A, *et al.* 2008. *Am J Pathol.* 173:536. (IHC) [PubMed](#)
3. Mochizuki K, *et al.* 2008. *J cell Sci.* 121:2148. (IF) [PubMed](#)
4. Xiao R, *et al.* 2007. *Mol Cell Biol.* 27:5393. (IF) [PubMed](#)
5. Rossi DJ, *et al.* 2007. *Nature.* 447:725. (IF) [PubMed](#)
6. Loidl J, *et al.* 2009. *Mol Cell Biol.* 20:2048. (IF) [PubMed](#)
7. Beels L, *et al.* 2009. *Circulation.* 120:1903. (IF) [PubMed](#)
8. Suzuki K, *et al.* 2010. *Nucleic Acids Res.* 38:e129. (IF) [PubMed](#)
9. Lukaszewicz A. 2010. *Chromasoma* Apr 27. [Epub ahead of print] (IF) [PubMed](#)
10. Yamada C, *et al.* 2010. *J. Biol. Chem.* 285:16693. (WB) [PubMed](#)
11. Bu Y, *et al.* 2010. *Biochem Biophys Res Commun.* 397:157. (WB) [PubMed](#)
12. Massignan T, *et al.* 2010. *J. Biol Chem.* 285:7752. (WB) [PubMed](#)
13. Banath JP, *et al.* 2010. *BMC Cancer* 10:4 (FC)
14. Zhang M., *et al.* 2011. *Cancer Res.* 23:7155. [PubMed](#)
15. Kuefner MA, *et al.* 2012. *Radiology* 264:59. [PubMed](#)
16. Yoshihara Y, *et al.* 2012. *Biochem Biophys Res Commun.* 421:57. [PubMed](#)
17. Titus S, *et al.* 2013. *Sci Transl Med.* 13:21. [PubMed](#)
18. Crown KN, *et al.* 2013. *G3.* 6:1927. [PubMed](#)
19. Schenkwein D, *et al.* 2013. *Nucleic Acids Res.* 41:e61. [PubMed](#)
20. Zhadanova NS, *et al.* 2014. *Mol Cell Biol.* 34:2786. [PubMed](#)
21. Horrell SA, *et al.* 2014. *Eukaryot Cell.* 13:1300. [PubMed](#)
22. Maya-Mendoza A, *et al.* 2015. *Mol Oncol.* 9:601. [PubMed](#)

Product Citations

1. Carpenter RS, *et al.* 2020. *Nat Commun.* 3.029166667. [PubMed](#)

RRID

AB_2616872 (BioLegend Cat. No. 613413)
AB_2616873 (BioLegend Cat. No. 613414)


Antigen Details

Structure	Basal histone, H2 histone family; 14 kD
Distribution	Nuclear
Function	Phosphorylated H2AX promotes DNA repair and maintains genomic stability. Important for recombination between immunoglobulin switch regions
Modification	Phosphorylation on Ser139 after double-stranded DNA breaks
Biology Area	Cell Biology, Chromatin Remodeling/Epigenetics, DNA Repair/Replication, Neuroscience
Molecular Family	Phospho-Proteins
Antigen References	<ol style="list-style-type: none">1. Mannironi C, <i>et al.</i> 1989. <i>Nucleic Acids Res.</i> 17:9113.2. Celeste A, <i>et al.</i> 2002. <i>Science</i> 296:922.3. Bassing CH, <i>et al.</i> 2002. <i>Proc. Natl. Acad. Sci. USA</i> 99:8173.4. Reina-San-Martin B, <i>et al.</i> 2003. <i>J. Exp. Med.</i> 197:1767.
Regulation	Synthesized in G1 and S-phase of cell cycle
Gene ID	3014

Other Formats

Purified anti-H2A.X Phospho (Ser139), FITC anti-H2A.X Phospho (Ser139), Alexa Fluor® 488 anti-H2A.X Phospho (Ser139), Alexa Fluor® 647 anti-H2A.X Phospho (Ser139), Alexa Fluor® 594 anti-H2A.X Phospho (Ser139), PE anti-H2A.X Phospho (Ser139), PerCP/Cyanine5.5 anti-H2A.X-Phosphorylated (Ser139), Direct-Blot™ HRP anti-H2A.X-Phosphorylated (Ser139), APC anti-H2A.X-Phosphorylated (Ser139), PE/Cyanine7 anti-H2A.X Phospho (Ser139), APC/Fire™ 750 anti-H2A.X Phospho (Ser139)

Product Data



Nocodazole treated HeLa cells (24 hours) were fixed and permeabilized with cold 70% ethanol, then intracellularly stained with anti-H2A.X Phospho (Ser139) (clone 2F3) PerCP/Cyanine5.5 (filled histogram) or mouse IgG1, κ PerCP/Cyanine5.5 isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587