

True-Nuclear™ Human Treg Flow™ Kit (FOXP3 Alexa Fluor® 488/CD4 PE-Cy5/CD25 PE)

Catalog# / Size	320403 / 25 tests
Clone	259D
Regulatory Status	RUO
Other Names	Forkhead box protein P3, Scurfin, JM2, IPEX, Zinc finger protein JM2
Description	<p>FOXP3 is a 50-55 kD transcription factor, also known as Forkhead box protein P3, Scurfin, JM2, or IPEX. It is proposed to be a master regulatory gene and more specific marker of T regulatory cells than most cell surface markers (such as CD4 and CD25). Transduced expression of FOXP3 in CD4⁺/CD25⁻ cells has been shown to induce GITR, CD103, and CTLA4 and impart a T regulatory cell phenotype. FOXP3 is mutated in X-linked autoimmunity-allergic dysregulation syndrome (XLAAD or IPEX) in humans and in "scurfy" mice. Overexpression of FOXP3 has been shown to lead to a hypoactive immune state suggesting that this transcriptional factor is a central regulator of T cell activity. In human, unlike in mouse, two isoforms of FOXP3 have been reported: one (FOXP3) corresponding to the canonical full-length sequence; the other (FOXP3 δ2) lacking exon 2. The 259D antibody recognizes human FOXP3 epitope in the region of amino acids 105-235.</p>

Product Details

Verified Reactivity	Human
Reported Reactivity	Cynomolgus, Rhesus, Baboon, Chimpanzee
Antibody Type	Monoclonal
Immunogen	Full-length FOXP3 protein
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	This kit is guaranteed for six months. Upon receipt, store between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Application Notes	<p>Materials Provided:</p> <ol style="list-style-type: none">1. Alexa Fluor® 488 anti-human FOXP3 - 25 tests2. Alexa Fluor® 488 Mouse IgG1, κ isotype control - 25 tests3. True-Nuclear™ Transcription Factor Buffer Set - 120 tests (Cat. No. 424401)4. anti-human CD4 PE-Cy5/CD25 PE Cocktail - 50 tests <p>Materials Not Included:</p> <ol style="list-style-type: none">1. Cell Staining Buffer (Cat. No. 420201)2. Single color compensation controls <p>Immunofluorescence Staining Procedures:</p> <ol style="list-style-type: none">1. Perform cell surface staining as described in BioLegend's Cell Surface Immunofluorescence Staining Protocol. Add 20 µL of the anti-human CD4 PE-Cy5/CD25 PE cocktail to each tube and incubate in the dark for 20 minutes.2. Add 2 mL of the cell staining buffer, centrifuge tubes at 400 x g at room temperature for five minutes, and discard the supernatant.3. Repeat Step 2, for a total of two washes.4. Add 1 mL of the Transcription Factor 1X Fix solution to each tube, vortex, and incubate at room temperature in the dark for 45-60 minutes.5. Without washing, add 2 mL of the Transcription Factor 1X Perm Buffer to each tube.6. Centrifuge tubes at 400 x g at room temperature for five minutes, and discard the supernatant.7. Add 2 mL of the Transcription Factor 1X Perm Buffer to each tube.8. Centrifuge tubes at 400 x g at room temperature for five minutes, and discard the supernatant.9. Resuspend the cell pellet in 100 µL of the Transcription Factor 1X Perm Buffer.10. Add 5 µL of Alexa Fluor® 488 anti-human FOXP3 antibody or 5 µL of Alexa Fluor® 488 mouse IgG1, κ isotype control into the appropriate tubes. Incubate in the dark at room temperature for at

least 30 minutes.

11. Add 2 mL of the Transcription Factor 1X Perm Buffer to each tube.
12. Centrifuge tubes at 400 x g at room temperature for five minutes, and discard the supernatant.
13. Add 2 mL of the cell staining buffer.
14. Centrifuge tubes at 400 x g at room temperature for five minutes, and discard the supernatant.
15. Resuspend in 0.5 mL cell staining buffer and then acquire tubes on a flow cytometer.

Caution: The True-Nuclear™ Transcription Factor Buffer Set contains paraformaldehyde, which is toxic and mutagenic. Please handle with caution. Wear gloves, lab coats, and necessary protection to avoid direct contact.

NOTE: For flow cytometric staining with this clone, True-Nuclear™ Transcription Factor Buffer Set (Cat. No. [424401](#)) offers improved staining and is highly recommended.

Application References

(PubMed link indicates BioLegend citation)

1. Roncador G, *et al.* 2005 *Eur. J. Immunol.* 35:1681.
2. Yang ZZ, *et al.* 2006. *Blood* 107:3639. [PubMed](#)
3. Gavin MA, *et al.* 2006. *P. Natl. Acad. Sci. USA* 103:6659. [PubMed](#)
4. Groh V, *et al.* 2006. *Nature Immunology* 7:755. [PubMed](#)
5. Tran DQ, *et al.* 2007. *Blood* doi:10.1182/blood-2007-06-094656. [PubMed](#)
6. Long SA, *et al.* 2008. *J Autoimmun.* 30:293. [PubMed](#)
7. Gong G, *et al.* 2009. *Blood* 113:837. [PubMed](#)
8. Long SA, *et al.* 2009. *Eur J. Immunol.* 39:612. [PubMed](#)
9. Long SA, *et al.* 2010. *Diabetes.* 59:407. [PubMed](#)
10. Ferraro A, *et al.* 2014. *PNAS.* 111:1111. [PubMed](#)
11. Vudattu NK, *et al.* 2014. *J Immunol.* 193:587. [PubMed](#)
12. Dupont G, *et al.* 2014. *Cytokine.* 69:146. [PubMed](#)


Antigen Details

Structure	Forkhead/winged-helix transcription factor family, approximately 50 kD, contains zinc finger and forkhead domains
Distribution	Nuclear; expressed in T regulatory cells
Function	Transcription factor proposed to be a master regulatory gene in T regulatory cell development and a critical factor for immune homeostasis
Interaction	Interacts with DNA
Cell Type	Tregs
Biology Area	Cell Biology, Immunology, Transcription Factors
Molecular Family	CD Molecules, Nuclear Markers
Antigen References	1. Hori S, <i>et al.</i> 2003. <i>Science</i> 299:1057.
Regulation	FOXP3 is present at high levels in T regulatory cell can also be induced by T cell activation
Gene ID	50943


Other Formats

PE anti-human FOXP3, Purified anti-human FOXP3, Alexa Fluor® 488 anti-human FOXP3, Alexa Fluor® 647 anti-human FOXP3, Pacific Blue™ anti-human FOXP3, True-Nuclear™ Human Treg Flow™ Kit (FOXP3 Alexa Fluor® 488/CD4 PE-Cy5/CD25 PE)

Product Data



Human peripheral blood lymphocytes were stained with True-Nuclear™ Human Treg Flow™ Kit (FOXP3 Alexa Fluor® 488/CD4 PE-Cy5/CD25 PE).



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587