

Ultra-LEAF™ Purified anti-mouse CD48 Antibody

Catalog# / Size	103456 / 100 mg 103454 / 25 mg 103455 / 50 mg 103452 / 1 mg 103453 / 5 mg
Clone	HM48-1
Regulatory Status	RUO
Other Names	BCM1, SLAMF2, Blast-1
Isotype	Armenian Hamster IgG
Description	CD48 is a 45 kD GPI-anchored glycoprotein also known as BCM1, Blast-1 (human), and OX-45 (rat). It is a member of the Ig superfamily, expressed on T and B cells and monocytes/macrophages. It plays a role in adhesion and T cell recognition. The primary ligands for CD48 are CD2 and CD244.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Armenian Hamster
Immunogen	Mouse T lymphoma MBL-2
Formulation	0.2 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is < 0.01 EU/µg of the protein (< 0.001 ng/µg of the protein) as determined by the LAL test.
Concentration	The antibody is bottled at the concentration indicated on the vial, typically between 2 mg/mL and 3 mg/mL. Older lots may have also been bottled at 1 mg/mL. To obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. This Ultra-LEAF™ solution contains no preservative; handle under aseptic conditions.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes	The HM48-1 antibody is useful for blocking <i>in vitro</i> and <i>in vivo</i> CD48 mediated interactions. Additional reported applications (for the relevant formats) include: immunoprecipitation ¹ , costimulation of T cell proliferation ^{1,2} , blocking of CD48-CD2 interaction ¹ , and inhibition of CTL activity and graft rejection ^{1,2} . For <i>in vivo</i> studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. Nos. 103452-103456) with a lower endotoxin limit (Endotoxin < 0.01 EU/µg).
Application References	1. Kato K, <i>et al.</i> 1992. <i>J. Exp. Med.</i> 176:1241. (IP, Costim, Block) 2. Qin L, <i>et al.</i> 1994. <i>J. Exp. Med.</i> 179:341. (Costim, Block)
(PubMed link indicates BioLegend citation)	
RRID	AB_2832312 (BioLegend Cat. No. 103456) AB_2832310 (BioLegend Cat. No. 103454) AB_2832311 (BioLegend Cat. No. 103455) AB_2832308 (BioLegend Cat. No. 103452) AB_2832309 (BioLegend Cat. No. 103453)

Antigen Details

Structure	Ig superfamily, 45 kD
Distribution	T cells and B cells, monocytes/macrophages
Function	Adhesion, T cell recognition
Ligand/Receptor	CD2, CD244
Cell Type	B cells, Macrophages, Monocytes, T cells
Biology Area	Immunology
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	<ol style="list-style-type: none">1. Barclay AN, <i>et al.</i> 1997. The Leukocyte Antigen FactsBook Academic Press.2. Flament C, <i>et al.</i> 1996. <i>Hum. Immunol.</i> 46:82.3. Van der Merwe PA, <i>et al.</i> 1995. <i>Curr. Biol.</i> 5:74.4. Latchman Y, <i>et al.</i> 1998. <i>J. Immunol.</i> 161:5809.
Gene ID	12506


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

FITC anti-mouse CD48, PE anti-mouse CD48, Purified anti-mouse CD48, Biotin anti-mouse CD48, APC anti-mouse CD48, Alexa Fluor® 488 anti-mouse CD48, Alexa Fluor® 647 anti-mouse CD48, Pacific Blue™ anti-mouse CD48, PE/Cyanine5 anti-mouse CD48, PerCP/Cyanine5.5 anti-mouse CD48, PE/Cyanine7 anti-mouse CD48, Alexa Fluor® 700 anti-mouse CD48, Brilliant Violet 421™ anti-mouse CD48, APC/Cyanine7 anti-mouse CD48, Purified anti-mouse CD48 (Maxpar® Ready), Alexa Fluor® 594 anti-mouse CD48, Brilliant Violet 605™ anti-mouse CD48, Brilliant Violet 510™ anti-mouse CD48, Brilliant Violet 711™ anti-mouse CD48, PE/Dazzle™ 594 anti-mouse CD48, APC/Fire™ 750 anti-mouse CD48, TotalSeq™-A0429 anti-mouse CD48, Brilliant Violet 785™ anti-mouse CD48, Ultra-LEAF™ Purified anti-mouse CD48, TotalSeq™-B0429 anti-mouse CD48, TotalSeq™-C0429 anti-mouse CD48

Product Data



C57BL/6 mouse splenocytes stained with Ultra-LEAF™ purified HM48-1, followed by anti-Armennian hamster IgG FITC

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587

