

Alexa Fluor® 647 anti-SOX2 Antibody

Catalog# / Size	656108 / 100 µg
Clone	14A6A34
Regulatory Status	RUO
Other Names	SRY-related HMG-box gene 2, SRY (sex determining region Y)-box 2, MCOPS3, ANOP3
Isotype	Mouse IgG1, κ
Description	SOX2 is the most studied member of SRY-related box transcription factor family. It binds to target genes through its highly conserved HMG box domain. Inactivation of the SOX2 gene causes lethality during embryonic development. SOX2 knockdown in embryonic stem cells results in their differentiation. Co-expression of SOX with OCT4, MYC, and KLF4 is sufficient to reprogram somatic cells to induced pluripotent stem cells (iPSCs), which exert similar characteristics as natural pluripotent stem cells. These findings indicate that SOX2 is crucial for the self-renewal and pluripotency of embryonic stem cells. In addition, over-expression of SOX2 has been found in various types of malignant cancer. Knockdown of SOX2 results in cell cycle arrest by downregulating cyclin D1 and inhibition of tumor cell proliferation, suggesting that SOX2 is involved in activating genes associated with tumor progression.

Product Details

Verified Reactivity	Human, Mouse
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Full length human SOX recombinant protein
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICC - Quality tested FC - Verified
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunocytochemistry. For immunocytochemistry, a concentration range of 1.0 - 5.0 µg/ml is recommended. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.</p> <p>* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633 nm / 635 nm.</p> <p>Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation.</p> <p>View full statement regarding label licenses</p>
Excitation Laser	Red Laser (633 nm)
Application Notes	This clone is not recommended for ChIP (Chromatin Immunoprecipitation) assays (as determined by in-house testing).
Product Citations	<ol style="list-style-type: none">Shrestha R, <i>et al.</i> 2020. Stem Cell Research. 41:101590.. PubMedShrestha R, <i>et al.</i> 2019. Cells. 8:. PubMed
RRID	AB_2563681 (BioLegend Cat. No. 656108)

Antigen Details

Structure	317 amino acids, predicted molecular weight of 34 kD, contains a HMG box domain responsible for DNA binding
Distribution	Nucleus
Function	Transcription factor that regulates the expression of the genes involved in embryonic development
Interaction	Interacts with FGFR1, SOX3, and ZSCAN10
Cell Type	Embryonic Stem Cells, Mesenchymal Stem Cells, Neural Stem Cells
Biology Area	Cell Biology, Cell Cycle/DNA Replication, Immunology, Neuroscience, Neuroscience Cell Markers, Stem Cells, Transcription Factors
Antigen References	<ol style="list-style-type: none">1. Rizzino A. 2009. <i>Wiley Interdiscip. Rev. Syst. Biol. Med.</i> 1:228.2. Stolzenburg S, et al. 2012. <i>Nucleic Acids Res.</i> 40:6725.3. Lai YS, et al. 2012. <i>Proc. Natl. Acad. Sci. USA.</i> 109:3772.4. Jeong CH, et al. 2010. <i>Stem Cells</i> 28:2141.5. Xiang R, et al. 2011. <i>Br. J. Cancer</i> 104:1410.6. Card DA, et al. 2008. <i>Mol. Cell Biol.</i> 28:6426.
Gene ID	6657


Related Protocols

[Immunocytochemistry Staining Protocol](#)


Other Formats

Purified anti-SOX2, PE anti-SOX2, Alexa Fluor® 594 anti-SOX2, Alexa Fluor® 488 anti-SOX2, Alexa Fluor® 647 anti-SOX2, Pacific Blue™ anti-SOX2, Brilliant Violet 421™ anti-SOX2

Product Data



NCCIT cells were fixed with 1% paraformaldehyde (PFA) for 10 minutes at 37°C, permeabilized with 0.5% Triton X-100 at room temperature for 10 minutes, blocked for 30 minutes at room temperature with 5% fetal bovine serum (FBS), then stained with anti-human SOX2 (clone 14A6A34) Alexa Fluor® 647 (red) overnight at 4°C. F-actin was stained with Phalloidin Alexa Fluor® 488 (green) and nuclei were counterstained with DAPI (blue).



Human embryonic carcinoma NCCIT cells were fixed and permeabilized with FOXP3 Fix/Perm Buffer Set, and then stained with SOX2 (clone 14A6A34) AF647 (filled histogram) or mouse IgG1, κ AF647 isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587