

APC anti-human IL-2 Antibody

Catalog# / Size	500309 / 25 tests 500311 / 50 µg 500310 / 100 tests
Clone	MQ1-17H12
Regulatory Status	RUO
Other Names	Interleukin-2, T cell growth factor (TCGF), Eosinophil differentiation factor (EDF), Killer cell helper factor (KHF), Macrophage-activating factor for cytotoxicity I (MAF-C I), Thymocyte differentiation factor (TDF)
Isotype	Rat IgG2a, κ
Description	IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells, promoting proliferation and maturation. Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK cells, LAK cells, monocytes, and oligodendrocytes.

Product Details

Verified Reactivity	Human
Reported Reactivity	Cat, Chimpanzee, Baboon, Cynomolgus, Rhesus, Sooty Mangabey
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	<i>E. coli</i> - expressed recombinant human IL-2
Formulation	test sizes: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA). µg size: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions.
Concentration	µg sizes: 0.2 mg/mL test sizes: lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.
Excitation Laser	Red Laser (633 nm)
Application Notes	ELISA or ELISPOT Capture^{2,3}: The purified MQ1-17H12 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the Biotin anti-human IL-2 antibody (Cat. No. 517605) as the detecting antibody. The Ultra-LEAF™ purified antibody is suggested for ELISPOT capture. For ELISPOT capture applications, a concentration range of 4.0 - 8.0 µg/mL is recommended. Additional reported applications (for the relevant formats) include: immunoprecipitation ² , immunohistochemical staining of paraformaldehyde-fixed, saponin-treated frozen tissue sections ^{1,4-6,8} , neutralization ¹³ , and immunocytochemistry. Note: For testing human IL-2 in serum or plasma, BioLegend's LEGEND MAX™ Kit (Cat. No. 431807) is specially developed and recommended. Clone MQ1-17H12 cross-reacts to Cat ¹⁵

Application References

(PubMed link indicates BioLegend citation)

1. Andersson J, *et al.* 1994. *Immunology* 83:16. (IHC)
2. Abrams J, *et al.* 1992. *Immunol. Rev.* 127:5. (IP)
3. Abrams JS. 1995. *Curr. Prot. Immunol.* Unit 6.20.
4. Fernandez V, *et al.* 1994. *Eur. J. Immunol.* 24:1808. (IHC)
5. Skansen-Saphir U, *et al.* 1994. *Eur. J. Immunol.* 24:916. (IHC)
6. Andersson U, *et al.* *Detection and Quantification of Gene Expression*. New York:Springer-Verlag. (IHC)
7. Prussin C, *et al.* 1995. *J. Immunol. Methods.* 188:117.
8. Raqib R, *et al.* 2002. *Infect. Immun.* 70:3199. (IHC)
9. Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113. [PubMed](#)
10. Colleton BA, *et al.* 2009. *J Virol.* 83:6288. [PubMed](#)
11. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
12. Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
13. Yeap SK, *et al.* 2013. *BMC Complement Altern. Med.* 13:145. (Neut)
14. Wu Z, *et al.* 2015. *J Virol.* 89:6435. [PubMed](#)
15. Maksaarekul S, *et al.* 2009. *Vaccine.* 28:3754 (FC) [PubMed](#)

Product Citations

1. Colleton B, *et al.* 2009. *J Virol.* 83:6288. [PubMed](#)
2. Hall VG, *et al.* 2022. *Nat Immunol.* 23:380. [PubMed](#)
3. D, *et al.* 2016. *Tuberculosis.* 95: 470-475. [PubMed](#)
4. Bauer J, *et al.* 2022. *Nat Commun.* 13:6401. [PubMed](#)
5. Zhou X, *et al.* 2021. *J Clin Transl Hepatol.* 9:283. [PubMed](#)
6. Qiu XM, *et al.* 2020. *Reproduction.* 251:159. [PubMed](#)
7. Cohen KW, *et al.* 2021. *Cell Reports Medicine.* 2(7):100354. [PubMed](#)
8. Zhang Z, *et al.* 2021. *Acta Pharm Sin B.* 11:1965. [PubMed](#)
9. Priyanto H, *et al.* 2021. *J Clin Tuberc Other Mycobact Dis.* 22:100214. [PubMed](#)
10. Heitmann JS, *et al.* 2021. *Nature.* Online ahead of print. [PubMed](#)
11. Zhang B, *et al.* 2021. *Nat Biomed Eng.* 5:1288. [PubMed](#)
12. Keeton R, *et al.* 2021. *Cell Host Microbe.* 29:1611. [PubMed](#)
13. Fuery A, *et al.* 2015. *PLoS One.* 10: 0133126. [PubMed](#)
14. Whitfield SJC, *et al.* 2017. *J Immunol.* 198:3989. [PubMed](#)
15. Waight JD, *et al.* 2018. *Cancer Cell.* 33:1033. [PubMed](#)
16. Motozono C, *et al.* 2021. *Cell Host Microbe.* . [PubMed](#)
17. Cassioli C, *et al.* 2021. *Front Cell Dev Biol.* 9:673446. [PubMed](#)
18. Wang Y, *et al.* 2021. *Cell Reports.* 36(6):109516. [PubMed](#)
19. Ferreira VH, *et al.* 2022. *Nat Commun.* 13:4489. [PubMed](#)
20. Rochman Y, *et al.* 2015. *PLoS One.* 10:122198. [PubMed](#)
21. Kuranda K, *et al.* 2018. *J Clin Invest.* 128:5267. [PubMed](#)
22. Faissner S, *et al.* 2022. *Front Immunol.* 13:980526. [PubMed](#)

RRID

AB_315096 (BioLegend Cat. No. 500309)
AB_315098 (BioLegend Cat. No. 500311)
AB_315097 (BioLegend Cat. No. 500310)

Antigen Details

Structure	Cytokine; 15.4 kD (Mammalian)
Bioactivity	Proliferation of T lymphocytes, B cells, anti-inflammatory, hematopoiesis, tumor surveillance
Cell Sources	T cells
Cell Targets	T cells, B cells, NK cells, LAK cells, monocytes, macrophages, oligodendrocytes
Receptors	High affinity heterotrimer of IL-2R α / β / γ , intermediate affinity homodimer IL-2R α (CD25; p55; Tac) and heterodimer IL-2R β (CD122)/ γ ; γ -subunit (CD132) in common with IL-4R, IL-7R, IL-13R, IL-15R
Cell Type	Tregs
Biology Area	Cell Biology, Immunology, Neuroinflammation, Neuroscience
Molecular Family	Cytokines/Chemokines
Antigen References	<ol style="list-style-type: none">1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook</i>. Academic Press, San Diego.2. Taniguchi T, <i>et al.</i> 1993. <i>Cell</i> 73:5.3. Nistico G. 1993. <i>Prog. Neurobiol.</i> 40:463.4. Waldmann T, <i>et al.</i> 1993. <i>Ann. NY Acad. Sci.</i> 685:603.
Regulation	Upregulated by NFAT; downregulated by TCF-8 and CIF (colostrums inhibitory factor)
Gene ID	3558

Related Protocols


[Surface and Intracellular Cytokine Staining for Flow Cytometry - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-human IL-2, FITC anti-human IL-2, PE anti-human IL-2, Purified anti-human IL-2, Alexa Fluor® 488 anti-human IL-2, Alexa Fluor® 647 anti-human IL-2, Alexa Fluor® 700 anti-human IL-2, PerCP/Cyanine5.5 anti-human IL-2, Pacific Blue™ anti-human IL-2, PE/Cyanine7 anti-human IL-2, Brilliant Violet 421™ anti-human IL-2, Brilliant Violet 605™ anti-human IL-2, Brilliant Violet 650™ anti-human IL-2, Brilliant Violet 510™ anti-human IL-2, Brilliant Violet 711™ anti-human IL-2, APC/Cyanine7 anti-human IL-2, Purified anti-human IL-2 (Maxpar® Ready), PE/Dazzle™ 594 anti-human IL-2, Brilliant Violet 785™ anti-human IL-2, PerCP anti-human IL-2, APC/Fire™ 750 anti-human IL-2, Ultra-LEAF™ Purified anti-human IL-2

Product Data



PMA + ionomycin-stimulated (6 hours)
human peripheral blood lymphocytes
intracellular stained with MQ1-17H12
APC and CD3 (UCHT1) PE

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587