

Biotin anti-mouse CD3 Antibody

Catalog# / Size	100243 / 50 µg 100244 / 500 µg
Clone	17A2
Regulatory Status	RUO
Other Names	T cell antigen receptor complex, T3
Isotype	Rat IgG2b, κ
Description	CD3, also known as T3, is a member of the Ig superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3 is composed of CD3ε, δ, γ and ζ chains. It forms a TCR complex by associating with TCR α/β or γ/δ chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and antigen recognition by binding the peptide/MHC antigen complex

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	γδTCR-positive T-T hybridoma D1
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with biotin under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	FC - Quality tested IHC-F - Verified
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤0.125 µg per million cells in 100 µl volume. For immunohistochemistry, a concentration range of 5.0 - 10 µg/mL is suggested. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes	Additional reported application (for relevant formats) include: spatial biology (IBEX) ^{1,2} .
Application References	1. Radtke AJ, <i>et al.</i> 2020. <i>Proc Natl Acad Sci U S A.</i> 117:33455-65. (SB) PubMed 2. Radtke AJ, <i>et al.</i> 2022. <i>Nat Protoc.</i> 17:378-401. (SB) PubMed
(PubMed link indicates BioLegend citation)	
Product Citations	1. Chappaz S, <i>et al.</i> 2021. <i>Cell Reports.</i> 36(3):109430. PubMed 2. Thierry GR, <i>et al.</i> 2018. <i>J Exp Med.</i> 215:2972. PubMed 3. Shao B, <i>et al.</i> 2018. <i>Mol Med Rep.</i> 18:920. PubMed 4. Barry KC, <i>et al.</i> 2018. <i>Nat Med.</i> 24:1178. PubMed 5. Blecher-Gonen R, <i>et al.</i> 2019. <i>Cell Syst.</i> 8:109. PubMed 6. Yu VWC <i>et al.</i> 2016. <i>Cell.</i> 167(5):1310-1322. PubMed 7. Zong L, <i>et al.</i> 2021. <i>NPJ Aging Mech Dis.</i> 7:25. PubMed 8. Joachim R, Suber F, and Kobzik L 2017. <i>Sci Rep.</i> . 10.1038/s41598-017-16743-1. PubMed 9. Zhao M, <i>et al.</i> 2019. <i>Cell Rep.</i> 26:652. PubMed 10. Dey A <i>et al.</i> 2019. <i>The EMBO journal.</i> 38(7) pii: e100293. PubMed 11. Niss K, <i>et al.</i> 2020. <i>Cell Reports.</i> 31(11):107763. PubMed 12. Wang Y, <i>et al.</i> 2021. <i>Signal Transduct Target Ther.</i> 6:409. PubMed 13. Shen Z, <i>et al.</i> 2021. <i>iScience.</i> 24:103014. PubMed

14. Yu WWC, *et al.* 2017. *Cell*. 168:944. [PubMed](#)
15. Chen X *et al.* 2017. *Cell stem cell*. 21(6):747-760 . [PubMed](#)
16. Dong S, *et al.* 2021. *Nature*. 591:117. [PubMed](#)
17. Comazetto S *et al.* 2018. *Cell stem cell*. 24(3):477-486 . [PubMed](#)
18. Lebert-Ghali C, *et al.* 2016. *Blood*. 127: 87-90. [PubMed](#)
19. Martínez-López M *et al.* 2019. *Immunity*. 50(2):446-461 . [PubMed](#)
20. Kyburz A, *et al.* 2019. *J Allergy Clin Immunol*. 143:1496. [PubMed](#)
21. Gamrekelashvili J, *et al.* 2021. *Bio Protoc*. 11:e4007. [PubMed](#)
22. Bost P, *et al.* 2020. *Cell*. 181(7):1475-1488. [PubMed](#)
23. Zeis P, *et al.* 2020. *Immunity*. 53:775. [PubMed](#)
24. Miyauchi K, *et al.* 2021. *Nat Commun*. 12:3789. [PubMed](#)
25. Sebina I, *et al.* 2016. *PLoS Pathog*. 12:e1005999. [PubMed](#)
26. Lefkopoulos S, *et al.* 2020. *Immunity*. 53(5):934-951.e9. [PubMed](#)
27. Zhu J *et al.* 2017. *Cell stem cell*. 20(3):374-384 . [PubMed](#)
28. Säwen P *et al.* 2018. *eLife*. 7 pii: e41258. [PubMed](#)
29. Lee A, *et al.* 2022. *Nat Commun*. 13:549. [PubMed](#)

RRID AB_2563946 (BioLegend Cat. No. 100243)
 AB_2563947 (BioLegend Cat. No. 100244)

Antigen Details

Structure	Ig superfamily, CD3/TCR, 20 kD
Distribution	Thymocytes (differentiation dependent), mature T cells, NK-T cells
Function	Antigen recognition, TCR signal transduction, T cell activation
Ligand/Receptor	Peptide antigen/MHC-complex
Antigen References	<ol style="list-style-type: none"> 1. Barclay A, <i>et al.</i> 1997. <i>The Leukocyte Antigen FactsBook</i> Academic Press. 2. Davis MM. 1990. <i>Annu. Rev. Biochem.</i> 59:475. 3. Weiss A, <i>et al.</i> 1994. <i>Cell</i> 76:263.
Gene ID	12502

Related Protocols



[Immunohistochemistry Protocol for Frozen Sections](#)

[Cell Surface Flow Cytometry Staining Protocol](#)


Other Formats

FITC anti-mouse CD3, PE anti-mouse CD3, Purified anti-mouse CD3, Alexa Fluor® 647 anti-mouse CD3, Alexa Fluor® 488 anti-mouse CD3, Pacific Blue™ anti-mouse CD3, Alexa Fluor® 700 anti-mouse CD3, PerCP/Cyanine5.5 anti-mouse CD3, PE/Cyanine7 anti-mouse CD3, APC/Cyanine7 anti-mouse CD3, Brilliant Violet 421™ anti-mouse CD3, Brilliant Violet 570™ anti-mouse CD3, Brilliant Violet 650™ anti-mouse CD3, Brilliant Violet 785™ anti-mouse CD3, Brilliant Violet 510™ anti-mouse CD3, APC anti-mouse CD3, Ultra-LEAF™ Purified anti-mouse CD3, Brilliant Violet 605™ anti-mouse CD3, Alexa Fluor® 594 anti-mouse CD3, Brilliant Violet 711™ anti-mouse CD3, Biotin anti-mouse CD3, PE/Dazzle™ 594 anti-mouse CD3, APC/Fire™ 750 anti-mouse CD3, Brilliant Violet 750™ anti-mouse CD3, TotalSeq™-A0182 anti-mouse CD3, TotalSeq™-B0182 anti-mouse CD3, Spark Blue™ 550 anti-mouse CD3, Spark NIR™ 685 anti-mouse CD3, TotalSeq™-C0182 anti-mouse CD3, APC/Fire™ 810 anti-mouse CD3, PE/Fire™ 640 anti-mouse CD3, Spark YG™ 570 anti-mouse CD3, PE/Fire™ 700 anti-mouse CD3, PE/Cyanine5 anti-mouse CD3, Spark Blue™ 574 anti-mouse CD3 Antibody, Spark Violet™ 423 anti-mouse CD3, PE/Fire™ 810 anti-mouse CD3, Spark Red™ 718 anti-mouse CD3

Product Data



C57BL/6 splenocytes were stained with CD19 APC and biotinylated CD3 (clone 17A2) (top) or biotinylated rat IgG2b, κ isotype control (bottom), followed by SAV-PE.



C57BL/6 mouse frozen spleen section was fixed with 4% paraformaldehyde (PFA) for 10 minutes at room temperature and blocked with 5% FBS for 30 minutes at room temperature. Then the section was stained with 10 μ g/ml of biotin anti-mouse CD3 (clone 17A2), Alexa Fluor[®] 594 anti-mouse B220 (clone RA3-6B2) (blue) and Alexa Fluor[®] 488 anti-mouse CD29 (clone HM β 1-1) (green) overnight at 4 $^{\circ}$ C, followed by 2.5 μ g/ml of Spark YG[™] 570 Streptavidin (red) for 2 hours at room temperature. The image was captured by 10X objective.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587