

FITC anti-human CD3 Antibody

Catalog# / Size	344803 / 25 tests 344804 / 100 tests
Clone	SK7
Regulatory Status	RUO
Workshop	HCDM listed
Other Names	T3, CD3ε
Isotype	Mouse IgG1, κ
Description	CD3ε is a 20 kD chain of the CD3/T-cell receptor (TCR) complex, which is composed of two CD3ε, one CD3γ, one CD3δ, one CD3ζ (CD247), and a T-cell receptor (α/β or γ/δ) heterodimer. It is found on all mature T cells, NK T cells, and some thymocytes. CD3, also known as T3, is a member of the immunoglobulin superfamily that plays a role in antigen recognition, signal transduction, and T cell activation.

Product Details

Verified Reactivity	Human
Reported Reactivity	Chimpanzee
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.
Excitation Laser	Blue Laser (488 nm)
Application Notes	Additional reported application (for the relevant formats) include: immunohistochemical staining of frozen tissue sections ^{4,5,8} , immunofluorescent staining ⁶ , and Western blotting ³ .
Application References	<ol style="list-style-type: none"> 1. Kan EA, <i>et al.</i> 1983. <i>J. Immunol.</i> 131:536. 2. Wood GS, <i>et al.</i> 1985. <i>Am. J. Pathol.</i> 120:371. 3. Van Dongen JJM, <i>et al.</i> 1988. <i>Blood</i> 71:603. (WB) 4. Haringman JJ, <i>et al.</i> 2005. <i>Arthritis Res. Ther.</i> 7:R862. (IHC) 5. Carbone A, <i>et al.</i> 1999. <i>Blood</i> 93:2319. (IHC) 6. Goval JJ, <i>et al.</i> 2006. <i>J. Histochem. Cytochem.</i> 54:75. (IF) 7. Rutjens E, <i>et al.</i> 2007. <i>J. Immunol.</i> 178:1702. 8. Kap Y, <i>et al.</i> 2009. <i>J. Histochem. Cytochem.</i> 57:1159. (IHC) 9. Yoshino N, <i>et al.</i> 2000. <i>Exp. Anim. (Tokyo)</i> 49:97. (FC)
Product Citations	<ol style="list-style-type: none"> 1. Chan S, <i>et al.</i> 2022. <i>Nat Cancer.</i> . PubMed 2. Chan JA, <i>et al.</i> 2022. <i>Nat Commun.</i> 13:4159. PubMed 3. Penkava F, <i>et al.</i> 2020. <i>Nat Commun.</i> 3.76875. PubMed

4. Montes de Oca M, *et al.* 2016. *Cell Rep.* 17:399-412. [PubMed](#)
5. Cao M, *et al.* 2020. *Aging Dis.* 1.479861111. [PubMed](#)
6. Hagan T, *et al.* 2020. *Cell.* 178(6):1313-1328.e13. [PubMed](#)
7. Björklund &, *et al.* 2016. *Nat Immunol.* 17:451-460. [PubMed](#)
8. Weaver JD, *et al.* 2022. *Oncoimmunology.* 11:2141007. [PubMed](#)
9. Thiel BA, *et al.* 2021. *PLoS One.* 16:e0249477. [PubMed](#)
10. Lu J, *et al.* 2021. *Front Immunol.* 12:732612. [PubMed](#)
11. Mudd PA, *et al.* 2022. *Cell.* 185:603. [PubMed](#)
12. Lima J, *et al.* 2016. *Reprod Sci.* 10.1177/1933719116653680. [PubMed](#)
13. Prota G, *et al.* 2020. *Cancer Immunol Res.* 8:685. [PubMed](#)
14. Chheda ZS, *et al.* 2018. *J Exp Med.* 9:19490. [PubMed](#)
15. Nagree MS, *et al.* 2022. *EMBO Mol Med.* 14:e14297. [PubMed](#)
16. Roudko V, *et al.* 2020. *Cell.* 183(6):1634-1649.e17. [PubMed](#)
17. Cimen Bozkus C, *et al.* 2021. *STAR Protoc.* 2:100758. [PubMed](#)
18. Qing G, *et al.* 2021. *Front Cell Dev Biol.* 9:761300. [PubMed](#)
19. Carter JA, *et al.* 2020. *Cell Systems.* 9(5):475-482.e4. [PubMed](#)

RRID AB_2043992 (BioLegend Cat. No. 344803)
 AB_2043993 (BioLegend Cat. No. 344804)

Antigen Details

Structure	Ig superfamily, with the subunits of CD3 γ , CD3 δ , CD3 ζ , (CD247) and TCR (α/β or γ/δ) forms CD3/TCR complex, 20 kD
Distribution	Mature T and NK T cells, during thymocyte differentiation
Function	Antigen recognition, signal transduction, T cell activation
Ligand/Receptor	Peptide antigen bound to MHC
Cell Type	NKT cells, T cells, Tregs
Biology Area	Immunology, Innate Immunity
Molecular Family	CD Molecules, TCRs
Antigen References	<ol style="list-style-type: none"> 1. Barclay N, <i>et al.</i> 1993. <i>The Leucocyte FactsBook.</i> Academic Press. San Diego. 2. Beverly P, <i>et al.</i> 1981. <i>Eur. J. Immunol.</i> 11:329. 3. Lanier L, <i>et al.</i> 1986. <i>J. Immunol.</i> 137:2501.
Gene ID	916

Related Protocols


[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC/Fire™ 750 anti-human CD3, Biotin anti-human CD3, Purified anti-human CD3, FITC anti-human CD3, PE anti-human CD3, Alexa Fluor® 488 anti-human CD3, APC anti-human CD3, PerCP/Cyanine5.5 anti-human CD3, PerCP anti-human CD3, PE/Cyanine7 anti-human CD3, APC/Cyanine7 anti-human CD3, Alexa Fluor® 700 anti-human CD3, Pacific Blue™ anti-human CD3, Alexa Fluor® 647 anti-human CD3, Brilliant Violet 510™ anti-human CD3, Brilliant Violet 421™ anti-human CD3, Brilliant Violet 605™ anti-human CD3, Brilliant Violet 711™ anti-human CD3, Brilliant Violet 785™ anti-human CD3, PE/Dazzle™ 594 anti-human CD3, Brilliant Violet 750™ anti-human CD3, TotalSeq™-A0049 anti-human CD3, TotalSeq™-C0049 anti-human CD3, Spark Blue™ 550 anti-human CD3, TotalSeq™-B0049 anti-human CD3, Alexa Fluor® 660 anti-human CD3, APC/Fire™ 810 anti-human CD3, Spark NIR™ 685 anti-human CD3, PE/Fire™ 640 anti-human CD3, PE/Fire™ 700 anti-human CD3, GMP FITC anti-human CD3, PE/Cyanine5 anti-human CD3 Antibody, GMP PE anti-human CD3, GMP APC anti-human CD3, GMP PerCP/Cyanine5.5 anti-human CD3, Spark YG™ 593 anti-human CD3, GMP PerCP anti-human CD3, Spark Violet™ 500 anti-human CD3

Product Data

Human peripheral blood lymphocytes
stained with SK7 FITC.



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587