

## FITC anti-mouse/human CD15 (SSEA-1) Antibody

<b>Catalog# / Size</b>	125611 / 25 tests 125612 / 100 tests
<b>Clone</b>	MC-480
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	Stage-Specific Embryonic Antigen 1, X-hapten, Lewis X, 3-FAL, CD15, SSEA1, SSEA-1
<b>Isotype</b>	Mouse IgM, κ
<b>Description</b>	The MC-480 antibody reacts with mouse and human Stage-Specific Embryonic Antigen-1 (SSEA-1). It is a Lewis blood group related carbohydrate antigen, also known as X-hapten, Lewis X, 3-FAL, 3-fucosyl-N-acetylactosamine, or CD15. The expression pattern of SSEA-1 antigen is different in humans than in mice. In mice, SSEA-1 is expressed on embryonic stem cells (ES), embryonal carcinoma cells (EC), 8-cell to blastocyst embryos, and a subset of embryonic inner cell mass. The expression on murine ES cells is decreased upon differentiation. In humans, however, SSEA-1 is not found on undifferentiated ES cells, but its expression is upregulated along with differentiation. CD15 is highly expressed on adult human granulocytes. It has been reported that SSEA-1 plays a role in cell adhesion and regulation of cell differentiation.

### Product Details

---

<b>Verified Reactivity</b>	Human, Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	Mouse F9 Teratocarcinoma Stem Cells (X-irradiated)
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.
<b>Excitation Laser</b>	Blue Laser (488 nm)
<b>Application Notes</b>	Additional reported applications (for the relevant formats) include: immunoprecipitation <sup>1</sup> , Western blotting <sup>1</sup> , and immunohistochemistry <sup>1</sup> of acetone-fixed frozen tissue sections and formalin-fixed paraffin-embedded sections.
<b>Application References</b> (PubMed link indicates BioLegend citation)	1. Solter D and Knowles BB. 1978. <i>Proc. Natl. Acad. Sci. USA.</i> 75:5565. (IHC, IP, WB)
<b>Product Citations</b>	1. Hamada N, <i>et al.</i> 2020. PLoS Genet. 16:e1008676. <a href="#">PubMed</a> 2. Medine C, <i>et al.</i> 2011. J Vis Exp. 56: 2969. <a href="#">PubMed</a> 3. Dumm RE, <i>et al.</i> 2019. Nat Commun. 10:779. <a href="#">PubMed</a>
<b>RRID</b>	AB_2561706 (BioLegend Cat. No. 125611)

## Antigen Details

<b>Structure</b>	Carbohydrate epitope, Lewis blood group antigens
<b>Distribution</b>	Murine embryonic stem cells (ES), embryonal carcinoma cells, 8-cell to blastocyst embryos; human differentiated cells, not on undifferentiated ES cells
<b>Function</b>	Adhesion, differentiation
<b>Cell Type</b>	Embryonic Stem Cells, Neural Stem Cells
<b>Biology Area</b>	Cell Biology, Immunology, Neuroscience, Neuroscience Cell Markers, Stem Cells
<b>Molecular Family</b>	CD Molecules
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Solter D and Knowles BB. 1978. <i>Proc. Natl. Acad. Sci. USA.</i> 75:5565.</li> <li>2. Harris JF, <i>et al.</i> 1984. <i>J. Immunol.</i> 132:2502.</li> <li>3. Gooi HC, <i>et al.</i> 1981. <i>Nature</i> 292:156.</li> <li>4. Cui L, <i>et al.</i> 2004. <i>J. Histochem. Cytochem.</i> 52:1447.</li> </ol>
<b>Gene ID</b>	<a href="#">14345</a> <a href="#">2526</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

PE anti-mouse/human CD15 (SSEA-1), Biotin anti-mouse/human CD15 (SSEA-1), Purified anti-mouse/human CD15 (SSEA-1), Alexa Fluor® 647 anti-mouse/human CD15 (SSEA-1), Alexa Fluor® 488 anti-mouse/human CD15 (SSEA-1), FITC anti-mouse/human CD15 (SSEA-1), TotalSeq™-A0076 anti-mouse/human CD15 (SSEA-1), APC anti-mouse/human CD15 (SSEA-1), TotalSeq™-C0076 anti-mouse/human CD15 (SSEA-1), TotalSeq™-B0076 anti-mouse/human CD15 (SSEA-1)

## Product Data



F9 cells (mouse embryonic carcinoma cell line) were stained with anti-mouse/human SSEA-1 (clone MC-480, filled histogram) FITC or mouse IgM FITC isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
 Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587

