

## Go-ChIP-Grade™ Purified anti-STAT1 Phospho (Tyr701) Antibody

<b>Catalog# / Size</b>	666405 / 25 µg 666406 / 100 µg
<b>Clone</b>	A17012A
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	Signal Transduced Activator of Transcription 1, Transcription Factor ISGF-3 components p91/p84
<b>Isotype</b>	Mouse IgG1, κ
<b>Description</b>	STAT1, also known as signal transduction and activator of transcription 1, is a ubiquitously expressed cytoplasmic protein and is activated in response to cytokine signaling, including IFN-α, IFN-γ, EGF, PDGF, and IL-6. Upon activation, STAT1 is phosphorylated at Tyrosine 701 (Tyr701) by receptor-associated kinases, including JAK1, JAK2, and TYK2. This results in STAT1 dimerization and subsequent translocation to the nucleus, where it functions as a transcriptional activator. STAT1 is involved in IFN-mediated immune responses, and STAT1-deficient mice are highly sensitive to bacterial and viral infections.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Preparation</b>	The antibody was purified by affinity chromatography.
<b>Concentration</b>	0.5 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C.
<b>Application</b>	<a href="#">ChIP - Quality tested</a> <a href="#">WB, ICFC, ICC - Verified</a>
<b>Recommended Usage</b>	<p>Each lot of this antibody is quality control tested by ChIP Assay. The suggested dilution for ChIP application is 1:50 by volume. For Western blotting, the suggested use of this reagent is 0.1 - 2.0 µg per ml. For intracellular flow cytometric staining, the suggested use of this reagent is ≤ 0.5 µg per million cells in 100 µl volume. For immunocytochemistry, a concentration of 5.0 µg/ml is recommended. It is recommended that the reagent to be titrated for optimal performance before each experiment.</p> <p>For intracellular flow cytometry our <a href="#">True-Nuclear™ Transcription Factor Staining Protocol</a> is recommended.</p>
<b>Application Notes</b>	<p>Clone A17012A recognizes STAT1 phosphorylated at Tyrosine 701 (Tyr701).</p> <p>When using this clone for ICC, we recommend using methanol to permeabilize fixed cells.</p>
<b>RRID</b>	AB_2728501 (BioLegend Cat. No. 666405) AB_2728502 (BioLegend Cat. No. 666406)

### Antigen Details

---

<b>Structure</b>	Two isoforms of STAT1 exist as a result of alternative splicing. Isoform α is a 750 amino acid protein with a predicted molecular weight of 87 kD; the β isoform is a 712 amino acid protein with a predicted molecular weight of 83 kD.
<b>Distribution</b>	Ubiquitous tissue expression, nucleoplasmic-cytosolic distribution

<b>Function</b>	Immune response activation, cell signaling
<b>Interaction</b>	STAT1, STAT2, IRF9, ERBB4, JAK1, JAK2, TYK2, TNK1, SHP2
<b>Biology Area</b>	Cell Biology
<b>Molecular Family</b>	Nuclear Markers, Phospho-Proteins

#### Antigen References

1. Moretti S, et al. 2017. J. Biol. Chem. 292: 1785.
2. Wei J, et al. 2015. J. Immunol. 195: 2870.
3. Sung PS, et al. 2015. Proc. Natl. Acad. Sci. USA. 112: 10443
4. Ooi EL, et al. 2014. Proc. Natl. Acad. Sci. USA. 111: 1909.
5. Wu TR, et al. 2002. J. Biol. Chem. 277: 47572.
6. Horvath, et al. 1996. J. Virol. 70: 647.
7. Haque SJ, et al. 1995. J. Biol. Chem. 270: 25709.

**Gene ID** [6772](#)

## Related Protocols

[BioLegend's Tools for Chromatin Immunoprecipitation \(ChIP\) Assays - Video](#)

[Chromatin Immunoprecipitation \(ChIP\) Assay Protocol](#)

[Immunocytochemistry Staining Protocol](#)

[Western Blotting Protocol](#)


[Intracellular Staining With True-Phos™ Perm Buffer in Cell Suspensions Protocol](#)

## Other Formats


Purified anti-STAT1 Phospho (Tyr701), Go-ChIP-Grade™ Purified anti-STAT1 Phospho (Tyr701), PE anti-STAT1 Phospho (Tyr701), Alexa Fluor® 647 anti-STAT1 Phospho (Tyr701), PerCP/Cyanine5.5 anti-STAT1 Phospho (Tyr701)

## Product Data


### Anti-STAT1 Phospho (Tyr701) antibody (A17012A)




Chromatin Immunoprecipitations (ChIP) were performed with cross-linked chromatin samples from  $4 \times 10^6$  of HT1080 cells treated with IFN $\gamma$  for 30 minutes with either A) 1:50 dilution of Go-ChIP-Grade™ Purified anti-STAT1 Phospho (Tyr701) (Clone A17012A) or B) equal amount of Purified Mouse IgG1,  $\kappa$  Isotype Control Antibody (Clone MG1-45, Cat. No. 401401) by using Go-ChIP-Grade™ Protein G Enzymatic Kit (Cat. No. 699904). The enriched DNA was purified and quantified by real-time qPCR using primers targeting human TAP1 gene region and  $\alpha$ -Satellite repeats. The amount of immunoprecipitated DNA in each sample is represented as signal relative to total amount of input chromatin.




Total cell lysates (15  $\mu$ g protein) from serum-starved HeLa S3 cells treated without (Lane 1) or with (Lane 2) 10 ng/mL human IFN- $\alpha$ 2 (Cat. No. 592702) for 10 minutes were resolved by 8% Bis-Tris gel electrophoresis, transferred to nitrocellulose, and probed with 2  $\mu$ g/mL (1:250 dilution) purified anti-STAT1 Phospho (Tyr701) antibody (clone A17012A). Proteins were visualized by chemiluminescence detection using HRP goat anti-mouse-IgG (Cat. No. 405301) at a 1:3000 dilution. Equal STAT1 loading was confirmed using a pan anti-STAT1 antibody that recognizes both STAT1  $\alpha$  (87 kD) and  $\beta$  (83 kD) isoforms.




Total cell lysates (15  $\mu$ g protein) from serum-starved HeLa S3 cells treated without (Lane 1) or with (Lane 2) 10 ng/mL human IFN- $\alpha$ 2 (Cat. No. 592702) for 10 minutes were resolved by 8% Bis-Tris gel electrophoresis, transferred to nitrocellulose, and probed with the indicated concentration of purified anti-STAT1 Phospho (Tyr701) antibody (clone A17012A). Proteins were visualized by chemiluminescence detection using HRP goat anti-mouse-IgG (Cat. No. 405301) at a 1:3000 dilution. Equal protein loading was confirmed using Direct-Blot™ HRP anti- $\beta$ -Actin antibody (Cat. No. 643807) used at a 1:5000 dilution.



Human peripheral blood mononuclear cells were treated with (left), or without (right) Recombinant Human Interferon- $\gamma$  (Cat. No. 570204) for 15 minutes, fixed with Fixation Buffer (Cat. No. 420801), permeabilized with True-Phos™ Perm Buffer (Cat. No. 425401) then stained with CD4 (clone A161A1) APC and purified anti-STAT1 Phospho (Tyr701) (clone A17012A), followed by anti-mouse IgG PE. Data was gated on lymphocyte and monocyte populations.



Human peripheral blood monocytes were treated with (filled histogram), or without (open histogram) Recombinant Human Interferon- $\gamma$  (Cat. No. 570204) for 15 minutes, fixed with Fixation Buffer (Cat. No. 420801), permeabilized with True-Phos™ Perm Buffer (Cat. No. 425401) then stained with purified anti-STAT1 Phospho (Tyr701) (clone A17012A), followed by anti-mouse IgG PE.



Serum starved HeLa cells were untreated (panel A) or stimulated with 100 ng/mL IFN- $\alpha$ 2 (Cat. No. 592702, panel B) for 20 minutes, fixed with 4% paraformaldehyde for 10 minutes, permeabilized with ice-cold methanol for 10 minutes, and blocked with 5% FBS for 60 minutes. Cells were then intracellularly stained with a 1:100 dilution (5  $\mu$ g/mL) of purified anti-STAT1 Phospho (Tyr701) Antibody, clone A17012A, for two hours at room temperature, followed by incubation with Alexa Fluor® 594 goat anti-mouse IgG Antibody (Cat. No. 405326) at 2.0  $\mu$ g/mL. Nuclei were counterstained with DAPI, and the image was captured with a 60X objective.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587