

## KIRAVIA Blue 520™ anti-human CD141 (Thrombomodulin) Antibody

<b>Catalog# / Size</b>	344129 / 25 tests 344130 / 100 tests
<b>Clone</b>	M80
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	Thrombomodulin, TM, THRM, THBD, Fetomodulin, BDCA-3
<b>Isotype</b>	Mouse IgG1, κ
<b>Description</b>	CD141 is a 75 kD, single chain, type I membrane glycoprotein also known as thrombomodulin, TM, THRM, THBD, and fetomodulin. CD141 is an important cofactor in the protein C anticoagulant system. After binding to its ligand thrombin, CD141 activates protein C, which degrades clotting factors Va and VIIIa, and as a consequence the amount of thrombin is reduced. CD141 is expressed on macrophages, monocytes, a subpopulation of myeloid dendritic cells, vascular endothelial cells, and keratinocytes. Besides anti-coagulation function, CD141 is also involved in embryonic and atherosclerotic plaque development.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Reported Reactivity</b>	African Green, Baboon
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with KIRAVIA Blue 520™ under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	<p>Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a>. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.</p> <p>* KIRAVIA Blue 520™ has an excitation maximum of 495 nm, and a maximum emission of 520 nm.</p> <p>KIRAVIA Blue™ 520 is a trademark of Sony. This product is subject to proprietary rights of Sony and is made and sold under license from Sony Corporation. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.</p> <p>Sony and the Sony logo are registered trademarks of Sony Corporation.</p>
<b>Excitation Laser</b>	Blue Laser (488 nm)
<b>RRID</b>	AB_2892417 (BioLegend Cat. No. 344129) AB_2892417 (BioLegend Cat. No. 344130)

### Antigen Details

---

<b>Structure</b>	Single chain, type I membrane glycoprotein, 75 kD
------------------	---

<b>Distribution</b>	Macrophages, monocytes, subset of myeloid dendritic cells, vascular endothelial cells, keratinocytes
<b>Function</b>	After thrombin binding, CD141 activates protein C, which degrades clotting factors Va and VIIIa and reduces the amount of thrombin generated.
<b>Interaction</b>	Protein C, Thrombin-activatable fibrinolysis inhibitor (TAFI), Platelet factor 4 (PF4)
<b>Ligand/Receptor</b>	Thrombin
<b>Cell Type</b>	Dendritic cells, Endothelial cells, Macrophages, Monocytes
<b>Biology Area</b>	Immunology, Innate Immunity
<b>Molecular Family</b>	CD Molecules
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Suzuki K, <i>et al.</i> 1987. <i>EMBO J.</i> 6:1891.</li> <li>2. Esmon CT, <i>et al.</i> 1989. <i>J. Biol. Chem.</i> 264:4743.</li> <li>3. Delvaeye M, <i>et al.</i> 2009. <i>N. Engl. J. Med.</i> 361:345.</li> <li>4. Shi CS, <i>et al.</i> 2008. <i>Blood</i> 112:3661.</li> <li>5. Chen LC, <i>et al.</i> 2009. <i>J. Infect.</i> 58:368.</li> </ol>
<b>Gene ID</b>	<a href="#">7056</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

Purified anti-human CD141 (Thrombomodulin), PE anti-human CD141 (Thrombomodulin), APC anti-human CD141 (Thrombomodulin), Biotin anti-human CD141 (Thrombomodulin), PE/Cyanine7 anti-human CD141 (Thrombomodulin), PerCP/Cyanine5.5 anti-human CD141 (Thrombomodulin), Brilliant Violet 421™ anti-human CD141 (Thrombomodulin), Brilliant Violet 785™ anti-human CD141 (Thrombomodulin), Brilliant Violet 605™ anti-human CD141 (Thrombomodulin), PE/Dazzle™ 594 anti-human CD141 (Thrombomodulin), TotalSeq™-A0163 anti-human CD141 (Thrombomodulin), Alexa Fluor® 647 anti-human CD141 (Thrombomodulin), TotalSeq™-C0163 anti-human CD141 (Thrombomodulin), TotalSeq™-B0163 anti-human CD141 (Thrombomodulin), KIRAVIA Blue 520™ anti-human CD141 (Thrombomodulin), TotalSeq™-D0163 anti-human CD141 (Thrombomodulin), PE/Fire™ 810 anti-human CD141 (Thrombomodulin)

## Product Data



Human peripheral blood monocytes were stained with anti-human CD141 (Thrombomodulin) (clone M80) KIRAVIA Blue 520™ (filled histogram) or mouse IgG1, κ KIRAVIA Blue 520™ control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587