

## PE anti-mouse Blimp-1 Antibody

<b>Catalog# / Size</b>	150005 / 25 µg 150006 / 100 µg
<b>Clone</b>	5E7
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	PR domain zinc finger protein 1, B lymphocyte-induced maturation protein-1, PRDM1, BLIMP1, PRDI-BF1, ZNFPR1A1, PRDM-1, BLIMP1
<b>Isotype</b>	Rat IgG2a, κ
<b>Description</b>	Blimp-1, also known as PRDM1, is a 98 kD protein containing 5 Kruppel-type zinc finger domains. Blimp-1 represses the transcription factors BCL6 and c-Myc. It is the master regulator of terminal B cell differentiation and is also involved in the differentiation and homeostasis of T cells and natural killer (NK) cells.

### Product Details

---

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Rat
<b>Immunogen</b>	Amino acids 255-395 from mouse Blimp-1 fused with GST.
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions.
<b>Concentration</b>	0.2 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">ICFC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by intracellular flow cytometry using our <a href="#">True-Nuclear™ Transcription Factor Staining Protocol</a> . For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.
<b>Excitation Laser</b>	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
<b>Product Citations</b>	1. Xu Y, <i>et al.</i> 2022. Nat Commun. 13:980. <a href="#">PubMed</a> 2. Leclerc M, <i>et al.</i> 2019. Nat Commun. 10:3345. <a href="#">PubMed</a>
<b>RRID</b>	AB_2565991 (BioLegend Cat. No. 150005) AB_2565992 (BioLegend Cat. No. 150006)

### Antigen Details

---

<b>Structure</b>	Five Kruppel-type zinc finger domains, 98 kD.
<b>Distribution</b>	Plasma cells, CD4 <sup>+</sup> and CD8 <sup>+</sup> effector/memory T cells, and natural killer cells.
<b>Function</b>	Terminal differentiation of B cells, effector/memory T cells, role in natural killer cells (NK) and T cell homeostasis, and repressor of BCL6 and c-Myc.

<b>Cell Type</b>	B cells, NK cells, Plasma cells, T cells
<b>Biology Area</b>	Cell Biology, Immunology, Transcription Factors
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Nakaki F, <i>et al.</i> 2013. <i>Nature</i> 501:222.</li> <li>2. Crotty S, <i>et al.</i> 2010. <i>Nat. Immunol.</i> 11:114.</li> <li>3. Zhao WL, <i>et al.</i> 2008. <i>Blood</i> 111:3867.</li> <li>4. Climmino L, <i>et al.</i> 2008. <i>J. Immunol.</i> 181:2338.</li> <li>5. Martins, G and Clarne, K. 2008. <i>Annu. Rev. Immunol.</i> 26:133.</li> </ol>
<b>Gene ID</b>	<a href="#">12142</a>

## Related Protocols


[Surface and Intracellular Cytokine Staining for Flow Cytometry - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)

## Other Formats

Alexa Fluor® 647 anti-mouse Blimp-1, PE anti-mouse Blimp-1, APC anti-mouse Blimp-1

## Product Data



C57BL/6 splenocytes were cultured for 4 days in the presence of LPS, stained with CD45R/B220 APC, fixed and permeabilized using True-Nuclear™ Transcription Factor Buffer, then stained with Blimp-1 (clone 5E7) PE (left) or Rat IgG2a, κ PE Isotype control (right)

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587