

PE anti-mouse CD28 Antibody

Catalog# / Size	102105 / 50 µg 102106 / 200 µg
Clone	37.51
Regulatory Status	RUO
Other Names	Tp44, T44
Isotype	Syrian Hamster IgG
Description	CD28 is a 44 kD glycoprotein, also known as Tp44 or T44. It is a member of the Ig superfamily, expressed on thymocytes, most peripheral T cells, and NK cells. In association with CD80 (B7-1) and CD86 (B7-2), CD28 acts as the second signal for T and NK cell activation and proliferation. The 37.51 antibody has been reported to augment <i>in vitro</i> T cell proliferation and cytokine production, and promote CTL development.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Syrian Hamster
Immunogen	C57BL/6 mouse T-cell lymphoma EL-4
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions.
Concentration	0.2 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤0.5 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	Additional reported applications (for the relevant formats) include: immunoprecipitation ¹ , <i>in vitro</i> costimulation of T and NK cells ¹ , <i>in vitro</i> blocking of allogeneic mixed leukocyte response and inhibition of MHC-unrestricted CTL cytotoxicity ^{3,4} , <i>in vitro</i> induction of thymocyte differentiation ^{2,5-9,11} , and immunohistochemical staining of acetone-fixed frozen sections. For <i>in vivo</i> studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) (Cat. No. 102116).
Application References	<ol style="list-style-type: none"> Gross JA, <i>et al.</i> 1992. <i>J. Immunol.</i> 149:380. (IP, Costim) Cibotti R, <i>et al.</i> 1997. <i>Immunity</i> 6:245. (Costim) Masten BJ, <i>et al.</i> 1997. <i>Am. J. Respir. Cell Mol. Biol.</i> 16:335. (Block) Nishio M, <i>et al.</i> 1996. <i>J. Immunol.</i> 157:4347. (Block) Zhang N and He Y-W, 2005. <i>J. Exp. Med.</i> 202:395. (Costim) Terrazas LI, <i>et al.</i> 2005. <i>Intl. J. Parasitology.</i> 35:1349. (Costim) Perchonock CE, <i>et al.</i> 2006. <i>Mol Cell Biol.</i> 26(16):6005. (Costim) Wang W, <i>et al.</i> 2007. <i>J. Immunol.</i> 178:4885. (Costim) Pua HH, <i>et al.</i> 2007. <i>J. Exp. Med.</i> 204:25. (Costim) Perchonock CE, <i>et al.</i> 2007. <i>J. Immunol.</i> 179:1768. Barbi J, <i>et al.</i> 2007. <i>Blood</i> 110:2215.
(PubMed link indicates BioLegend citation)	

12. Milpied P, *et al.* 2011. *Blood* 118:2993. [PubMed](#)
13. Cunningham NR, *et al.* 2011. *Int Immunol.* 23:693. [PubMed](#)
14. Crispin JC, *et al.* 2012. *J. Immunol.* 188:3567. [PubMed](#)
15. Li CR, *et al.* 2014. *J Immunol.* 192:1425. [PubMed](#)
16. Blankenhaus B, *et al.* 2014. *PLoS Pathog.* 10:1003913. [PubMed](#)

Product Citations

1. Qi Z, *et al.* 2022. *Nat Commun.* 13:182. [PubMed](#)
2. Xie A *et al.* 2017. *Endocrinology.* 158(10):3140-3151. [PubMed](#)
3. Paiva RA, *et al.* 2021. *Cell Reports.* 35(2):108967. [PubMed](#)
4. Shissler SC, *et al.* 2020. *Sci Rep.* 10:8218. [PubMed](#)
5. Martina M, *et al.* 2016. *J Am Soc Nephrol.* 27: 1113-1123. [PubMed](#)
6. Hu J, *et al.* 2022. *J Immunother Cancer.* 10:. [PubMed](#)
7. Sasayama S, *et al.* 2018. *Int J Mol Sci.* 19: [PubMed](#)
8. Timilshina M, *et al.* 2020. *Cell Reports.* 27(10):2948-2961.e7.. [PubMed](#)
9. Li X, *et al.* 2021. *Front Cell Dev Biol.* 9:647713. [PubMed](#)

RRID

AB_312870 (BioLegend Cat. No. 102105)
 AB_312871 (BioLegend Cat. No. 102106)

Disclaimer

This product may be used for research purposes only. It is not licensed for resale and may only be used by the buyer. This product may not be used and is not licensed for clinical assays where the results of such assays are provided as a diagnostic service. If a diagnostic or therapeutic use is anticipated then a license must be requested from the University of California. The availability of such diagnostic and therapeutic use license(s) cannot be guaranteed from the University of California.

Antigen Details

Structure	Ig superfamily, 44 kD
Distribution	Thymocytes, CD4 ⁺ , CD8 ⁺ peripheral T cells, NK cells
Function	Costimulates T and NK cells
Ligand/Receptor	CD80 (B7-1), CD86 (B7-2)
Cell Type	NK cells, T cells, Thymocytes, Tregs
Biology Area	Costimulatory Molecules, Immunology
Molecular Family	CD Molecules
Antigen References	<ol style="list-style-type: none"> 1. Barclay AN, <i>et al.</i> 1997. <i>The Leukocyte Antigen FactsBook</i> Academic Press. 2. Lenschow DJ, <i>et al.</i> 1996. <i>Annu. Rev. Immunol.</i> 14:233. 3. Gross JA, <i>et al.</i> 1992. <i>J. Immunol.</i> 149:380.
Gene ID	12487

Related Protocols


[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-mouse CD28, Biotin anti-mouse CD28, PE anti-mouse CD28, PE/Cyanine5 anti-mouse CD28, Purified anti-mouse CD28, PerCP/Cyanine5.5 anti-mouse CD28, Ultra-LEAF™ Purified anti-mouse CD28, Purified anti-mouse CD28 (Maxpar® Ready), PE/Cyanine7 anti-mouse CD28, PE/Dazzle™ 594 anti-mouse CD28, Brilliant Violet 421™ anti-mouse CD28, TotalSeq™-C0204 anti-mouse CD28, KIRAVIA Blue 520™ anti-mouse CD28

Product Data

C57BL/6 splenocytes stained with 37.51 PE and CD3 FITC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587