

PE/Cyanine7 anti-human CD141 (Thrombomodulin) Antibody

Catalog# / Size	344109 / 25 tests 344110 / 100 tests
Clone	M80
Regulatory Status	RUO
Other Names	Thrombomodulin, TM, THRM, THBD, Fetomodulin, BDCA-3
Isotype	Mouse IgG1, κ
Description	CD141 is a 75 kD, single chain, type I membrane glycoprotein also known as thrombomodulin, TM, THRM, THBD, and fetomodulin. CD141 is an important cofactor in the protein C anticoagulant system. After binding to its ligand thrombin, CD141 activates protein C, which degrades clotting factors Va and VIIIa, and as a consequence the amount of thrombin is reduced. CD141 is expressed on macrophages, monocytes, a subpopulation of myeloid dendritic cells, vascular endothelial cells, and keratinocytes. Besides anti-coagulation function, CD141 is also involved in embryonic and atherosclerotic plaque development.

Product Details

Verified Reactivity	Human
Reported Reactivity	African Green, Baboon
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Cyanine7 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Additional Product Notes	BioLegend is in the process of converting the name PE/Cy7 to PE/Cyanine7. The dye molecule remains the same, so you should expect the same quality and performance from our PE/Cyanine7 products. Please contact Technical Service if you have any questions.

Product Citations

1. Alcántara-Hernández M *et al.* 2017. *Immunity*. 47(6):1037-1050. [PubMed](#)
2. Hearnden R, *et al.* 2021. *STAR Protocols*. 2(2):100422. [PubMed](#)
3. Forbester JL, *et al.* 2020. *J Virol*. 94:.. [PubMed](#)
4. Leylek R, *et al.* 2019. *Cell Rep*. 29:3736. [PubMed](#)
5. Mishra A, *et al.* 2018. *J Allergy Clin Immunol*. 142:1066. [PubMed](#)
6. Janela B, *et al.* 2019. *Immunity*. 50:1069. [PubMed](#)
7. Lee YS, *et al.* 2021. *J Immunother Cancer*. 9:.. [PubMed](#)
8. Chakraborty M, *et al.* 2021. *Cell Reports*. 34(2):108609. [PubMed](#)
9. Lee J, *et al.* 2015. *J Exp Med*. 212:385. [PubMed](#)
10. Breton G, *et al.* 2015. *J Exp Med*. 212:401. [PubMed](#)
11. Leylek R, *et al.* 2020. *Cell Rep*. 32:108180. [PubMed](#)

RRID AB_2561622 (BioLegend Cat. No. 344109)
AB_2561623 (BioLegend Cat. No. 344110)

Antigen Details

Structure	Single chain, type I membrane glycoprotein, 75 kD
Distribution	Macrophages, monocytes, subset of myeloid dendritic cells, vascular endothelial cells, keratinocytes
Function	After thrombin binding, CD141 activates protein C, which degrades clotting factors Va and VIIIa and reduces the amount of thrombin generated.
Interaction	Protein C, Thrombin-activatable fibrinolysis inhibitor (TAFI), Platelet factor 4 (PF4)
Ligand/Receptor	Thrombin
Cell Type	Dendritic cells, Endothelial cells, Macrophages, Monocytes
Biology Area	Immunology, Innate Immunity
Molecular Family	CD Molecules
Antigen References	<ol style="list-style-type: none">1. Suzuki K, <i>et al.</i> 1987. <i>EMBO J.</i> 6:1891.2. Esmon CT, <i>et al.</i> 1989. <i>J. Biol. Chem.</i> 264:4743.3. Delvaeye M, <i>et al.</i> 2009. <i>N. Engl. J. Med.</i> 361:345.4. Shi CS, <i>et al.</i> 2008. <i>Blood</i> 112:3661.5. Chen LC, <i>et al.</i> 2009. <i>J. Infect.</i> 58:368.
Gene ID	7056


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD141 (Thrombomodulin), PE anti-human CD141 (Thrombomodulin), APC anti-human CD141 (Thrombomodulin), Biotin anti-human CD141 (Thrombomodulin), PE/Cyanine7 anti-human CD141 (Thrombomodulin), PerCP/Cyanine5.5 anti-human CD141 (Thrombomodulin), Brilliant Violet 421™ anti-human CD141 (Thrombomodulin), Brilliant Violet 785™ anti-human CD141 (Thrombomodulin), Brilliant Violet 605™ anti-human CD141 (Thrombomodulin), PE/Dazzle™ 594 anti-human CD141 (Thrombomodulin), TotalSeq™-A0163 anti-human CD141 (Thrombomodulin), Alexa Fluor® 647 anti-human CD141 (Thrombomodulin), TotalSeq™-C0163 anti-human CD141 (Thrombomodulin), TotalSeq™-B0163 anti-human CD141 (Thrombomodulin), KIRAVIA Blue 520™ anti-human CD141 (Thrombomodulin), TotalSeq™-D0163 anti-human CD141 (Thrombomodulin), PE/Fire™ 810 anti-human CD141 (Thrombomodulin)

Product Data



LPS-stimulated (overnight) human peripheral blood monocytes were stained with CD141 (clone M80) PE/Cyanine7 (filled histogram) or mouse IgG1, κ PE/Cyanine7 isotype control (open histogram). Data shown was gated on the CD14+ cell population.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587