

PerCP anti-human CD14 Antibody

Catalog# / Size	367151 / 25 tests 367152 / 100 tests
Clone	63D3
Regulatory Status	RUO
Other Names	Monocyte differentiation antigen CD14, myeloid cell-specific leucine-rich glycoprotein, LPS receptor
Isotype	Mouse IgG1, κ
Description	CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein that is also known as the LPS receptor. CD14 is expressed at high levels on monocytes and macrophages, and at lower levels on granulocytes. Some dendritic cell populations such as interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells have also been reported to express CD14. As a high-affinity receptor for LPS, CD14 is involved in the clearance of gram-negative pathogens and in the upregulation of adhesion molecules and cytokine expression in monocytes and neutrophils.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Purified human peripheral blood monocytes.
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PerCP under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * PerCP has a maximum absorption of 482 nm and a maximum emission of 675 nm.
Excitation Laser	Blue Laser (488 nm)
Application References	1. Fridlender ZG, <i>et al.</i> 1999. <i>Hum. Immunol.</i> 11:1028. 2. Devitt A, <i>et al.</i> 1998. <i>Nature</i> 6675:505.
(PubMed link indicates BioLegend citation)	
Product Citations	1. Tarke A, <i>et al.</i> 2022. <i>Cell.</i> 185:847. PubMed
RRID	AB_2876693 (BioLegend Cat. No. 367151) AB_2876693 (BioLegend Cat. No. 367152)

Antigen Details

Structure	GPI-linked membrane glycoprotein.
Distribution	Monocytes, macrophages, dendritic cells, and granulocytes.
Function	LPS receptor, clearance of Gram-negative pathogens.
Interaction	LPS.
Ligand/Receptor	LPS.
Cell Type	Dendritic cells, Granulocytes, Macrophages, Monocytes, Neutrophils
Biology Area	Cell Biology, Immunology, Neuroinflammation, Neuroscience
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	<ol style="list-style-type: none">1. Stocks SC, <i>et al.</i> 1990. <i>Biochem. J.</i> 268:275.2. Wright SD, <i>et al.</i> 1990. <i>Science</i> 4975:1431.

Gene ID [929](#)


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD14, PE anti-human CD14, Biotin anti-human CD14, APC/Cyanine7 anti-human CD14, PE/Cyanine7 anti-human CD14, PerCP/Cyanine5.5 anti-human CD14, Alexa Fluor® 700 anti-human CD14, APC anti-human CD14, FITC anti-human CD14, APC/Fire™ 750 anti-human CD14, Pacific Blue™ anti-human CD14, Brilliant Violet 510™ anti-human CD14, Brilliant Violet 605™ anti-human CD14, Alexa Fluor® 647 anti-human CD14, Alexa Fluor® 488 anti-human CD14, TotalSeq™-A0051 anti-human CD14, PE/Dazzle™ 594 anti-human CD14, Brilliant Violet 750™ anti-human CD14, TotalSeq™-C0051 anti-human CD14, Brilliant Violet 421™ anti-human CD14, Brilliant Violet 785™ anti-human CD14, Brilliant Violet 711™ anti-human CD14, TotalSeq™-B0051 anti-human CD14, Spark Blue™ 550 anti-human CD14, Spark NIR™ 685 anti-human CD14, PE/Fire™ 640 anti-human CD14, APC/Fire™ 810 anti-human CD14, PerCP anti-human CD14, PE/Fire™ 700 anti-human CD14

Product Data



Human peripheral blood monocytes were stained with CD14 (clone 63D3) PerCP (filled histogram) or mouse IgG1, κ PerCP isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587