

## PerCP anti-human CD3 Antibody

<b>Catalog# / Size</b>	300325 / 25 tests 300326 / 100 tests
<b>Clone</b>	HIT3a
<b>Regulatory Status</b>	RUO
<b>Workshop</b>	V CD03.05
<b>Other Names</b>	T3, CD3ε
<b>Isotype</b>	Mouse IgG2a, κ
<b>Description</b>	CD3ε is a 20 kD chain of the CD3/T-cell receptor (TCR) complex which is composed of two CD3ε, one CD3γ, one CD3δ, one CD3ζ (CD247), and a T-cell receptor (α/β or γ/δ) heterodimer. It is found on all mature T lymphocytes, NK-T cells, and some thymocytes. CD3, also known as T3, is a member of the immunoglobulin superfamily that plays a role in antigen recognition, signal transduction, and T cell activation.

### Product Details

<b>Verified Reactivity</b>	Human
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with PerCP under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.  * PerCP has a maximum absorption of 482 nm and a maximum emission of 675 nm.
<b>Excitation Laser</b>	Blue Laser (488 nm)
<b>Application Notes</b>	Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections, immunoprecipitation, and activation of T lymphocytes <sup>4-7</sup> . The HIT3a antibody is able to stimulate T cell activation. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 300314). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 300332) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).
<b>Application References</b>	<ol style="list-style-type: none"> <li>Schlossman S, <i>et al.</i> Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.</li> <li>Knapp W. 1989. Leucocyte Typing IV. Oxford University Press New York.</li> <li>Barclay N, <i>et al.</i> 1997. The Leucocyte Antigen Facts Book. Academic Press Inc. San Diego.</li> <li>Sedelies KA, <i>et al.</i> 2004. <i>J. Biol. Chem.</i> 279:26581. (Activ)</li> <li>Rivollier A, <i>et al.</i> 2004. <i>Blood</i> 104:4029. (Activ)</li> <li>Scharschmidt E, <i>et al.</i> 2004. <i>Mol. Cell Biol.</i> 24:3860. (Activ)</li> <li>Smeltz RB. 2007. <i>J. Immunol.</i> 178:4786. (Activ)</li> </ol>
<b>(PubMed link indicates BioLegend citation)</b>	
<b>Product Citations</b>	<ol style="list-style-type: none"> <li>Zheng G, <i>et al.</i> 2021. Signal Transduct Target Ther. 6:236. <a href="#">PubMed</a></li> </ol>

2. Zhang H, *et al.* 2021. *Front Immunol.* 12:644520. [PubMed](#)
3. Schmitz T, *et al.* 2021. *J Reprod Immunol.* 148:103424. [PubMed](#)
4. Lu Y, *et al.* 2019. *Anticancer Res.* 39:5911. [PubMed](#)
5. DeKosky B, *et al.* 2016. *Proc Natl Acad Sci U S A.* 113: 2636 - 2645. [PubMed](#)
6. Sekhri P, *et al.* 2022. *Cancers (Basel).* 14: [PubMed](#)
7. Arora JK, *et al.* 2022. *iScience.* 25:104034. [PubMed](#)
8. Cheng B, *et al.* 2022. *Cancer Commun (Lond).* 42:17. [PubMed](#)
9. Yang L, *et al.* 2020. *Genes Dis.* 7:128. [PubMed](#)
10. Chen Z, *et al.* 2021. *Dis Markers.* 2021:5838582. [PubMed](#)
11. Zhang X, *et al.* 2021. *Front Immunol.* 12:602492. [PubMed](#)
12. Saraiva DP, *et al.* 2018. *Front Immunol.* 2:184027778. [PubMed](#)
13. Wei J, *et al.* 2019. *J Immunother Cancer.* 7:209. [PubMed](#)
14. Li X, *et al.* 2021. *Front Cell Dev Biol.* 9:647713. [PubMed](#)

**RRID** AB\_2616609 (BioLegend Cat. No. 300325)  
 AB\_2616610 (BioLegend Cat. No. 300326)

## Antigen Details

<b>Structure</b>	Ig superfamily, with the subunits of CD3 $\gamma$ , CD3 $\delta$ , CD3 $\zeta$ (CD247) and TCR ( $\alpha/\beta$ or $\gamma/\delta$ ) forms CD3/TCR complex, 20 kD
<b>Distribution</b>	Mature T and NK-T cells, thymocyte differentiation
<b>Function</b>	Antigen recognition, signal transduction, T cell activation
<b>Ligand/Receptor</b>	Peptide antigen bound to MHC
<b>Cell Type</b>	T cells, NKT cells
<b>Biology Area</b>	Immunology
<b>Molecular Family</b>	CD Molecules, TCRs
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Barclay N, <i>et al.</i> 1993. <i>The Leucocyte FactsBook.</i> Academic Press. San Diego.</li> <li>2. Beverly P, <i>et al.</i> 1981. <i>Eur. J. Immunol.</i> 11:329.</li> <li>3. Lanier L, <i>et al.</i> 1986. <i>J. Immunol.</i> 137:2501-2507.</li> </ol>
<b>Gene ID</b>	<a href="#">916</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

APC anti-human CD3, Biotin anti-human CD3, FITC anti-human CD3, PE anti-human CD3, PE/Cyanine5 anti-human CD3, Purified anti-human CD3, APC/Cyanine7 anti-human CD3, PE/Cyanine7 anti-human CD3, Alexa Fluor® 488 anti-human CD3, Alexa Fluor® 647 anti-human CD3, Alexa Fluor® 700 anti-human CD3, PerCP anti-human CD3, PerCP/Cyanine5.5 anti-human CD3, Pacific Blue™ anti-human CD3, Ultra-LEAF™ Purified anti-human CD3, PE/Dazzle™ 594 anti-human CD3

## Product Data



Human, lysed whole blood was stained with CD3 (clone HIT3a) PerCP (filled histogram) or Mouse IgG2a,  $\kappa$  PerCP isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587