

PerCP/Cyanine5.5 anti-mouse PLZF Antibody

Catalog# / Size	145807 / 25 µg 145808 / 100 µg
Clone	9E12
Regulatory Status	RUO
Other Names	Promyelocytic leukemia zinc finger, Zbtb16, Zinc finger and BTB domain-containing protein 16, Zfp145
Isotype	Armenian Hamster IgG
Description	PLZF (promyelocytic leukemia zinc finger), also known as Zbtb16, Zinc finger and BTB domain-containing protein 16, Zfp145, is a member of the BTB-POZ family of transcription factors. It was first identified in a patient with acute promyelocytic leukemia, where a reciprocal chromosomal translocation <i>t</i> (11;17)(q23;q21) resulted in a fusion with <i>RARA</i> gene encoding retinoic acid receptor alpha. Expression of this transcriptional repressor in immune cells differs between human and mouse. In humans, PLZF is expressed in CD34 ⁺ progenitor cells and in primitive multipotent hematopoietic cell lines, NK cells, γδ T cells, CD4 ⁺ and CD8 ⁺ T cells. It is also expressed in MR1-specific mucosal-associated invariant T cells as well as in MHC class II-restricted T cells that develop via a thymocyte-thymocyte interaction. PLZF is involved in NK cell function, cellular quiescence, and growth suppression. It also inhibits gene expression induced by retinoic acid receptor. In mice, PLZF is highly expressed in immature CD1d-restricted invariant NKT (iNKT) cells, a subset of γδ (Vg1.1 ⁺ Vd6.3 ⁺) T cells, and non-invariant CD1d-restricted T cells. PLZF exists as a homodimer or in complex with PLZP, and has been shown to be involved in the development of NKT cells. It is also reported to be expressed in embryonic tissues, giving rise to hematopoietic progenitors.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Armenian Hamster
Immunogen	Combination of peptides covering the amino, carboxyl and hinge regions of the PLZF.
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Cyanine5.5 under optimal conditions.
Concentration	0.2 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤0.06 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.
Additional Product Notes	BioLegend is in the process of converting the name PerCP/Cy5.5 to PerCP/Cyanine5.5. The dye molecule remains the same, so you should expect the same quality and performance from our PerCP/Cyanine5.5 products. Contact Technical Service if you have any questions.
Product Citations	<ol style="list-style-type: none"> 1. Harsha Krovi S, <i>et al.</i> 2020. Nat Commun. 4.790277778. PubMed 2. Trittel S, <i>et al.</i> 2019. Sci Rep. 9:16362. PubMed 3. Kim TJ, <i>et al.</i> 2019. Nat Commun. 10:3258. PubMed 4. Toubal A, <i>et al.</i> 2020. Nat Commun. 3755:11. PubMed
RRID	AB_2566166 (BioLegend Cat. No. 145807)

Antigen Details

Structure	BTB domain-containing
Distribution	iNKT cells, γ/δ (Vg1.1 ⁺ Vd6.3 ⁺) T cells, non-invariant CD1d-restricted T cells
Function	Involved in NKT cell development
Cell Type	NKT cells, T cells, Mesenchymal Stem Cells
Biology Area	Cell Biology, Immunology, Signal Transduction, Stem Cells, Transcription Factors
Antigen References	<ol style="list-style-type: none"> 1. Constantinides MG, <i>et al.</i> 2011. <i>J. Immunol.</i> 187:309. 2. Kovalovsky D, <i>et al.</i> 2008. <i>Nat. Immunol.</i> 9:1055. 3. Chen Z, <i>et al.</i> 1993. <i>EMBO J.</i> 12:1161. 4. Chen Z, <i>et al.</i> 1994. <i>Proc. Natl. Acad. Sci. USA</i> 91:1178.
Gene ID	235320

Related Protocols



[Surface and Intracellular Cytokine Staining for Flow Cytometry - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

PE anti-mouse PLZF, PE/Cyanine7 anti-mouse PLZF, PerCP/Cyanine5.5 anti-mouse PLZF

Product Data



BALB/c thymocytes were surface stained with CD3 FITC, then fixed and permeabilized using Fixation buffer and Permeabilization wash buffer, and then were stained with PLZF (clone 9E12) PerCP/Cyanine5.5 (top) or Armenian hamster IgG PerCP/Cyanine5.5 Isotype control (bottom)

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license).

BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587