

Purified anti-C/EBP β (3 isoforms C/EBP β , LAP, LIP) Antibody

Catalog# / Size	606202 / 100 μ g
Clone	1H7
Regulatory Status	RUO
Other Names	CCAAT/enhancer-binding protein beta, Alpha-1-acid glycoprotein/enhancer binding protein, Interleukin 6-dependent DNA-binding protein (IL-6DBP), Nuclear factor of interleukin 6 (NF-IL6), Liver-enriched transcriptional activator protein (LAP)
Isotype	Mouse IgG2a, κ
Description	C/EBP β (also known as CCAAT/enhancer-binding protein beta, interleukin 6-dependent DNA-binding protein, and nuclear factor of interleukin 6 [NF-IL6]) is a member of the basic leucine zipper C/EBP transcription factor family. C/EBP has three isoforms that include C/EBP β , LAP, and LIP (35 kD, 32 kD, and 20 kD, respectively). This protein is localized in the nucleus and functions as a transcription factor in immune, inflammatory, acute-phase, cytokine, collagen type I gene expression. It is upregulated by growth hormones and has been shown to interact with CEBP- α , CEBP- δ , CEBP- γ , CEBP- ϵ , and CEBP- ζ . The C/EBP β protein can be modified by phosphorylation and ubiquitination. The 1H7 monoclonal antibody has been reported to be useful for Western blotting, immunoprecipitation, and immunofluorescence of mouse and rat C/EBP β , LIP, and LAP isoforms. The 1H7 antibody does not recognize human C/EBP β , LIP or LAP.

Product Details

Verified Reactivity	Mouse, Rat
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Recombinant LIP
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography.
Concentration	0.5 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C.
Application	WB - Quality tested ChIP, IP, ICC - Reported in the literature, not verified in house
Recommended Usage	Each lot of this antibody is quality control tested by Western blotting . Western blotting, suggested working dilution(s): Use 5 μ g antibody per 5 mL antibody dilution buffer for each mini-gel. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes	Recognizes three mouse isoforms: C/EBP β , LAP, LIP. Does not cross-react with human.
Application References (PubMed link indicates BioLegend citation)	1. Su, <i>et al.</i> 2003. <i>J. Biol. Chem.</i> 278:51150. (ChIP) 2. Blidner AG, <i>et al.</i> 2015. <i>J Immunol.</i> 194:3452. PubMed
Product Citations	1. Huang JQ, <i>et al.</i> 2021. <i>Front Mol Biosci.</i> 8:725275. PubMed 2. Ebert SM, <i>et al.</i> 2020. <i>J Biol Chem.</i> 295:2787. PubMed 3. Blidner A, <i>et al.</i> 2015. <i>J Immunol.</i> 194:3452. PubMed 4. Sharma N, <i>et al.</i> 2021. <i>Front Mol Biosci.</i> 7:603168. PubMed
RRID	AB_315675 (BioLegend Cat. No. 606202)

Antigen Details

Structure	C/EBP transcription factor family, basic leucine zipper, dimer. Isoforms C/EBP β , LAP, LIP, approximately 35 kD, 32 kD, 20 kD, respectively
Distribution	Nuclear
Function	Transcription factor in immune, inflammatory, acute-phase, cytokine, collagen type I genes
Interaction	CEBP- α , CEBP- β , CEBP- δ , CEBP- γ , CEBP- ϵ , CEBP- ζ
Modification	Phosphorylation, Ubiquitination
Biology Area	Cell Biology, Transcription Factors
Molecular Family	Nuclear Markers
Antigen References	1. Chang C, <i>et al.</i> 1990. <i>Mol. Cell Biol.</i> 10:6642. 2. Eaton E, <i>et al.</i> 2001. <i>J. Cell Physiol.</i> 189:91. 3. Piwien-Pilipuk G, <i>et al.</i> 2002. <i>J. Biol. Chem.</i> 277:44557. 4. Hattori T, <i>et al.</i> 2003. <i>Oncogene.</i> 22:1273.
Regulation	Growth hormone
Gene ID	12608 25301


Related Protocols

[Western Blotting Protocol](#)

Other Formats

Purified anti-C/EBP β (3 isoforms C/EBP β , LAP, LIP), Biotin anti-C/EBP β (3 isoforms C/EBP β , LAP, LIP)

Product Data



Total cell lysates (15 μ g total protein) from THP-1 (negative control), Raw264.7, P388D1 and NIH/3T3 cells were resolved by 4-12% Bis-Tris gel electrophoresis, transferred to a PVDF membrane, and probed with 1.0 μ g/mL of Purified anti-cEBP β Antibody, clone 1H7, overnight at 4°C. Proteins were visualized by chemiluminescence detection using HRP goat anti-mouse IgG Antibody (Cat. No. 405306) at a 1:3000 dilution. Direct-Blot™ HRP anti-GAPDH Antibody (Cat. No. 607904) was used as a loading control at a 1:50000 dilution (lower). Lane M: Molecular Weight marker.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587