

## Purified anti-mouse CD317 (BST2, PDCA-1) Antibody

<b>Catalog# / Size</b>	127102 / 100 µg
<b>Clone</b>	129C1
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	BST2, tetherin, HM1.2 antigen, bone marrow stromal antigen 2, PDCA-1
<b>Isotype</b>	Rat IgG2b, κ
<b>Description</b>	CD317, known as BST2, tetherin, HM1.2 antigen, bone marrow stromal antigen 2, or PDCA-1, is type II transmembrane glycoprotein with a molecular mass of 29-33 kD. It is predominantly expressed on Type I IFN-producing cells (IPCs) in naïve mice, but is up-regulated on most cell types following stimulation with type I IFNs and IFN-gamma. It is highly expressed on terminally differentiated normal plasmacytoid dendritic cells and some tumor cells, such as multiple myeloma, renal cell carcinoma, and melanoma cells. BST2 is a recently identified as IFN-induced cellular response factor that blocks release of HIV-1 and other retroviruses from infected cells. BST2 has been found to be the natural ligand of ILT7 in human model.

### Product Details

---

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Rat
<b>Immunogen</b>	Mouse plasmacytoid dendritic cells (DCs)
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Preparation</b>	The antibody was purified by affinity chromatography.
<b>Concentration</b>	0.5 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C.
<b>Application</b>	<a href="#">FC - Quality tested</a> <a href="#">IP - Reported in the literature, not verified in house</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 µg per 10 <sup>6</sup> cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.
<b>Application References</b> (PubMed link indicates BioLegend citation)	<ol style="list-style-type: none"> <li>Blasius AL, <i>et al.</i> 2006. <i>J. Immunol.</i> 177:3260</li> <li>Cocka LJ, <i>et al.</i> 2012. <i>PLoS Pathog.</i> 8:e1002931. <a href="#">PubMed</a>.</li> </ol>
<b>Product Citations</b>	<ol style="list-style-type: none"> <li>Cocka L, Bates P 2012. <i>PLoS One.</i> 8:e1002931. <a href="#">PubMed</a></li> <li>Abboud G, <i>et al.</i> 2018. <i>Front Immunol.</i> 9:1973. <a href="#">PubMed</a></li> <li>Prabakaran T, <i>et al.</i> 2021. <i>EBioMedicine.</i> 66:103314. <a href="#">PubMed</a></li> <li>Olety B, <i>et al.</i> 2021. <i>Sci Adv.</i> 7:eabj7398. <a href="#">PubMed</a></li> </ol>
<b>RRID</b>	AB_1953279 (BioLegend Cat. No. 127102)

### Antigen Details

---

<b>Structure</b>	Type II transmembrane glycoprotein with a molecular mass of 29-33 kD.
<b>Distribution</b>	Expressed on type I IFN-producing cells, plasmacytoid dendritic cells, and neoplastic B cells, such

as multiple myeloma.

<b>Function</b>	Recently identified as antiviral protein that blocks the release of nascent retrovirus or other particles from infected cells.
<b>Cell Type</b>	Dendritic cells, B cells
<b>Biology Area</b>	Immunology, Innate Immunity, Costimulatory Molecules
<b>Molecular Family</b>	CD Molecules
<b>Antigen References</b>	1. Douglas JL. <i>et al.</i> 2009. <i>J Virol.</i> 83(16):7931 2. Cao W <i>et al.</i> 2009. <i>J. Exp. Med.</i> 206(7):1603 3. Neil SJ. <i>et al.</i> 2008. <i>Nature</i> 451:425
<b>Gene ID</b>	<a href="#">69550</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)


## Other Formats

Purified anti-mouse CD317 (BST2, PDCA-1), PE anti-mouse CD317 (BST2, PDCA-1), Pacific Blue™ anti-mouse CD317 (BST2, PDCA-1), Alexa Fluor® 647 anti-mouse CD317 (BST2, PDCA-1)

## Product Data



SJL mouse splenocytes double stained with 129C1 PE and CD11c (N418) APC



SJL mouse splenocytes double stained with rat IgG2b isotype control PE and CD11c (N418) APC

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587

