

TotalSeq™-B0163 anti-human CD141 (Thrombomodulin) Antibody

Catalog# / Size	344127 / 10 µg
Clone	M80
Regulatory Status	RUO
Other Names	Thrombomodulin, TM, THRM, THBD, Fetomodulin, BDCA-3
Isotype	Mouse IgG1, κ
Barcode Sequence	GGATAACCGCGCTTT
Description	CD141 is a 75 kD, single chain, type I membrane glycoprotein also known as thrombomodulin, TM, THRM, THBD, and fetomodulin. CD141 is an important cofactor in the protein C anticoagulant system. After binding to its ligand thrombin, CD141 activates protein C, which degrades clotting factors Va and VIIIa, and as a consequence the amount of thrombin is reduced. CD141 is expressed on macrophages, monocytes, a subpopulation of myeloid dendritic cells, vascular endothelial cells, and keratinocytes. Besides anti-coagulation function, CD141 is also involved in embryonic and atherosclerotic plaque development.

Product Details

Verified Reactivity	Human
Reported Reactivity	African Green, Baboon
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA
Preparation	The antibody was purified by chromatography and conjugated with TotalSeq™-B oligomer under optimal conditions.
Concentration	0.5 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	PG - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis and the oligomer sequence is confirmed by sequencing. TotalSeq™-B antibodies are compatible with 10x Genomics Single Cell Gene Expression Solutions.</p> <p>To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.</p> <p>Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.</p>
Additional Product Notes	<p>TotalSeq™ reagents are designed to profile protein levels at a single cell level following an optimized protocol similar to the CITE-seq workflow. A compatible single cell device (e.g. 10x Genomics Chromium System and Reagents) and sequencer (e.g. Illumina analyzers) are required. Please contact technical support for more information, or visit biolegend.com/totalseq.</p> <p>The barcode flanking sequences are GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTNNNNNNNNNN (PCR handle), and NNNNNNNNGCTTTAAGGCCGGTCCTAGC*A*A (capture sequence). N represents either</p>

randomly selected A, C, G, or T, and * indicates a phosphorothioated bond, to prevent nuclease degradation.

View more applications data for this product in our [Scientific Poster Library](#).

RRID AB_2832671 (BioLegend Cat. No. 344127)

Antigen Details

Structure	Single chain, type I membrane glycoprotein, 75 kD
Distribution	Macrophages, monocytes, subset of myeloid dendritic cells, vascular endothelial cells, keratinocytes
Function	After thrombin binding, CD141 activates protein C, which degrades clotting factors Va and VIIIa and reduces the amount of thrombin generated.
Interaction	Protein C, Thrombin-activatable fibrinolysis inhibitor (TAFI), Platelet factor 4 (PF4)
Ligand/Receptor	Thrombin
Cell Type	Dendritic cells, Endothelial cells, Macrophages, Monocytes
Biology Area	Immunology, Innate Immunity
Molecular Family	CD Molecules
Antigen References	<ol style="list-style-type: none">1. Suzuki K, <i>et al.</i> 1987. <i>EMBO J.</i> 6:1891.2. Esmon CT, <i>et al.</i> 1989. <i>J. Biol. Chem.</i> 264:4743.3. Delvaeye M, <i>et al.</i> 2009. <i>N. Engl. J. Med.</i> 361:345.4. Shi CS, <i>et al.</i> 2008. <i>Blood</i> 112:3661.5. Chen LC, <i>et al.</i> 2009. <i>J. Infect.</i> 58:368.
Gene ID	7056

Related Protocols

[TotalSeq™-B or -C with 10x Feature Barcoding Technology](#)

Other Formats

Purified anti-human CD141 (Thrombomodulin), PE anti-human CD141 (Thrombomodulin), APC anti-human CD141 (Thrombomodulin), Biotin anti-human CD141 (Thrombomodulin), PE/Cyanine7 anti-human CD141 (Thrombomodulin), PerCP/Cyanine5.5 anti-human CD141 (Thrombomodulin), Brilliant Violet 421™ anti-human CD141 (Thrombomodulin), Brilliant Violet 785™ anti-human CD141 (Thrombomodulin), Brilliant Violet 605™ anti-human CD141 (Thrombomodulin), PE/Dazzle™ 594 anti-human CD141 (Thrombomodulin), TotalSeq™-A0163 anti-human CD141 (Thrombomodulin), Alexa Fluor® 647 anti-human CD141 (Thrombomodulin), TotalSeq™-C0163 anti-human CD141 (Thrombomodulin), TotalSeq™-B0163 anti-human CD141 (Thrombomodulin), KIRAVIA Blue 520™ anti-human CD141 (Thrombomodulin), TotalSeq™-D0163 anti-human CD141 (Thrombomodulin), PE/Fire™ 810 anti-human CD141 (Thrombomodulin)

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587

