

APC/Cyanine7 anti-human IL-2 Antibody

Catalog# / Size	500341 / 25 tests 500342 / 100 tests
Clone	MQ1-17H12
Regulatory Status	RUO
Other Names	Interleukin-2, T cell growth factor (TCGF), Eosinophil differentiation factor (EDF), Killer cell helper factor (KHF), Macrophage-activating factor for cytotoxicity I (MAF-C I), Thymocyte differentiation factor (TDF)
Isotype	Rat IgG2a, κ
Description	IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells, promoting proliferation and maturation. Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK cells, LAK cells, monocytes, and oligodendrocytes.

Product Details

Verified Reactivity	Human
Reported Reactivity	Cat, Chimpanzee, Baboon, Cynomolgus, Rhesus, Sooty Mangabey
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	<i>E. coli</i> - expressed recombinant human IL-2
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with APC/Cyanine7 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.
Excitation Laser	Red Laser (633 nm)
Application Notes	<p>ELISA or ELISPOT Capture^{2,3}: The purified MQ1-17H12 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the Biotin anti-human IL-2 antibody (Cat. No. 517605) as the detecting antibody. The Ultra-LEAF™ purified antibody is suggested for ELISPOT capture. For ELISPOT capture applications, a concentration range of 4.0 - 8.0 µg/mL is recommended.</p> <p>Additional reported applications (for the relevant formats) include: immunoprecipitation², immunohistochemical staining of paraformaldehyde-fixed, saponin-treated frozen tissue sections^{1,4-6,8}, neutralization¹³, and immunocytochemistry.</p> <p>Note: For testing human IL-2 in serum or plasma, BioLegend's LEGEND MAX™ Kit (Cat. No. 431807) is specially developed and recommended.</p> <p>Clone MQ1-17H12 cross-reacts to Cat¹⁵</p>
Additional Product Notes	BioLegend is in the process of converting the name APC/Cy7 to APC/Cyanine7. The dye molecule remains the same, so you should expect the same quality and performance from our APC/Cyanine7 products. Please contact Technical Service if you have any questions.

Application References

(PubMed link indicates
BioLegend citation)

1. Andersson J, *et al.* 1994. *Immunology* 83:16. (IHC)
2. Abrams J, *et al.* 1992. *Immunol. Rev.* 127:5. (IP)
3. Abrams JS. 1995. *Curr. Prot. Immunol.* Unit 6.20.
4. Fernandez V, *et al.* 1994. *Eur. J. Immunol.* 24:1808. (IHC)
5. Skansen-Saphir U, *et al.* 1994. *Eur. J. Immunol.* 24:916. (IHC)
6. Andersson U, *et al.* *Detection and Quantification of Gene Expression*. New York:Springer-Verlag. (IHC)
7. Prussin C, *et al.* 1995. *J. Immunol. Methods.* 188:117.
8. Raqib R, *et al.* 2002. *Infect. Immun.* 70:3199. (IHC)
9. Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113. [PubMed](#)
10. Colleton BA, *et al.* 2009. *J Virol.* 83:6288. [PubMed](#)
11. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
12. Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
13. Yeap SK, *et al.* 2013. *BMC Complement Altern. Med.* 13:145. (Neut)
14. Wu Z, *et al.* 2015. *J Virol.* 89:6435. [PubMed](#)
15. Maksaarekul S, *et al.* 2009. *Vaccine.* 28:3754 (FC) [PubMed](#)

Product Citations

1. Woolsey C, *et al.* 2020. *Scientific Reports.* 10(1):3071.. [PubMed](#)
2. Sibener LV *et al.* 2018. *Cell.* 174(3):672-687 . [PubMed](#)
3. Zhu Y, *et al.* 2019. *Cell Stem Cell.* 25:542. [PubMed](#)
4. Li YR, *et al.* 2021. *Cell Rep Med.* 2:100449. [PubMed](#)
5. Ferreira-Gomes M, *et al.* 2021. *Nat Commun.* 12:1961. [PubMed](#)

RRID

AB_2562854 (BioLegend Cat. No. 500341)
AB_2562855 (BioLegend Cat. No. 500342)

Antigen Details

Structure	Cytokine; 15.4 kD (Mammalian)
Bioactivity	Proliferation of T lymphocytes, B cells, anti-inflammatory, hematopoiesis, tumor surveillance
Cell Sources	T cells
Cell Targets	T cells, B cells, NK cells, LAK cells, monocytes, macrophages, oligodendrocytes
Receptors	High affinity heterotrimer of IL-2R α / β / γ , intermediate affinity homodimer IL-2R α (CD25; p55; Tac) and heterodimer IL-2R β (CD122)/ γ ; γ -subunit (CD132) in common with IL-4R, IL-7R, IL-13R, IL-15R
Cell Type	Tregs
Biology Area	Cell Biology, Immunology, Neuroinflammation, Neuroscience
Molecular Family	Cytokines/Chemokines
Antigen References	<ol style="list-style-type: none">1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook</i>. Academic Press, San Diego.2. Taniguchi T, <i>et al.</i> 1993. <i>Cell</i> 73:5.3. Nistico G. 1993. <i>Prog. Neurobiol.</i> 40:463.4. Waldmann T, <i>et al.</i> 1993. <i>Ann. NY Acad. Sci.</i> 685:603.
Regulation	Upregulated by NFAT; downregulated by TCF-8 and CIF (colostrums inhibitory factor)
Gene ID	3558

Related Protocols

[Surface and Intracellular Cytokine Staining for Flow Cytometry - Video](#)


[Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-human IL-2, FITC anti-human IL-2, PE anti-human IL-2, Purified anti-human IL-2, Alexa Fluor® 488 anti-human IL-2, Alexa Fluor® 647 anti-human IL-2, Alexa Fluor® 700 anti-human IL-2, PerCP/Cyanine5.5 anti-human IL-2, Pacific Blue™ anti-human IL-2, PE/Cyanine7 anti-human IL-2, Brilliant Violet 421™ anti-human IL-2, Brilliant Violet 605™ anti-human IL-2, Brilliant Violet 650™ anti-human IL-2, Brilliant Violet 510™ anti-human IL-2, Brilliant Violet 711™ anti-human IL-2, APC/Cyanine7 anti-human IL-2, Purified

anti-human IL-2 (Maxpar® Ready), PE/Dazzle™ 594 anti-human IL-2, Brilliant Violet 785™ anti-human IL-2, PerCP anti-human IL-2, APC/Fire™ 750 anti-human IL-2, Ultra-LEAF™ Purified anti-human IL-2

Product Data



Human peripheral blood lymphocytes were stimulated with PMA + Ionomycin for 6 hours (in the presence of monensin), surface stained with CD3 PE, fixed, permeabilized, and then stained with IL-2 (clone MQ1-17H12) APC/Cyanine7 (left) or rat IgG2a, κ isotype control (right).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587