

APC/Fire™ 750 Annexin V

Catalog# / Size	640952 / 25 tests 640953 / 100 tests
Regulatory Status	RUO
Other Names	Annexin A5
Description	Annexin V (or Annexin A5) is a member of the annexin family of intracellular proteins that binds to phosphatidylserine (PS) in a calcium-dependent manner. PS is normally only found on the intracellular leaflet of the plasma membrane in healthy cells, but during early apoptosis, membrane asymmetry is lost and PS translocates to the external leaflet. Fluorochrome-labeled Annexin V can then be used to specifically target and identify apoptotic cells. Annexin V Binding Buffer (Cat. No. 422201) is recommended for use with Annexin V staining. Annexin V binding alone cannot differentiate between apoptotic and necrotic cells. Therefore, we recommend using our Helix NP™ Blue (Cat. No. 425305), Helix NP™ Green (Cat. No. 425303) or Helix NP™ NIR (Cat. No. 425301). Early apoptotic cells will exclude 7-AAD and PI, while late stage apoptotic cells and necrotic cells will stain positively, due to the passage of these dyes into the nucleus where they bind to DNA.

Product Details

Verified Reactivity	All mammalian species
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this product is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric application, the suggested use of this reagent is 5 - 15 µg per 100,000 - million cells in a 100 µl volume of Annexin V Binding Buffer (Cat No. 422201). It is recommended that the reagent be titrated for optimal performance for each application. * APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.
Excitation Laser	Red Laser (633 nm)
Application Notes	<p>Annexin V Staining</p> <ol style="list-style-type: none"> 1. Wash cells twice with cold BioLegend Cell Staining Buffer (Cat. No. 420201) and then resuspend cells in Annexin V Binding Buffer (Cat. No. 422201) at a concentration of 1x10⁶ cells/mL. 2. Transfer 100 µL of cell suspension in 5 mL test tube. 3. Add 5 µL of APC/Fire™ 750 Annexin V. 4. Add 10 µL of PI solution (Cat. No. 421301) or 7-AAD (Cat. No. 420403/420404). 5. Gently vortex the cells, and incubate for 15 min at room temperature (25°C), in the dark. 6. Add 400 µL of Annexin V Binding Buffer (Cat. No. 422201) to each tube. Analyze by flow cytometry. <p>For a better experience detecting apoptosis, we now recommend Apotracker™. Cell staining with Apotracker™ is Calcium independent. Thus, no special buffers are required, and the protocol can be shortened for single-step co-staining with other reagents.</p>
Application References	<ol style="list-style-type: none"> 1. Koopman G, et al. 1994. <i>Blood</i> 84:1415. 2. Vermes I, et al. 1995. <i>J. Immunol. Methods</i> 184:39. 3. Dachary-Prigent J, et al. 1993. <i>Blood</i> 81:2554. 4. Sekine C, et al. 2009. <i>Int Immunol.</i> PubMed 5. Grujic M, et al. 2010. <i>J. Immunol.</i> 185:1730. PubMed
(PubMed link indicates BioLegend citation)	

Product Citations

1. de Winde CM, *et al.* 2021. *iScience*. 24:102976. [PubMed](#)
2. Parayath NN, *et al.* 2020. *Nat Commun.* 4.680555556. [PubMed](#)

RRID

not an antibody (BioLegend Cat. No. 640952)
not an antibody (BioLegend Cat. No. 640953)

Antigen Details

Biology Area

Apoptosis/Tumor Suppressors/Cell Death, Cell Biology, Neuroscience


Gene ID

[308](#)

Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Product Data



Human T leukemia cell line Jurkat, treated (left) or non-treated (right) with BioLegend's anti-human CD95 (EOS9.1) mAb (cat. 305704) for 4 hours at 37°C, then stained with Annexin V- APC/Fire™ 750 and Helix NP Green (cat. 425303 at 1.25nM) for 15 minutes at 37°C in Annexin Binding buffer.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587