

## APC/Fire™ 750 anti-human IL-2 Antibody

<b>Catalog# / Size</b>	500351 / 25 tests 500352 / 100 tests
<b>Clone</b>	MQ1-17H12
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	Interleukin-2, T cell growth factor (TCGF), Eosinophil differentiation factor (EDF), Killer cell helper factor (KHF), Macrophage-activating factor for cytotoxicity I (MAF-C I), Thymocyte differentiation factor (TDF)
<b>Isotype</b>	Rat IgG2a, κ
<b>Description</b>	IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells, promoting proliferation and maturation. Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK cells, LAK cells, monocytes, and oligodendrocytes.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Reported Reactivity</b>	Cat, Chimpanzee, Baboon, Cynomolgus, Rhesus, Sooty Mangabey
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Rat
<b>Immunogen</b>	<i>E. coli</i> - expressed recombinant human IL-2
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">ICFC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">intracellular immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood.  * APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.
<b>Excitation Laser</b>	Red Laser (633 nm)
<b>Application Notes</b>	<b>ELISA or ELISPOT Capture<sup>2,3</sup>:</b> The purified MQ1-17H12 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the Biotin anti-human IL-2 antibody (Cat. No. 517605) as the detecting antibody. The Ultra-LEAF™ purified antibody is suggested for ELISPOT capture. For ELISPOT capture applications, a concentration range of 4.0 - 8.0 µg/mL is recommended. <b>Additional reported applications (for the relevant formats) include:</b> immunoprecipitation <sup>2</sup> , immunohistochemical staining of paraformaldehyde-fixed, saponin-treated frozen tissue sections <sup>1,4,6,8</sup> , neutralization <sup>13</sup> , and immunocytochemistry.  <b>Note:</b> For testing human IL-2 in serum or plasma, BioLegend's LEGEND MAX™ Kit (Cat. No. 431807) is specially developed and recommended.  Clone MQ1-17H12 cross-reacts to Cat <sup>15</sup>
<b>Application References</b>	1. Andersson J, <i>et al.</i> 1994. <i>Immunology</i> 83:16. (IHC)

(PubMed link indicates BioLegend citation)

- Abrams J, *et al.* 1992. *Immunol. Rev.* 127:5. (IP)
- Abrams JS. 1995. *Curr. Prot. Immunol.* Unit 6.20.
- Fernandez V, *et al.* 1994. *Eur. J. Immunol.* 24:1808. (IHC)
- Skansen-Saphir U, *et al.* 1994. *Eur. J. Immunol.* 24:916. (IHC)
- Andersson U, *et al.* *Detection and Quantification of Gene Expression.* New York:Springer-Verlag. (IHC)
- Prussin C, *et al.* 1995. *J. Immunol. Methods.* 188:117.
- Raqib R, *et al.* 2002. *Infect. Immun.* 70:3199. (IHC)
- Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113. [PubMed](#)
- Colleton BA, *et al.* 2009. *J Virol.* 83:6288. [PubMed](#)
- Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
- Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
- Yeap SK, *et al.* 2013. *BMC Complement Altern. Med.* 13:145. (Neut)
- Wu Z, *et al.* 2015. *J Virol.* 89:6435. [PubMed](#)
- Maksaarekul S, *et al.* 2009. *Vaccine.* 28:3754 (FC) [PubMed](#)

RRID AB\_2820082 (BioLegend Cat. No. 500351)  
AB\_2820083 (BioLegend Cat. No. 500352)

## Antigen Details

---

<b>Structure</b>	Cytokine; 15.4 kD (Mammalian)
<b>Bioactivity</b>	Proliferation of T lymphocytes, B cells, anti-inflammatory, hematopoiesis, tumor surveillance
<b>Cell Sources</b>	T cells
<b>Cell Targets</b>	T cells, B cells, NK cells, LAK cells, monocytes, macrophages, oligodendrocytes
<b>Receptors</b>	High affinity heterotrimer of IL-2R $\alpha$ / $\beta$ / $\gamma$ , intermediate affinity homodimer IL-2R $\alpha$ (CD25; p55; Tac) and heterodimer IL-2R $\beta$ (CD122)/ $\gamma$ ; $\gamma$ -subunit (CD132) in common with IL-4R, IL-7R, IL-13R, IL-15R
<b>Cell Type</b>	Tregs
<b>Biology Area</b>	Cell Biology, Immunology, Neuroinflammation, Neuroscience
<b>Molecular Family</b>	Cytokines/Chemokines
<b>Antigen References</b>	<ol style="list-style-type: none"><li>Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook.</i> Academic Press, San Diego.</li><li>Taniguchi T, <i>et al.</i> 1993. <i>Cell</i> 73:5.</li><li>Nistico G. 1993. <i>Prog. Neurobiol.</i> 40:463.</li><li>Waldmann T, <i>et al.</i> 1993. <i>Ann. NY Acad. Sci.</i> 685:603.</li></ol>
<b>Regulation</b>	Upregulated by NFAT; downregulated by TCF-8 and CIF (colostrums inhibitory factor)
<b>Gene ID</b>	<a href="#">3558</a>

## Related Protocols

---

[Intracellular Flow Cytometry Staining Protocol](#)


## Other Formats

---

APC anti-human IL-2, FITC anti-human IL-2, PE anti-human IL-2, Purified anti-human IL-2, Alexa Fluor® 488 anti-human IL-2, Alexa Fluor® 647 anti-human IL-2, Alexa Fluor® 700 anti-human IL-2, PerCP/Cyanine5.5 anti-human IL-2, Pacific Blue™ anti-human IL-2, PE/Cyanine7 anti-human IL-2, Brilliant Violet 421™ anti-human IL-2, Brilliant Violet 605™ anti-human IL-2, Brilliant Violet 650™ anti-human IL-2, Brilliant Violet 510™ anti-human IL-2, Brilliant Violet 711™ anti-human IL-2, APC/Cyanine7 anti-human IL-2, Purified anti-human IL-2 (Maxpar® Ready), PE/Dazzle™ 594 anti-human IL-2, Brilliant Violet 785™ anti-human IL-2, PerCP anti-human IL-2, APC/Fire™ 750 anti-human IL-2, Ultra-LEAF™ Purified anti-human IL-2

## Product Data

---



PMA + ionomycin stimulated (6 hours) human peripheral blood mononuclear cells were stained with CD3 PerCP/Cyanine5.5 and then fixed, permeablized and stained with IL-2 (clone MQ1-17H12) APC/Fire™ 750 (left) or rat IgG2a, κ APC/Fire™ 750 isotype control (right).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587