

## Biotin anti-mouse CD154 Antibody

<b>Catalog# / Size</b>	106503 / 50 µg
<b>Clone</b>	MR1
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	CD40 ligand (CD40L), gp39, T-BAM, TRAP, Ly-62, TNFSF5
<b>Isotype</b>	Armenian Hamster IgG
<b>Description</b>	CD154 is a 39 kD TNF superfamily member also known as CD40 ligand, gp39, T-BAM, TRAP, and Ly-62. CD154 is an accessory molecule expressed predominantly on activated CD4 <sup>+</sup> lymphocytes that bind CD40. CD154 plays an important role in T-B cell costimulation. The MR1 antibody has been reported to inhibit the activation of T and B lymphocytes <i>in vitro</i> and antigen-specific lymphocyte responses <i>in vivo</i> .

### Product Details

---

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Armenian Hamster
<b>Immunogen</b>	Activated mouse Th1 clone D1.6
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Preparation</b>	The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions.
<b>Concentration</b>	0.5 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per 10 <sup>6</sup> cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.
<b>Application Notes</b>	Additional reported applications (for the relevant formats) include: immunohistochemical staining <sup>1,2</sup> of acetone-fixed frozen sections, and <i>in vitro</i> and <i>in vivo</i> blocking of ligand binding <sup>3-5</sup> . For most successful immunofluorescent staining results, it may be important to maximize signal over background by using a relatively bright fluorochrome-antibody conjugate (Cat. No. 106506) or by using a high sensitivity, three-layer staining technique (e.g., including a biotinylated antibody (Cat. No. 106504) or biotinylated anti-Armenian hamster IgG (Cat. No. 405501) second step, followed by SA <sub>v</sub> -PE (Cat. No. 405204). The Ultra-LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. Nos. 106515-106520).
<b>Application References</b> (PubMed link indicates BioLegend citation)	<ol style="list-style-type: none"><li>Lettesjö H, <i>et al.</i> 2000. <i>J. Immunol.</i> 165:4095. (IHC)</li><li>Dunn RJ, <i>et al.</i> 1997. <i>J. Histochem. Cytochem.</i> 45:129. (IHC)</li><li>Noelle RJ, <i>et al.</i> 1992. <i>P. Natl. Acad. Sci. USA</i> 89:6550. (Block)</li><li>Roy M, <i>et al.</i> 1995. <i>Eur. J. Immunol.</i> 25:596. (Block)</li><li>Foy TM, <i>et al.</i> 1994. <i>J. Exp. Med.</i> 180:157. (Block)</li><li>Lawson BR, <i>et al.</i> 2007. <i>J. Immunol.</i> 178:5366.</li></ol>
<b>Product Citations</b>	<ol style="list-style-type: none"><li>Silva-Cayetano A, <i>et al.</i> 2020. <i>Med.</i> 2(3):243-262.e8. <a href="#">PubMed</a></li></ol>
<b>RRID</b>	AB_313268 (BioLegend Cat. No. 106503)

## Antigen Details

---

<b>Structure</b>	TNF superfamily, 39 kD
<b>Distribution</b>	Activated CD4 <sup>+</sup> T cells
<b>Function</b>	T-B cell costimulation
<b>Ligand/Receptor</b>	CD40
<b>Cell Type</b>	T cells, Tregs
<b>Biology Area</b>	Costimulatory Molecules, Immunology
<b>Molecular Family</b>	Adhesion Molecules, CD Molecules
<b>Antigen References</b>	1. Barclay A, <i>et al.</i> 1997. The Leukocyte Antigen FactsBook Academic Press. 2. Noelle RJ, <i>et al.</i> 1992. <i>P. Natl. Acad. Sci. USA</i> 89:6550. 3. Bancherou J, <i>et al.</i> 1994. <i>Annu. Rev. Immunol.</i> 12:881. 4. Clark EA, <i>et al.</i> 1996. <i>P. Natl. Acad. Sci. USA</i> 83:4494.
<b>Gene ID</b>	<a href="#">21947</a>

## Related Protocols

---

[Cell Surface Flow Cytometry Staining Protocol](#)


## Other Formats

---

Biotin anti-mouse CD154, PE anti-mouse CD154, APC anti-mouse CD154, PE/Cyanine7 anti-mouse CD154, PerCP/Cyanine5.5 anti-mouse CD154, Ultra-LEAF™ Purified anti-mouse CD154

## Product Data

---



PMA- and ionomycin-stimulated (6 hrs)  
BALB/c T cells stained with MR1 PE

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587