

PE/Cyanine7 anti-human CD253 (Trail) Antibody

Catalog# / Size	308215 / 25 tests 308216 / 100 tests
Clone	RIK-2
Regulatory Status	RUO
Other Names	Apo-2 ligand, Apo-2L, TNFSF10, CD253
Isotype	Mouse IgG1, κ
Description	TRAIL is a 30 kD protein known as TNF-related apoptosis inducing ligand, CD253, TNFSF10, or Apo-2L. It can be expressed as a cell surface (30 kD) and soluble ligand (19-20 kD) produced by megakaryocytes and platelets, monocytes, neutrophils, NK cells and activated T cells. This antigen is also expressed on tumor cells and transformed cell lines. TRAIL has been reported to induce apoptosis in tumor and transformed cell lines by a caspase-dependent process. This antibody is useful for flow cytometric staining and blocking studies.

Product Details

Verified Reactivity	Human
Reported Reactivity	African Green
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Human TRAIL-transfected mouse cell line
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Cyanine7 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	Additional reported applications (for the relevant formats) include: blocking ¹⁻⁴ of TRAIL-induced apoptosis and T cell mediated cytotoxicity. The Ultra-LEAF™ purified antibody (Endotoxin <0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 308214).
Application References (PubMed link indicates BioLegend citation)	<ol style="list-style-type: none"> 1. Kayagaki N, <i>et al.</i> 1999. <i>J. Immunol.</i> 162:2639. (Block) 2. Uno K, <i>et al.</i> 2003. <i>Blood</i> 101:3658. (Block) 3. Sato K, <i>et al.</i> 2005. <i>J. Immunol.</i> 174:4025. (Block) 4. Denny MF, <i>et al.</i> 2007. <i>Blood</i> 110:2907. (Block) 5. Kemter E, <i>et al.</i> 2011. <i>Xenotransplantation.</i> 19:40. PubMed
Product Citations	<ol style="list-style-type: none"> 1. Kerdidani D, <i>et al.</i> 2022. <i>J Exp Med.</i> 219:. PubMed
RRID	AB_2832606 (BioLegend Cat. No. 308215)

Antigen Details

Structure	TNF ligand superfamily member, approximately 30 kD
Distribution	Widespread, highest expression in spleen, lung, prostate; NK, monocytes, neutrophils, activated T cells
Function	Induces apoptosis in tumorigenic and transformed cell lines, involved in T cell mediated cytotoxicity, TRAIL ligand of cognate receptors induces NF- κ B signaling in a variety of cell types
Ligand/Receptor	DR4 and DR5 cognate receptors, also binds to the decoy receptors DcR1 and DcR2
Cell Type	Monocytes, Neutrophils, NK cells, T cells
Biology Area	Immunology
Molecular Family	CD Molecules
Antigen References	<ol style="list-style-type: none"> 1. Pitti RM, <i>et al.</i> 1996. <i>J. Biol. Chem.</i> 271:12687. 2. Mariani SM, <i>et al.</i> 1998. <i>Eur. J. Immunol.</i> 28:973. 3. Dorothee G, <i>et al.</i> 2002. <i>J. Immunol.</i> 169:809. 4. Crist SA, <i>et al.</i> 2004. <i>Exp. Hematology</i> 32:1073.
Gene ID	8743


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD253 (TRAIL), PE anti-human CD253 (TRAIL), APC anti-human CD253 (TRAIL), TotalSeq™-A0803 anti-human CD253 (Trail), Ultra-LEAF™ Purified anti-human CD253 (Trail), PE/Cyanine7 anti-human CD253 (Trail)

Product Data



hTRAIL transfected L5178Y cells were stained with CD253 (Trail) (clone RIK-2) PE/Cyanine7 (filled histogram) or mouse IgG1, ? PE/Cyanine7 isotype control (open histogram)

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

