



# PE/Cyanine7 anti-human CD68 Antibody

**Catalog# / Size** 333815 / 25 tests

333816 / 100 tests

Clone Y1/82A

Regulatory Status RUO

Workshop VI MR23

Other Names Macrosialin

**Isotype** Mouse IgG2b, κ

**Description** CD68 is a 110 kD glycoprotein, also known as macrosialin, belonging to the sialomucin family.

It is closely related to the family of acidic, highly glycosylated lysosomal-associated membrane

proteins (LAMPs). CD68 is predominately expressed in cytoplasmic granules of

monocytes/macrophages, dendritic cells, and granulocytes. It is one of the useful myeloid cell

markers. Further studies have shown that CD68 is also expressed by a subset of

hematopoietic progenitors,  $\gamma/\delta$  T cells, NK cells, LAK cells, subset of B cells, fibroblasts, and endothelial cells. The biological function of CD68 is still unknown.

#### **Product Details**

Verified Reactivity Human

Antibody Type Monoclonal

Host Species Mouse

Formulation Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)

Preparation The antibody was purified by affinity chromatography and conjugated with PE/Cyanine7 under

optimal conditions.

Concentration Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration

and Expiration Lookup or Certificate of Analysis online tools.)

Storage & Handling The antibody solution should be stored undiluted between 2°C and 8°C, and protected from

prolonged exposure to light. Do not freeze.

Application <u>ICFC - Quality tested</u>

Recommended Usage Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with

flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per

million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

**Excitation Laser** Blue Laser (488 nm)

Green Laser (532 nm)/Yellow-Green Laser (561 nm)

Application Notes Additional reported application: immunohistochemical staining of frozen tissue sections. This clone

was tested in-house and does not work on formalin fixed paraffin-embedded (FFPE) tissue.

Additional Product Notes BioLegend is in the process of converting the name PE/Cy7 to PE/Cyanine7. The dye molecule

remains the same, so you should expect the same quality and performance from our PE/Cyanine7 products. Please contact <u>Technical Service</u> if you have any questions.

Application References

1. Doussis IA, et al. 1993. J. Clin. Pathol. 46:334.

(PubMed link indicates BioLegend citation) 2. Davey FR, et al. 1988. J. Clin. Pathol. 41:753.

3. Bushway ME, et al. 2014. Biol Reprod. 90(5): 110. (IF) PubMed

**Product Citations** 

1. Klemm F, et al. 2020. Cell. 181(7):1643-1660.e17. PubMed

2. Trujillo-Alonso V, et al. 2019. Nat Nanotechnol. 1.011111111. PubMed

3. Borowicz S, *et al.* 2021. PLOS ONE. 16(6):e0252197. <u>PubMed</u>

4. Deniset JF et al. 2019. Immunity. 51(1):131-140 . PubMed

**RRID** AB\_2562935 (BioLegend Cat. No. 333815)

AB\_2562936 (BioLegend Cat. No. 333816)

# **Antigen Details**

Structure Sialomucin family, 110 kD

Distribution Monocytes/macrophages, dendritic cells, granulocytes, subset of hematopoietic progenitors, γ/δ T

cells, NK cells, LAK cells, subset of B cells, fibroblasts, endothelial cells

Cell Type B cells, Dendritic cells, Endothelial cells, Fibroblasts, Granulocytes, Hematopoietic stem and

progenitors, Macrophages, Monocytes, T cells

Biology Area Cell Biology, Immunology, Neuroscience, Neuroscience Cell Markers

Molecular Family CD Molecules

Antigen References 1. Holness CL and Simmons DL. 1993. Blood 81:1607.

2. Gottfried E, et al. 2008. Scand. J. Immunol. 67:453.

3. Hameed A, et al. 1994. Hum. Pathol. 25:872.

Gene ID 968

### **Related Protocols**


Surface and Intracellular Cytokine Staining for Flow Cytometry - Video

Intracellular Flow Cytometry Staining Protocol

#### **Other Formats**

PerCP/Cyanine5.5 anti-human CD68, Purified anti-human CD68, Biotin anti-human CD68, FITC anti-human CD68, PE anti-human CD68, APC anti-human CD68, Alexa Fluor® 488 anti-human CD68, PE/Cyanine7 anti-human CD68, Alexa Fluor® 647 anti-human CD68, APC/Cyanine7 anti-human CD68, APC/Fire™ 750 anti-human CD68, Brilliant Violet 785™ anti-human CD68, Brilliant Violet 421™ anti-human CD68, TotalSeq™-B0234 anti-human CD68

## **Product Data**



Human peripheral blood mononuclear cells were fixed and permeabilized with BioLegend Fixation Buffer (Cat. No. 420801) and Permeabilization Wash Buffer (Cat. No. 421002), and then stained with anti-human CD68 (clone Y1/82A) PE/Cyanine7 (filled histogram) or mouse IgG2b, K PE/Cyanine7 isotype control (open histogram). Data shown was gated on the monocyte population.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587