

PE/Dazzle™ 594 anti-mouse CD3ε Antibody

Catalog# / Size	100347 / 25 µg 100348 / 100 µg
Clone	145-2C11
Regulatory Status	RUO
Other Names	CD3ε, T3, CD3
Isotype	Armenian Hamster IgG
Description	CD3ε is a 20 kD transmembrane protein, also known as CD3 or T3. It is a member of the Ig superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3ε forms a TCR complex by associating with the CD3δ, γ and ζ chains, as well as the TCR α/β or γ/δ chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and antigen recognition by binding the peptide/MHC antigen complex.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Armenian Hamster
Immunogen	H-2K ^b -specific mouse cytotoxic T lymphocyte clone BM10-37
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions.
Concentration	0.2 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application. * PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	Clone 145-2C11 is useful for <i>in vitro</i> blocking of target-specific CTL-mediated cell lysis ¹ , as well as T cell activation assays, inducing proliferation and cytokine production ^{1,2,7,12,16} . It also induces apoptosis in immature thymocytes ³² , and <i>in vivo</i> T cell depletion ⁸⁻¹⁰ . Additional reported applications (for relevant formats of this clone) include: immunoprecipitation ¹ , immunohistochemical staining ^{14,15} of acetone-fixed frozen sections and zinc-fixed paraffin-embedded sections, Western blotting ⁴ , complement-mediated cytotoxicity ⁶ , <i>in vitro</i> and <i>in vivo</i> stimulation of T cells ^{1,2,7,12,16} , immunofluorescent staining ⁵ , and <i>in vivo</i> T cell depletion ⁸⁻¹⁰ . The 145-2C11 antibody has been reported to block the binding of 17A2 antibody to CD3 epsilon-specific T cells ¹¹ . Clone 145-2C11 is not recommended for formalin-fixed paraffin embedded sections. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 100314). For <i>in vivo</i> studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 100340) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).

Application References

1. Leo O, et al. 1987. *P. Natl. Acad. Sci. USA* 84:1374. (IP, Activ, Block)

(PubMed link indicates BioLegend citation)

2. Kruisbeek AM, et al. 1991. *In Current Protocols in Immunology*. 3.12.1. (Activ)
3. Duke RC, et al. 1995. *Current Protocols in Immunology*. 3.17.1.
4. Salvadori S, et al. 1994. *J. Immunol.* 153:5176. (WB)
5. Payer E, et al. 1991. *J. Immunol.* 146:2536. (IF)
6. Jacobs H, et al. 1994. *Eur. J. Immunol.* 24:934. (CMCD)
7. Vossen ACTM, et al. 1995. *Eur. J. Immunol.* 25:1492. (Activ)
8. Henrickson M, et al. 1995. *Transplantation* 60:828. (Deplete)
9. Kinnaert P, et al. 1996. *Transpl. Int.* 9:386. (Deplete)
10. Han WR, et al. 1999. *Transpl. Immunol.* 7:207. (Deplete)
11. Miescher GC, et al. 1989. *Immunol. Lett.* 23:113. (Block)
12. Terrazas LI, et al. 2005. *Intl. J. Parasitology.* 35:1349. (Activ)
13. Ko SY, et al. 2005. *J. Immunol.* 175:3309.
14. Podd BS, et al. 2006. *J. Immunol.* 176:6532. (IHC-F)
15. Tilley SL, et al. 2007. *J. Immunol.* 178:3208. (IHC-F)
16. Wang W, et al. 2007. *J. Immunol.* 178:4885. (Activ)
17. Xiao S, et al. 2007. *J. Exp. Med.* 204:1691.
18. Chappaz S, et al. 2007. *Blood* doi:10.1182/blood-2007-02-074245. (FC) [PubMed](#).
19. Curtis JM, et al. 2005. *J. Immunol.* 175:4392. [PubMed](#)
20. Guo Y, et al. 2008. *Blood* 112:480. [PubMed](#)
21. Kenna TJ, et al. 2008. *Blood* 111:2091.
22. Perchonock CE, et al. 2007. *J. Immunol.* 179:1768. [PubMed](#)
23. Perchonock GE, et al. 2006. *Mol. Cell. Biol.* 26:6005. [PubMed](#)
24. Kanaya T, et al. 2008. *Am. J. Physiol. Gastrointest. Liver Physiol.* 295:G273. [PubMed](#)
25. de Koning BA, et al. 2006. *Int. Immunol.* 18:941. [PubMed](#)
26. Schulteis RD, et al. 2008. *Blood* 295:G273. [PubMed](#)
27. Qi Q, et al. 2009. *Blood* 114:564. [PubMed](#)
28. Helmersson S, et al. 2013. *Am J Pathol.* 9440:123. [PubMed](#)
29. Wu S, et al. 2014. *Clin Vaccine Immunol.* 21:156. [PubMed](#)
30. Yan J, et al. 2014. *Vaccine.* 32:2833. [PubMed](#)
31. Guiterrez DA, et al. 2014. *Diabetes.* 63:3827. [PubMed](#)
32. Shi YF, et al. 1991. *J Immunol.* 146:3340. (Apop)

Product Citations

1. Sharma NS, et al. 2020. *J Clin Invest.* 130:451. [PubMed](#)
2. Liu X, et al. 2021. *eLife.* 0.4166666666666667. [PubMed](#)
3. Sitnik S, et al. 2020. *Mol Ther Oncolytics.* 17:190. [PubMed](#)
4. Gagnon JD, et al. 2019. *Cell Rep.* 28:2169. [PubMed](#)
5. Port JR, et al. 2020. *J Virol.* 94:. [PubMed](#)
6. Fearon AE, et al. 2021. *iScience.* 24:103143. [PubMed](#)

RRID

AB_2564028 (BioLegend Cat. No. 100347)
AB_2564029 (BioLegend Cat. No. 100348)

Antigen Details

Structure	Ig superfamily, forms CD3/TCR complex with CD3 δ , γ and ζ subunits and TCR (α/β and γ/δ), 20 kD
Distribution	Thymocytes (differentiation dependent), mature T cells, NK-T cells
Function	TCR signal transduction, T cell activation, antigen recognition
Ligand/Receptor	Peptide antigen/MHC-complex
Cell Type	NKT cells, T cells, Thymocytes, Tregs
Biology Area	Immunology
Molecular Family	CD Molecules, TCRs
Antigen References	<ol style="list-style-type: none">1. Barclay A, et al. 1997. <i>The Leukocyte Antigen FactsBook</i> Academic Press.2. Davis MM. 1990. <i>Annu. Rev. Biochem.</i> 59:475.3. Weiss A, et al. 1994. <i>Cell</i> 76:263.
Gene ID	12501


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-mouse CD3 ϵ , Biotin anti-mouse CD3 ϵ , FITC anti-mouse CD3 ϵ , PE anti-mouse CD3 ϵ , PE/Cyanine5 anti-mouse CD3 ϵ , Purified anti-mouse CD3 ϵ , PE/Cyanine7 anti-mouse CD3 ϵ , Alexa Fluor® 488 anti-mouse CD3 ϵ , Alexa Fluor® 647 anti-mouse CD3 ϵ , PerCP anti-mouse CD3 ϵ , PerCP/Cyanine5.5 anti-mouse CD3 ϵ , Purified anti-mouse CD3 ϵ (Maxpar® Ready), APC/Cyanine7 anti-mouse CD3 ϵ , Pacific Blue™ anti-mouse CD3 ϵ , Brilliant Violet 421™ anti-mouse CD3 ϵ , Ultra-LEAF™ Purified anti-mouse CD3 ϵ , PE/Dazzle™ 594 anti-mouse CD3 ϵ , Brilliant Violet 510™ anti-mouse CD3 ϵ , Brilliant Violet 605™ anti-mouse CD3 ϵ , Brilliant Violet 711™ anti-mouse CD3 ϵ , Brilliant Violet 785™ anti-mouse CD3 ϵ , APC/Fire™ 750 anti-mouse CD3 ϵ , GolnVivo™ Purified anti-mouse CD3 ϵ , Spark YG™ 593 anti-mouse CD3

Product Data



C57BL/6 mouse splenocytes were stained with CD3 ϵ (clone 145-2C11) PE/Dazzle™ 594 or Armenian hamster IgG PE/Dazzle™ 594 isotype control.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587