

Purified anti-human CD134 (OX40) (Maxpar[®] Ready) Antibody

Catalog# / Size	350015 / 100 µg
Clone	Ber-ACT35 (ACT35)
Regulatory Status	RUO
Other Names	ACT35 antigen, tumor necrosis factor receptor superfamily member 4 (TNFRSF4)
Isotype	Mouse IgG1, κ
Description	CD134, also known as OX40 and TNFRSF4, is a 50 kD type I transmembrane glycoprotein. It is a member of the TNF receptor family. OX40 is expressed on activated T lymphocytes including Th1, Th2, Th17, and Treg cells. The interaction of OX40 with OX40L results in B cell proliferation and antibody secretion, regulation of primary T cell expansion, and T cell survival. OX40 influences the size of the T cell memory pool and regulation of CD4 ⁺ T cell tolerance.

Product Details

Verified Reactivity	Human
Reported Reactivity	Chimpanzee
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	HTLV 1-transformed HUT 102 cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and EDTA.
Preparation	The antibody was purified by affinity chromatography.
Concentration	1.0 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C.
Application	FC - Quality tested CyTOF[®] - Verified
Recommended Usage	This product is suitable for use with the Maxpar[®] Metal Labeling Kits . For metal labeling using Maxpar [®] Ready antibodies, proceed directly to the step to Partially Reduce the Antibody by adding 100 µl of Maxpar [®] Ready antibody to 100 µl of 4 mM TCEP-R in a 50 kDa filter and continue with the protocol. Always refer to the latest version of Maxpar [®] User Guide when conjugating Maxpar [®] Ready antibodies.
Application Notes	Additional reported applications (for the relevant formats) include: Western blotting ¹ , immunoprecipitation ¹ , immunohistochemical staining ^{2,3} of paraffin embedded ⁷ and frozen tissue sections, ELISA ⁴ , and functional assay ⁵ . The LEAF [™] or Ultra-LEAF [™] purified antibody is recommended for functional assays (contact our custom solutions team).
Additional Product Notes	Maxpar [®] is a registered trademark of Standard BioTools Inc.
Application References	<ol style="list-style-type: none"> 1. Latza U, <i>et al.</i> 1994. <i>Eur. J. Immunol.</i> 24:677. (WB, IP) 2. Durkop H, <i>et al.</i> 1995. <i>Brit. J. Haematol.</i> 91:927. (IHC) 3. Durkop H, <i>et al.</i> 1997. <i>Brit. J. Haematol.</i> 98:863. (IHC) 4. Willett B, <i>et al.</i> 2007. <i>J. Virol.</i> 81:9665. (ELISA) 5. Li M and Zhang Y. <i>et al.</i> 2005. <i>Cell. Mol. Immunol.</i> 2:467. (FA) 6. Głowiczki ML, <i>et al.</i> 2012. <i>Clin. J. Am. Soc. Nephrol.</i> 8:546. PubMed 7. Domingos PL, <i>et al.</i> 2012. <i>An. Bras. Dermatol.</i> 87:851. (IHC)
(PubMed link indicates BioLegend citation)	
Product Citations	<ol style="list-style-type: none"> 1. Gide TN, <i>et al.</i> 2019. <i>Cancer Cell.</i> 35:238. PubMed
RRID	AB_2563718 (BioLegend Cat. No. 350015)

Antigen Details

Structure	50 kD, TNF receptor family
Distribution	Activated T cells
Function	Receptor for OX40 ligand, co-stimulatory signal for T cell proliferation and survival.
Ligand/Receptor	OX40L (CD252)
Cell Type	T cells, Tregs
Biology Area	Apoptosis/Tumor Suppressors/Cell Death, Cell Biology, Costimulatory Molecules, Immunology
Molecular Family	CD Molecules, Immune Checkpoint Receptors
Antigen References	<ol style="list-style-type: none">1. Smith CA, <i>et al.</i> 1994. <i>Cell</i>. 76:959.2. Chen AL, <i>et al.</i> 1999. <i>Immunity</i>. 11:689.3. Croft M. 2010. <i>Annu. Rev. Immunol.</i> 28:57.4. Ruby CE, <i>et al.</i> 2009. <i>J. Immunol.</i> 183:5079.5. Klinger M, <i>et al.</i> 2009. <i>J. Immunol.</i> 182:4581.
Gene ID	7293



Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

FITC anti-human CD134 (OX40), Purified anti-human CD134 (OX40), PE anti-human CD134 (OX40), APC anti-human CD134 (OX40), PerCP/Cyanine5.5 anti-human CD134 (OX40), PE/Cyanine7 anti-human CD134 (OX40), Brilliant Violet 421™ anti-human CD134 (OX40), Purified anti-human CD134 (OX40) (Maxpar® Ready), Alexa Fluor® 647 anti-human CD134 (OX40), PE/Dazzle™ 594 anti-human CD134 (OX40), APC/Cyanine7 anti-human CD134 (OX40), Biotin anti-human CD134 (OX40), Brilliant Violet 510™ anti-human CD134 (OX40), Brilliant Violet 605™ anti-human CD134 (OX40), APC/Fire™ 750 anti-human CD134 (OX40), Brilliant Violet 711™ anti-human CD134 (OX40), TotalSeq™-A0158 anti-human CD134 (OX40), TotalSeq™-C0158 anti-human CD134 (OX40), TotalSeq™-B0158 anti-human CD134 (OX40), TotalSeq™-D0158 anti-human CD134 (OX40), PE/Cyanine5 anti-human CD134 (OX40), PE/Fire™ 810 anti-human CD134 (OX40)

Product Data



Human PBMCs were incubated for 3 days in media alone (top) or with PHA (bottom) and then stained with 154Sm-anti-CD45 (HI30) and 158Gd-anti-CD134 (ACT35). CD45+CD14- cells are displayed in the analysis. Data provided by DVS Sciences.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587