

TotalSeq™-A0080 anti-human CD8a Antibody

Catalog# / Size	301067 / 10 µg
Clone	RPA-T8
Regulatory Status	RUO
Workshop	IV T171
Other Names	T8, Leu2
Isotype	Mouse IgG1, κ
Barcode Sequence	GCTGCGCTTCCATT
Description	CD8a is a 32-34 kD type I glycoprotein. It forms a homodimer (CD8a/a) or heterodimer (CD8a/b) with CD8b. CD8, also known as T8 and Leu2, is a member of the immunoglobulin superfamily found on the majority of thymocytes, a subset of peripheral blood T cells, and NK cells (which express almost exclusively CD8a homodimers). CD8 acts as a co-receptor with MHC class I-restricted T cell receptors in antigen recognition and T cell activation, and has been shown to play a role in thymic differentiation. Two domains in CD8a are important for function: the extracellular IgSF domain binds the α3 domain of MHC class I and the cytoplasmic CXCP motif binds the tyrosine kinase p56 Lck.

Product Details

Verified Reactivity	Human, Cynomolgus, Rhesus
Reported Reactivity	Chimpanzee, Baboon, Pigtailed Macaque, Sooty Mangabey
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA.
Preparation	The antibody was purified by chromatography and conjugated with TotalSeq™-A oligomer under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	PG - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis and the oligomer sequence is confirmed by sequencing. TotalSeq™-A antibodies are compatible with 10x Genomics Single Cell Gene Expression Solutions .

To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.

Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.

Application Notes	The RPA-T8 antibody does not block the binding of HIT8a antibody to CD8a. Additional reported applications of this antibody (for the relevant formats) include: immunohistochemical staining of paraformaldehyde-fixed frozen sections ³ and costimulation of T cell responses ⁴ . This clone was tested in-house and does not work on formalin fixed paraffin-embedded (FFPE) tissue. The Ultra-LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for
--------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

functional assays (Cat. Nos. 301073 & 301074).

Additional Product Notes

TotalSeq™ reagents are designed to profile protein levels at a single cell level following an optimized protocol similar to the CITE-seq workflow. A compatible single cell device (e.g. [10x Genomics Chromium System and Reagents](#)) and sequencer (e.g. Illumina analyzers) are required. Please contact [technical support](#) for more information, or visit [biolegend.com/totalseq](#).

The barcode flanking sequences are CCTTGGCACCCGAGAATTCCA (PCR handle), and BAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA*A*A (capture sequence). B represents either C, G, or T, and * indicates a phosphorothioated bond, to prevent nuclease degradation.

View more applications data for this product in our [Scientific Poster Library](#).

Application References

(PubMed link indicates BioLegend citation)

1. Knapp W, *et al.* Eds. 1989. Leucocyte Typing IV. Oxford University Press. New York.
2. Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
3. Mack CL, *et al.* 2004. *Pediatr. Res.* 56:79. (IHC)
4. Magidovich E, *et al.* 2007. *P. Natl. Acad. Sci. USA* 104:13022.
5. Thakral D, *et al.* 2008. *J. Immunol.* 180:7431. [PubMed](#)
6. Kmiecik M, *et al.* 2009. *J. Transl. Med.* 7:89. (FC) [PubMed](#)
7. Thakral D, *et al.* 2008. *J. Immunol.* 180:7431. (FC) [PubMed](#)
8. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
9. Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
10. Stoeckius M, *et al.* 2017. *Nat. Methods.* 14:865. (PG)

Product Citations

1. Stuart T, *et al.* 2019. *Cell.* 177:1888. [PubMed](#)
2. Swanson E, *et al.* 2021. *eLife.* 10:00. [PubMed](#)
3. Hao Y, *et al.* 2021. *Cell.* 184:3573. [PubMed](#)

RRID

AB_2734248 (BioLegend Cat. No. 301067)

Antigen Details

Structure	Ig superfamily, homodimer or heterodimer with CD8 β , 32-34 kD
Distribution	Majority of thymocytes, T cell subset, NK cells
Function	MHC class I co-receptor, thymic differentiation, T cell activation
Ligand/Receptor	MHC Class I molecules
Cell Type	Dendritic cells, NK cells, T cells, Thymocytes, Tregs
Biology Area	Immunology
Molecular Family	CD Molecules
Antigen References	1. Barclay N, <i>et al.</i> 1993. The Leucocyte Antigen FactsBook. Academic Press Inc. San Diego.
Gene ID	925

Related Protocols

[TotalSeq™-A Antibodies and Cell Hashing with 10x Single Cell 3' Reagent Kit v3 3.1 Protocol](#)

Other Formats

APC anti-human CD8a, APC/Cyanine7 anti-human CD8a, Biotin anti-human CD8a, FITC anti-human CD8a, PE anti-human CD8a, PE/Cyanine5 anti-human CD8a, PE/Cyanine7 anti-human CD8a, Purified anti-human CD8a, Alexa Fluor® 488 anti-human CD8a, Alexa Fluor® 647 anti-human CD8a, Pacific Blue™ anti-human CD8a, Alexa Fluor® 700 anti-human CD8a, PerCP anti-human CD8a, PerCP/Cyanine5.5 anti-human CD8a, Brilliant Violet 421™ anti-human CD8a, Brilliant Violet 570™ anti-human CD8a, Brilliant Violet 605™ anti-human CD8a, Brilliant Violet 650™ anti-human CD8a, Brilliant Violet 711™ anti-human CD8a, Brilliant Violet 785™ anti-human CD8a, Brilliant Violet 510™ anti-human CD8a, Purified anti-human CD8a (Maxpar® Ready), Alexa Fluor® 594 anti-human CD8a, PE/Dazzle™ 594 anti-human CD8a, APC/Fire™ 750 anti-human CD8a, TotalSeq™-A0080 anti-human CD8a, TotalSeq™-B0080 anti-human CD8a, TotalSeq™-C0080 anti-human CD8a, Ultra-LEAF™ Purified anti-human CD8a, Spark Violet™ 423 anti-human CD8a Antibody

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587