

TotalSeq™-D0146 anti-human CD69 Antibody

Catalog# / Size	310963 / 10 µg
Clone	FN50
Regulatory Status	RUO
Workshop	IV A91
Other Names	Very Early Activation Antigen (VEA), Activation inducer molecule (AIM)
Isotype	Mouse IgG1, κ
Barcode Sequence	GTCTCTTGGCTTAAA
Description	CD69 is a 27-33 kD type II transmembrane protein also known as activation inducer molecule (AIM), very early activation antigen (VEA), and MLR3. It is a member of the C-type lectin family, expressed as a disulfide-linked homodimer. Other members of this receptor family include NKG2, NKR-P1 CD94, and Ly49. CD69 is transiently expressed on activated leukocytes including T cells, thymocytes, B cells, NK cells, neutrophils, and eosinophils. CD69 is constitutively expressed by a subset of medullary mature thymocytes, platelets, mantle B cells, and certain CD4 ⁺ T cells in germinal centers of normal lymph nodes. CD69 is involved in early events of lymphocyte, monocyte, and platelet activation, and has a functional role in redirected lysis mediated by activated NK cells.

Product Details

Verified Reactivity	Human
Reported Reactivity	African Green, Baboon, Chimpanzee, Cynomolgus, Pigtailed Macaque, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA
Preparation	The antibody was purified by chromatography and conjugated with TotalSeq™-D oligomer under optimal conditions.
Concentration	0.5 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	PG - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis and the oligomer sequence is confirmed by sequencing. TotalSeq™-D antibodies are compatible with Mission Bio's Tapestry Single-Cell Sequencing Platform for simultaneous detection of DNA and Protein.</p> <p>To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.</p> <p>Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.</p>
Application Notes	Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections ² , immunofluorescence microscopy ³ , and spatial biology (IBEX) ^{8,9} .

Additional Product Notes TotalSeq™-D reagents are designed to profile protein expression at single cell level. The [Mission Bio Tapestri platform](#) and sequencer (e.g. Illumina analyzers) are required. Please contact [technical support](#) for more information, or visit biolegend.com/totalseq/single-cell-dna

The barcode flanking sequences are CGAGATGACTACGCTACTCATGG (PCR handle), and GAGCCGATCTAGTATCTCAGT*C*G (capture sequence). * indicates a phosphorothioated bond, to prevent nuclease degradation.

View more applications data for this product in our [Application Technical Notes](#).

Application References

(PubMed link indicates BioLegend citation)

1. Knapp WB, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
2. Sakkas LI, *et al.* 1998. *Clin. and Diag. Lab. Immunol.* 5:430. (IHC)
3. Kim JR, *et al.* 2005. *BMC Immunol.* 6:3. (IF)
4. Verjans GM, *et al.* 2007. *P. Natl. Acad. Sci. USA* 104:3496.
5. Lu H, *et al.* 2009. *Toxicol Sci.* 112:363. (FC) [PubMed](#)
6. Thakral D, *et al.* 2008. *J. Immunol.* 180:7431. (FC) [PubMed](#)
7. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
8. Radtke AJ, *et al.* 2020. *Proc Natl Acad Sci USA.* 117:33455-33465. (SB) [PubMed](#)
9. Radtke AJ, *et al.* 2022. *Nat Protoc.* 17:378-401. (SB) [PubMed](#)

RRID AB_2892376 (BioLegend Cat. No. 310963)

Antigen Details

Structure	C-type lectin, type II glycoprotein, 28/32 kD
Distribution	Activated T cells, B cells, NK cells, granulocytes, thymocytes, platelets, Langerhans cells
Function	Lymphocyte, monocyte, and platelet activation, NK cell killing
Cell Type	B cells, Granulocytes, Langerhans cells, NK cells, Platelets, T cells, Thymocytes, Tregs
Biology Area	Costimulatory Molecules, Immunology
Molecular Family	CD Molecules
Antigen References	1. Schlossman S, <i>et al.</i> Eds. 1995. Leucocyte Typing V. Oxford University Press. New York. 2. Testi R, <i>et al.</i> 1994. <i>Immunol. Today</i> 15:479.
Gene ID	969

Related Protocols

[TotalSeq™-D with Mission Bio Tapestri® Single-Cell DNA + Protein Protocol](#)

Other Formats

Purified anti-human CD69, FITC anti-human CD69, PE anti-human CD69, PE/Cyanine5 anti-human CD69, APC anti-human CD69, APC/Cyanine7 anti-human CD69, PE/Cyanine7 anti-human CD69, Alexa Fluor® 488 anti-human CD69, Alexa Fluor® 647 anti-human CD69, Pacific Blue™ anti-human CD69, Alexa Fluor® 700 anti-human CD69, Biotin anti-human CD69, PerCP/Cyanine5.5 anti-human CD69, PerCP anti-human CD69, Brilliant Violet 421™ anti-human CD69, Brilliant Violet 785™ anti-human CD69, Brilliant Violet 650™ anti-human CD69, Brilliant Violet 510™ anti-human CD69, Brilliant Violet 605™ anti-human CD69, Purified anti-human CD69 (Maxpar® Ready), PE/Dazzle™ 594 anti-human CD69, Brilliant Violet 711™ anti-human CD69, APC/Fire™ 750 anti-human CD69, TotalSeq™-A0146 anti-human CD69, TotalSeq™-B0146 anti-human CD69, TotalSeq™-C0146 anti-human CD69, Brilliant Violet 750™ anti-human CD69, KIRAVIA Blue 520™ anti-human CD69, Spark NIR™ 685 anti-human CD69 Antibody, PE/Fire™ 640 anti-human CD69, Spark YG™ 581 anti-human CD69, TotalSeq™-D0146 anti-human CD69, Spark Blue™ 550 anti-human CD69

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license).

BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587