

## Brilliant Violet 711™ anti-human CD223 (LAG-3) Antibody

<b>Catalog# / Size</b>	369319 / 25 tests 369320 / 100 tests
<b>Clone</b>	11C3C65
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	CD223, LAG-3, LAG3, lymphocyte-activation gene-3
<b>Isotype</b>	Mouse IgG1, $\kappa$
<b>Description</b>	CD223, also known as LAG-3, is a 70 kD type I transmembrane glycoprotein that is involved in T-cell signaling. Similar to CD4, CD223 binds MHC class II, but with a higher affinity. CD223 negatively regulates T-cell activation. It is expressed by activated T-cells and natural killer cells (NKs), as well as regulatory T-cells. It is transiently expressed on the surface of activated T-cells in acute conditions but high expression is maintained under tolerizing conditions. CD223 deficiency results in reduced tumor growth. CD223 and PD-1 can act in synergy and reverse exhausted phenotypes, improve tumor rejection, and control viral load.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	Human LAG-3 transfected cells.
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 711™ under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	<p>Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a>. For flow cytometric staining, the suggested use of this reagent is <math>\leq 5 \mu\text{l}</math> per million cells in 100 <math>\mu\text{l}</math> staining volume or 5 <math>\mu\text{l}</math> per 100 <math>\mu\text{l}</math> of whole blood.</p> <p>Brilliant Violet 711™ excites at 405 nm and emits at 711 nm. The bandpass filter 710/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 711™ is a trademark of Sirigen Group Ltd.</p> <p><a href="#">Learn more about Brilliant Violet™.</a></p> <p>This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.</p>
<b>Excitation Laser</b>	Violet Laser (405 nm)
<b>Application Notes</b>	The staining of clone 11C3C65 cannot be blocked by clone 7H2C65, which is another anti-human CD223 (LAG-3) antibody.

## Product Citations

1. Myers JA, *et al.* 2022. JCI Insight. :. [PubMed](#)
2. Carnevale J, *et al.* 2022. Nature. 609:174. [PubMed](#)
3. Evgin L, *et al.* 2020. Nat Commun. 2.671527778. [PubMed](#)

## RRID

AB\_2716124 (BioLegend Cat. No. 369319)  
AB\_2716125 (BioLegend Cat. No. 369320)

## Antigen Details

<b>Structure</b>	70 kD transmembrane glycoprotein, Ig superfamily, highly homologous to CD4.
<b>Distribution</b>	Activated T-cells and natural killer cells (NKs) and regulatory T cells.
<b>Function</b>	Negatively regulates T-cell activation.
<b>Ligand/Receptor</b>	Binds MHC class II molecules.
<b>Cell Type</b>	Dendritic cells, NK cells, T cells, Tregs
<b>Biology Area</b>	Immunology, Inhibitory Molecules
<b>Molecular Family</b>	CD Molecules, Immune Checkpoint Receptors
<b>Antigen References</b>	<ol style="list-style-type: none"><li>1. Castelli C, <i>et al.</i> 2014. <i>Oncoimmunology</i>. 3(11):e967146.</li><li>2. Poirier N, <i>et al.</i> 2011. <i>Clin. Exp. Immunol.</i> 164:265.</li><li>3. Juno JA, <i>et al.</i> 2015. <i>Retrovirology</i>. 12:17.</li><li>4. Casati C, <i>et al.</i> 2006. <i>Cancer Res</i>. 66:4450.</li></ol>
<b>Gene ID</b>	<a href="#">3902</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)


## Other Formats

Purified anti-human CD223 (LAG-3), Alexa Fluor® 647 anti-human CD223 (LAG-3), PE anti-human CD223 (LAG-3), FITC anti-human CD223 (LAG-3), PE/Cyanine7 anti-human CD223 (LAG-3), PerCP/Cyanine5.5 anti-human CD223 (LAG-3), Brilliant Violet 421™ anti-human CD223 (LAG-3), Brilliant Violet 650™ anti-human CD223 (LAG-3), Brilliant Violet 510™ anti-human CD223 (LAG-3), Brilliant Violet 785™ anti-human CD223 (LAG-3), Brilliant Violet 711™ anti-human CD223 (LAG-3), Brilliant Violet 605™ anti-human CD223 (LAG-3), Alexa Fluor® 488 anti-human CD223 (LAG-3), Biotin anti-human CD223 (LAG-3), PE/Dazzle™ 594 anti-human CD223 (LAG-3), APC/Fire™ 750 anti-human CD223 (LAG-3), TotalSeq™-A0152 anti-human CD223 (LAG-3), TotalSeq™-C0152 anti-human CD223 (LAG-3), TotalSeq™-B0152 anti-human CD223 (LAG-3), TotalSeq™-D0152 anti-human CD223 (LAG-3), Alexa Fluor® 700 anti-human CD223 (LAG-3), Pacific Blue™ anti-human CD223 (LAG-3), PE/Cyanine5 anti-human CD223 (LAG-3), APC/Cyanine7 anti-human CD223 (LAG-3), APC/Fire™ 810 anti-human CD223 (LAG-3)

## Product Data



CD3/CD28/IL-2 stimulated (three days) peripheral blood mononuclear cells (PBMCs) were stained with CD8 APC/Cy7 and CD223 (clone 11C3C65) Brilliant Violet 711™ (top) or mouse IgG1, κ Brilliant Violet 711™ isotype control (bottom).



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587