

PE/Cyanine7 anti-human CD107a (LAMP-1) Antibody

Catalog# / Size	328617 / 25 tests 328618 / 100 tests
Clone	H4A3
Regulatory Status	RUO
Workshop	P PR-63; BP 473; P P008
Other Names	Lysosome-Associated Membrane Protein 1, LGP-120, LAMP-1
Isotype	Mouse IgG1, κ
Description	CD107a, also known as Lysosome-Associated Membrane Protein 1 (LAMP-1) or LGP-120, is a 110-140 kD type I membrane glycoprotein. Mature CD107a is heavily glycosylated from a 40 kD core protein. This molecule is located on the luminal side of lysosomes. Upon activation, CD107a is transferred to the cell membrane surface of activated platelets, activated lymphocytes, macrophages, epithelial cells, endothelial cells, and some tumor cells. CD107a has been suggested to play a role in the protection of lysosomal membrane from lysosomal hydrolases which is involved in cell adhesion and regulation of tumor metastasis, and mediates autoimmune disease progression. CD107a is a ligand for galactin and E-selectin. Surface expression of LAMP-1 has been shown to correlate with CD8 ⁺ T cell and NK cell cytotoxicity.

Product Details

Verified Reactivity	Human
Reported Reactivity	African Green, Baboon, Chimpanzee, Cynomolgus, Pigtailed Macaque, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Human adult adherent peripheral blood cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Cyanine7 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	Additional reported applications (for the relevant formats) include: Western blotting ⁸ , immunohistochemical staining ² , immunofluorescence ^{5,7} , and immunoprecipitation ⁵ . This antibody is specific to human LAMP-1. Positive control: HeLa cells; LAMP-1 molecular weight appears to be at ~110 kDa on the gel due to high glycosylation.
Additional Product Notes	BioLegend is in the process of converting the name PE/Cy7 to PE/Cyanine7. The dye molecule remains the same, so you should expect the same quality and performance from our PE/Cyanine7 products. Please contact Technical Service if you have any questions.
Application References	

1. Misse D, *et al.* 1999. *Blood* 93:2454.
2. Furuta K, *et al.* 2001. *Am. J. Pathol.* 159:449. (IHC)
3. Watanabe A, *et al.* 2011. *J. Biol. Chem.* 286:10702. [PubMed](#)
4. Baron Gaillard CL, *et al.* 2011. *Mol. Cell. Biol.* 22:5459. [PubMed](#)
5. Hauck CR and Meyer TF. 1997. *FEBS Lett.* 405:86. (IF, IP)
6. De Keersmaecker B, *et al.* 2012. *J. Virol.* 86:9351. [PubMed](#)
7. Knodler LA, *et al.* 2010. *P. Natl. Acad. Sci. USA.* 107:17733. (IF)
8. Oh J, *et al.* 2000. *Hum. Mol. Genet.* 9:375. (WB)
9. Salio M, *et al.* 2013 *PNAS.* 110:4753. [PubMed](#)

Product Citations

1. Rappazzo CG, *et al.* 2020. bioRxiv. . [PubMed](#)
2. Wang H, *et al.* 2022. *Front Immunol.* 13:852436. [PubMed](#)
3. Leeansyah E, *et al.* 2015. *PLoS Pathog.* 11: 1005072. [PubMed](#)
4. Demers K, *et al.* 2016. *PLoS Pathog.* 12: 1005805. [PubMed](#)
5. Blandine Monel *et al.* 2019. *Cell reports.* 27(1):142-153 . [PubMed](#)
6. Herter JM, *et al.* 2022. *Strahlenther Onkol.* Online ahead of print. [PubMed](#)
7. Pace M, *et al.* 2016. *PLoS Pathog.* 12: 1005782. [PubMed](#)
8. Maas RJ, *et al.* 2020. *Oncoimmunology.* 9:1843247. [PubMed](#)
9. Port JR, *et al.* 2020. *J Virol.* 94:. [PubMed](#)
10. Davis Z, *et al.* 2016. *PLoS Pathog.* 12: 1005421. [PubMed](#)
11. Dai Z, *et al.* 2022. *Signal Transduct Target Ther.* 7:85. [PubMed](#)
12. Tomić A, *et al.* 2016. *PLoS Pathog.* 12:e1006015. [PubMed](#)
13. Salzberger W, *et al.* 2015. *PLoS One.* 10: 0145324. [PubMed](#)
14. Bak S, *et al.* 2019. *Front Immunol.* 9:2953. [PubMed](#)
15. Melacarne A, *et al.* 2021. *Cell Reports.* 36(1):109312. [PubMed](#)
16. Van der Meer JMR, *et al.* 2021. *Oncoimmunology.* 10:1981049. [PubMed](#)
17. Simonetta F, *et al.* 2015. *J Immunol.* 195: 4712 - 4720. [PubMed](#)
18. Harris LD, *et al.* 2020. *Front Cell Infect Microbiol.* 10:120. [PubMed](#)
19. Tang X, *et al.* 2020. *Sci Adv.* 6:eaaz0374. [PubMed](#)

RRID

AB_11147761 (BioLegend Cat. No. 328617)
 AB_11147955 (BioLegend Cat. No. 328618)

Antigen Details

Structure	LAMP-1 is a 417 amino acid protein with a molecular mass of 45 kD.
Distribution	Macrophages, epithelial cells, endothelial cells, some tumor cells; located on the luminal side of lysosomes or on the surface of cell membranes
Function	Protect lysosomal membrane from lysosomal hydrolases, adhesion
Ligand/Receptor	Galaptin
Cell Type	Endothelial cells, Epithelial cells, Macrophages
Biology Area	Cell Biology, Immunology, Neurodegeneration, Neuroscience, Protein Trafficking and Clearance
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	<ol style="list-style-type: none"> 1. Sarafian V, <i>et al.</i> 2006. <i>Arch. Dermatol. Res.</i> 298:7381. 2. Schlossman SF, <i>et al.</i> 1995. <i>Leukocyte Typing V:White Cell Differentiation Antigens.</i> New York:Oxford University Press. 3. Sawada R, <i>et al.</i> 1993. <i>J. Biol. Chem.</i> 268:12675. 4. Chen JW, <i>et al.</i> 1988. <i>J. Biol. Chem.</i> 263:8754. 5. Chen JW, <i>et al.</i> 1986. <i>Biochem. Soc. Symp.</i> 51:97112.
Gene ID	3916

Related Protocols


[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Biotin anti-human CD107a (LAMP-1), Purified anti-human CD107a (LAMP-1), FITC anti-human CD107a (LAMP-1), PE anti-human CD107a (LAMP-1), Alexa Fluor® 488 anti-human CD107a (LAMP-1), Alexa Fluor® 647 anti-human CD107a (LAMP-1),

PerCP/Cyanine5.5 anti-human CD107a (LAMP-1), APC anti-human CD107a (LAMP-1), Pacific Blue™ anti-human CD107a (LAMP-1), Brilliant Violet 421™ anti-human CD107a (LAMP-1), PE/Cyanine7 anti-human CD107a (LAMP-1), APC/Cyanine7 anti-human CD107a (LAMP-1), Brilliant Violet 510™ anti-human CD107a (LAMP-1), Brilliant Violet 605™ anti-human CD107a (LAMP-1), Purified anti-human CD107a (LAMP-1) (Maxpar® Ready), Brilliant Violet 650™ anti-human CD107a (LAMP-1), Brilliant Violet 711™ anti-human CD107a (LAMP-1), PerCP anti-human CD107a (LAMP-1), Brilliant Violet 785™ anti-human CD107a (LAMP-1), PE/Dazzle™ 594 anti-human CD107a (LAMP-1), TotalSeq™-A0155 anti-human CD107a (LAMP-1), TotalSeq™-C0155 anti-human CD107a (LAMP-1), TotalSeq™-B0155 anti-human CD107a (LAMP-1), APC/Fire™ 750 anti-human CD107a (LAMP-1) Antibody, PE/Cyanine5 anti-human CD107a (LAMP-1)

Product Data



Thrombin-activated human peripheral blood platelets were stained with CD107a (clone H4A3) PE/Cyanine7 (filled histogram) or mouse IgG1, κ PE/Cyanine7 isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587