

PE/Dazzle™ 594 anti-mouse CD4 Antibody

| | |
|--------------------------|---|
| Catalog# / Size | 100455 / 25 µg 100456 / 100 µg |
| Clone | GK1.5 |
| Regulatory Status | RUO |
| Other Names | L3T4, T4 |
| Isotype | Rat IgG2b, κ |
| Description | CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosin kinase, lck. |

Product Details

| | |
|-------------------------------|--|
| Verified Reactivity | Mouse |
| Antibody Type | Monoclonal |
| Host Species | Rat |
| Immunogen | Mouse CTL clone V4 |
| Formulation | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide. |
| Preparation | The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. |
| Concentration | 0.2 mg/ml |
| Storage & Handling | The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze. |
| Application | FC - Quality tested |
| Recommended Usage | Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application. * PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm. |
| Excitation Laser | Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm) |
| Application Notes | Additional reported applications (for the relevant formats) include: blocking of CD4 ⁺ T cell activation ^{1,4,11} , thymocyte costimulation ³ , <i>in vitro</i> and <i>in vivo</i> depletion ^{2,5-8} , blocking of egg-sperm cell adhesion ^{1,4} , immunohistochemical staining of acetone-fixed frozen sections ^{9,10} , immunoprecipitation ^{1,2} , and spatial biology (IBEX) ^{12,13} . The GK1.5 antibody is able to block CD4 mediated cell adhesion and T cell activation. Binding of GK1.5 antibody to CD4 T cells can be blocked by RM4-5 antibody, but not RM4-4 antibody. For <i>in vivo</i> studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 100442) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin < 0.01 EU/µg). |
| Application References | <ol style="list-style-type: none"> Dialynas DP, <i>et al.</i> 1983. <i>J. Immunol.</i> 131:2445. (Block, IP) Dialynas DP, <i>et al.</i> 1983. <i>Immunol. Rev.</i> 74:29. (IP, Deplete) Wu L, <i>et al.</i> 1991. <i>J. Exp. Med.</i> 174:1617. (Costim) Godfrey DI, <i>et al.</i> 1994. <i>J. Immunol.</i> 152:4783. (Block) Gavett SH, <i>et al.</i> 1994. <i>Am. J. Respir. Cell. Mol. Biol.</i> 10:587. (Deplete) Schuyler M, <i>et al.</i> 1994. <i>Am. J. Respir. Crit. Care Med.</i> 149:1286. (Deplete) Ghobrial RR, <i>et al.</i> 1989. <i>Clin. Immunol. Immunopathol.</i> 52:486. (Deplete) Israelski DM, <i>et al.</i> 1989. <i>J. Immunol.</i> 142:954. (Deplete) |

9. Zheng B, *et al.* 1996. *J. Exp. Med.* 184:1083. (IHC)
10. Frei K, *et al.* 1997. *J. Exp. Med.* 185:2177. (IHC)
11. Felix NJ, *et al.* 2007. *Nat. Immunol.* 8:388. (Block)
12. Radtke AJ, *et al.* 2020. *Proc Natl Acad Sci U S A.* 117:33455-65. (SB) [PubMed](#)
13. Radtke AJ, *et al.* 2022. *Nat Protoc.* 17:378-401. (SB) [PubMed](#)

Product Citations

1. Wu FTH, *et al.* 2019. *Br J Cancer.* 120:196. [PubMed](#)
2. Lebratti T, *et al.* 2021. *eLife.* 10:00. [PubMed](#)
3. Pandey V, *et al.* 2021. *eLife.* 0.4166666666666667. [PubMed](#)
4. Siamishi I, *et al.* 2020. *Cell Reports.* 31(11):107756. [PubMed](#)
5. Selvaraj UM, *et al.* 2021. *Brain Behav Immun.* 95:502. [PubMed](#)
6. Lewandowski JP, *et al.* 2020. *Genome Biol.* 1.039583333. [PubMed](#)
7. Fang H, *et al.* 2020. *Nat Commun.* 4.661805556. [PubMed](#)
8. Wang S, *et al.* 2021. *Parasite.* 28:13. [PubMed](#)
9. Hoover AR, *et al.* 2022. *Clin Transl Med.* 12:e937. [PubMed](#)
10. Voeller J, *et al.* 2019. *J Immunother Cancer.* 0.530555556. [PubMed](#)
11. Iwanami N, *et al.* 2020. *iScience.* 23:101260. [PubMed](#)
12. Li X, *et al.* 2022. *Oncoimmunology.* 11:2118210. [PubMed](#)
13. van Nieuwenhuijze A, *et al.* 2018. *Front Immunol.* 9:2234. [PubMed](#)

RRID

AB_2565844 (BioLegend Cat. No. 100455)
 AB_2565845 (BioLegend Cat. No. 100456)

Antigen Details

| | |
|---------------------------|--|
| Structure | Ig superfamily, 55 kD |
| Distribution | Majority of thymocytes, T cell subset |
| Function | TCR co-receptor, T cell activation |
| Ligand/Receptor | MHC class II molecule |
| Cell Type | Dendritic cells, T cells, Thymocytes, Tregs |
| Biology Area | Immunology |
| Molecular Family | CD Molecules |
| Antigen References | <ol style="list-style-type: none"> 1. Barclay A, <i>et al.</i> 1997. <i>The Leukocyte Antigen FactsBook</i> Academic Press. 2. Bierer BE, <i>et al.</i> 1989. <i>Annu. Rev. Immunol.</i> 7:579. 3. Janeway CA. 1992. <i>Annu. Rev. Immunol.</i> 10:645. |
| Gene ID | 12504 |



Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-mouse CD4, Biotin anti-mouse CD4, FITC anti-mouse CD4, PE anti-mouse CD4, PE/Cyanine5 anti-mouse CD4, Purified anti-mouse CD4, PE/Cyanine7 anti-mouse CD4, APC/Cyanine7 anti-mouse CD4, Alexa Fluor® 647 anti-mouse CD4, Alexa Fluor® 488 anti-mouse CD4, Pacific Blue™ anti-mouse CD4, Alexa Fluor® 700 anti-mouse CD4, PerCP anti-mouse CD4, PerCP/Cyanine5.5 anti-mouse CD4, Brilliant Violet 421™ anti-mouse CD4, Ultra-LEAF™ Purified anti-mouse CD4, Alexa Fluor® 594 anti-mouse CD4, Brilliant Violet 711™ anti-mouse CD4, Brilliant Violet 510™ anti-mouse CD4, Brilliant Violet 605™ anti-mouse CD4, Brilliant Violet 785™ anti-mouse CD4, PE/Dazzle™ 594 anti-mouse CD4, APC/Fire™ 750 anti-mouse CD4, GoInVivo™ Purified anti-mouse CD4, Brilliant Violet 750™ anti-mouse CD4, Brilliant Violet 650™ anti-mouse CD4, Spark Blue™ 550 anti-mouse CD4, Spark NIR™ 685 anti-mouse CD4, KIRAVIA Blue 520™ anti-mouse CD4, PE/Fire™ 640 anti-mouse CD4, APC/Fire™ 810 anti-mouse CD4, PE/Fire™ 700 anti-mouse CD4, Spark Violet™ 538 anti-mouse CD4, Spark YG™ 593 anti-mouse CD4, Spark Blue™ 574 anti-mouse CD4 Antibody, Spark UV™ 387 anti-mouse CD4

Product Data



C57BL/6 mouse splenocytes were stained with CD3ε APC and CD4 (clone GK1.5) PE/Dazzle™ 594 (top) or rat IgG2b, κ PE/Dazzle™ 594 isotype control (bottom).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
 Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587