

Alexa Fluor® 488 anti-mouse CD4 Antibody

Catalog# / Size	100532 / 25 µg 100529 / 100 µg
Clone	RM4-5
Regulatory Status	RUO
Other Names	L3T4, T4
Isotype	Rat IgG2a, κ
Description	CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes and a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a co-receptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosine kinase lck.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	BALB/c mouse thymocytes
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 488 under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested IHC-F, 3D IHC - Verified
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤ 0.06 µg per 10 ⁶ cells in 100 µl volume. For immunohistochemistry on frozen tissue sections, a concentration range of 5.0 - 10 µg/ml is suggested. For 3D immunohistochemistry on formalin-fixed tissues, a concentration of 5.0 µg/mL is suggested. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm.

Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation.

[View full statement regarding label licenses](#)

Excitation Laser	Blue Laser (488 nm)
Application Notes	The RM4-5 antibody blocks the binding of GK1.5 antibody and H129.19 antibody to CD4 ⁺ T cells, but not RM4-4 antibody. Additional reported applications (for the relevant formats) include: blocking of ligand binding, <i>in vivo</i> depletion of CD4 ⁺ cells ¹ , and immunohistochemistry of acetone-fixed frozen tissue sections ^{2,3,11} and paraffin-embedded sections ¹¹ . Clone RM4-5 is not recommended for immunohistochemistry of formalin-fixed paraffin sections. Instead, acetone frozen or zinc-fixed paraffin sections are recommended. The Ultra-LEAF™ Purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 100575 and 100576).
Application References	1. Kruisbeek AM. 1991. <i>In Curr. Protocols Immunol.</i> pp. 4.1.1-4.1.5. (Block, Deplete)
(PubMed link indicates	2. Nitta H, et al. 1997. <i>Cell Vision</i> 4:73. (IHC)

BioLegend citation)

3. Fan WY, *et al.* 2001. *Exp. Biol. Med.* 226:1045.
4. Muraille E, *et al.* 2003. *Infect. Immun.* 71:2704. (IHC)
5. León-Ponte M, *et al.* 2007. *Blood* 109:3139. (FC)
6. Bourdeau A, *et al.* 2007. *Blood* doi:10.1182/blood-2006-08-044370. (FC)
7. Matsumoto M, *et al.* 2007. *J. Immunol.* 178:2499. [PubMed](#)
8. Shigeta A, *et al.* 2008. *Blood* 112:4915. [PubMed](#)
9. Zaborsky N, *et al.* 2010. *J. Immunol.* 184:725. [PubMed](#)
10. Rodrigues-Manzanet R, *et al.* 2010. *P. Natl Acad Sci USA* 107:8706. [PubMed](#)
11. Whiteland JL, *et al.* 1995. *J. Histochem. Cytochem.* 43:313. (IHC)

Product Citations

1. Stolp B, *et al.* 2022. *Cell Rep.* 38:110387. [PubMed](#)
2. Lin JR *et al.* 2018. *eLife.* 7 pii: e31657. [PubMed](#)
3. Chen C, *et al.* 2019. *Cell Rep.* 29:4200. [PubMed](#)
4. Stewart JM, *et al.* 2020. *ACS Biomater Sci Eng.* 6:5941. [PubMed](#)
5. Wang Y, *et al.* 2021. *Cell Rep.* 35:108948. [PubMed](#)
6. Formaglio P, *et al.* 2021. *Immunity.* 54:2724. [PubMed](#)
7. Hrdinka M, *et al.* 2016. *PLoS One.* 11: 0162863. [PubMed](#)
8. Lee CG, *et al.* 2019. *JCI Insight.* 4:5. [PubMed](#)
9. Malik A *et al.* 2018. *Immunity.* 49(3):515-530. [PubMed](#)
10. White JM, *et al.* 2020. *Mol Pharm.* 2.165972222. [PubMed](#)
11. Qadir AS, *et al.* 2021. *iScience.* 24:103348. [PubMed](#)
12. Wallace-Povirk A, *et al.* 2022. *Sci Rep.* 12:11346. [PubMed](#)
13. Kuo P, *et al.* 2016. *J Am Heart Assoc.* 5: 002610. [PubMed](#)
14. Nakamura K, *et al.* 2018. *Cancer Cell.* 33:634. [PubMed](#)
15. Song W, *et al.* 2022. *Immunity.* 55:290. [PubMed](#)

RRID

AB_493373 (BioLegend Cat. No. 100532)
 AB_389303 (BioLegend Cat. No. 100529)

Antigen Details

Structure	Ig superfamily, 55 kD
Distribution	Majority of thymocytes, T cell subset
Function	TCR co-receptor, T cell activation
Ligand/Receptor	MHC class II molecule
Cell Type	Dendritic cells, T cells, Thymocytes, Tregs
Biology Area	Immunology
Molecular Family	CD Molecules
Antigen References	<ol style="list-style-type: none"> 1. Barclay A, <i>et al.</i> 1997. <i>The Leukocyte Antigen FactsBook</i> Academic Press. 2. Bierer BE, <i>et al.</i> 1989. <i>Annu. Rev. Immunol.</i> 7:579. 3. Janeway CA. 1992. <i>Annu. Rev. Immunol.</i> 10:645.
Gene ID	12504

Related Protocols

[Immunohistochemistry Protocol for Frozen Sections](#)

[Cell Surface Flow Cytometry Staining Protocol](#)


[Ce3D™ Tissue Clearing Kit](#)

Other Formats


APC anti-mouse CD4, Biotin anti-mouse CD4, FITC anti-mouse CD4, PE anti-mouse CD4, PE/Cyanine5 anti-mouse CD4, Purified anti-mouse CD4, PE/Cyanine7 anti-mouse CD4, APC/Cyanine7 anti-mouse CD4, Alexa Fluor® 647 anti-mouse CD4, Alexa Fluor® 488 anti-mouse CD4, Pacific Blue™ anti-mouse CD4, Alexa Fluor® 700 anti-mouse CD4, PerCP anti-mouse CD4, PerCP/Cyanine5.5 anti-mouse CD4, Brilliant Violet 421™ anti-mouse CD4, APC/Fire™ 750 anti-mouse CD4, Brilliant Violet 570™ anti-mouse CD4, Brilliant Violet 605™ anti-mouse CD4, Brilliant Violet 650™ anti-mouse CD4, Brilliant Violet 711™ anti-mouse CD4,

Brilliant Violet 785™ anti-mouse CD4, Brilliant Violet 510™ anti-mouse CD4, Purified anti-mouse CD4 (Maxpar® Ready), PE/Dazzle™ 594 anti-mouse CD4, TotalSeq™-A0001 anti-mouse CD4, TotalSeq™-B0001 anti-mouse CD4, TotalSeq™-C0001 anti-mouse CD4, Ultra-LEAF™ Purified anti-mouse CD4, Spark Violet™ 423 anti-mouse CD4 (L3T4) Antibody


Product Data



C57BL/6 mouse splenocytes stained with CD4 (clone RM4-5) Alexa Fluor® 488 (solid line) or rat IgG2a, κ Alexa Fluor® 488 isotype control (dotted line).



C57BL/6 mouse frozen spleen section was fixed with 4% paraformaldehyde (PFA) for 10 minutes at room temperature and blocked with 5% FBS for 30 minutes at room temperature. Then the section was stained with 10 µg/ml of anti-mouse CD4 (clone RM4-5) Alexa Fluor® 488 (green), anti-mouse CD8b (clone YTS156.7.7) Alexa Fluor® 647 (red) and anti-mouse B220 (clone RA3-6B2) Alexa Fluor® 594 (blue) overnight at 4°C. The image was captured by 10X objective.



Paraformaldehyde-fixed (4%), 500 µm-thick mouse thymus tissue section was processed according to the Ce3DTM Tissue Clearing Kit protocol (Cat. No. 427701). The section was costained with anti-mouse CD4 Antibody (clone RM4-5) Alexa Fluor® 488 at 5 µg/mL (green), anti-mouse CD68 Antibody (clone FA-11) Alexa Fluor® 594 at 5 µg/mL (yellow), and anti-mouse CD8a Antibody (clone 53-6.7) Alexa Fluor® 647 at 5 µg/mL (magenta) and counterstained with DAPI (blue). The section was then optically cleared and mounted in a sample chamber. The image was captured with a 10X objective using Zeiss 780 confocal microscope and processed by Imaris image analysis software.

[Watch the video.](#)

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587