

Alexa Fluor® 647 anti-Bcl-2 Antibody

Catalog# / Size	658705 / 25 tests 658706 / 100 tests
Clone	100
Regulatory Status	RUO
Other Names	B-Cell CLL/Lymphoma 2 (Bcl-2), Protein Phosphatase 1, Regulatory Subunit 50
Isotype	Mouse IgG1
Description	Bcl-2 is a conserved anti-apoptotic protein that plays important roles in normal immunity. It forms homodimers or heterodimers with other Bcl-2 family members. This protein regulates apoptosis through controlling mitochondrial fusion and fission. Phosphorylation of Bcl-2 has been shown to enhance activity to allow response to extracellular growth-factor-mediated signals. Bcl-2 is also regulated by IRES, miR-15a, miR-16-1 and RNA-BP nucleolin. Chromosome 18q21.3 translocation and overexpression of Bcl-2 is frequently observed in follicular lymphomas and some diffuse large B-cell lymphomas.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Synthetic peptide of amino acids 41-54 of human bcl-2 oncoprotein
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested SB - Reported in the literature, not verified in house
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood. * Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633 nm / 635 nm. Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation. View full statement regarding label licenses
Excitation Laser	Red Laser (633 nm)
Application Notes	Additional reported application (for relevant formats) include: spatial biology (IBEX) ^{3,4} .
Additional Product Notes	Iterative Bleaching Extended multi-plexity (IBEX) is a fluorescent imaging technique capable of highly-multiplexed spatial analysis. The method relies on cyclical bleaching of panels of fluorescent antibodies in order to image and analyze many markers over multiple cycles of staining, imaging, and, bleaching. It is a community-developed open-access method developed by the Center for Advanced Tissue Imaging (CAT-I) in the National Institute of Allergy and Infectious Diseases (NIAID, NIH).
Application References	1. Pezzella F, <i>et al.</i> 1993. <i>New Eng. J. Med.</i> 329:690. (IHC)

(PubMed link indicates BioLegend citation)

1. Pezzella F, *et al.* 1990. *Am. J. Pathol.* 137:225. (IHC)
2. Radtke AJ, *et al.* 2020. *Proc Natl Acad Sci U S A.* 117:33455-65. (SB) [PubMed](#)
3. Radtke AJ, *et al.* 2022. *Nat Protoc.* 17:378-401. (SB) [PubMed](#)

Product Citations

1. Lin JR *et al.* 2018. *eLife.* 7 pii: e31657. [PubMed](#)
2. Lin J, Sorger M 2015. *Sci Rep.* 6: 8390. [PubMed](#)
3. Herrera FG, *et al.* 2019. *Int J Radiat Oncol Biol Phys.* 103:320. [PubMed](#)
4. Cai J, *et al.* 2021. *eLife.* 10:00. [PubMed](#)

RRID

AB_2563279 (BioLegend Cat. No. 658705)
AB_2563280 (BioLegend Cat. No. 658706)

Antigen Details

Structure	Two isoforms, 239 or 205 amino acids in length, with a predicted molecular weight of 26 kD or 22 kD and four 'BH' domains.
Distribution	Outer mitochondrial membrane, nuclear membrane, endoplasmic reticulum membrane.
Function	Regulates apoptosis through controlling mitochondrial fusion and fission.
Interaction	BAX, BAD, BAK, Bcl-X(L), EI24, APAF1, BBC3, BCL2L1, BNIP1, MRPL41, TP53BP2, FKBP8 BAG1, RAF1, EGLN3, G0S2, and BOP.
Cell Type	B cells
Biology Area	Apoptosis/Tumor Suppressors/Cell Death, Cell Biology, Cell Cycle/DNA Replication, Chromatin Remodeling/Epigenetics, Immunology, Mitochondrial Function, Neuroscience, Signal Transduction, Transcription Factors, Ubiquitin/Protein Degradation
Antigen References	<ol style="list-style-type: none">1. Lin P, <i>et al.</i> 2013. <i>Curr. Hematol. Malign. Rep.</i> 8:243.2. Tomita N. 2011. <i>J. Clin. Exp. Hematop.</i> 51:7.3. Kelly PN, <i>et al.</i> 2011. <i>Cell Death Differ.</i> 18:1414.4. Willimott S, <i>et al.</i> 2010. <i>Biochem. Soc. Trans.</i> 38:1571.5. Rolland SG, <i>et al.</i> 2010. <i>Curr. Opin. Cell Biol.</i> 22:852.
Gene ID	596

Related Protocols


[Surface and Intracellular Cytokine Staining for Flow Cytometry - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)


Other Formats

Purified anti-Bcl-2, Alexa Fluor® 647 anti-Bcl-2, Alexa Fluor® 488 anti-Bcl-2, PE anti-Bcl-2, Brilliant Violet 421™ anti-Bcl-2

Product Data



Human peripheral blood lymphocytes were treated with Fixation Buffer (Cat. No. 420801) and Permeabilization Wash Buffer (Cat. No. 421002), then stained with Bcl-2 (clone 100) Alexa Fluor® 647 (filled histogram) or mouse IgG1, κ Alexa Fluor® 647 isotype control (open histogram).



Confocal image of human lymph node sample acquired using the IBEX method of highly multiplexed antibody-based imaging: BCL2 (cyan) in Cycle 1, CD3 (purple) in Cycle 4. Tissues were prepared using ~1% (vol/vol) formaldehyde and a detergent. Following fixation, samples are immersed in 30% (wt/vol) sucrose for cryoprotection. Images are courtesy of Drs. Andrea J. Radtke and Ronald N. Germain of the Center for Advanced Tissue Imaging (CAT-I) in the National Institute of Allergy and Infectious Diseases (NIAID, NIH).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587