

## APC anti-human CD183 (CXCR3) Antibody

<b>Catalog# / Size</b>	353707 / 25 tests 353708 / 100 tests
<b>Clone</b>	G025H7
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	CXCR3, G protein-coupled receptor 9 (GPR9), CKR-L2
<b>Isotype</b>	Mouse IgG1, $\kappa$
<b>Description</b>	Human CXCR3, also known as GPR9, is a chemokine receptor that binds CXCL9, CXCL10, and CXCL11. It is a 38 kD seven-pass transmembrane receptor coupled to G-protein. CXCR3 is highly expressed by T cells (Th1), natural killer cells (NK cells), dendritic cells, mast cells, alveolar macrophages, eosinophils, and human airway epithelial cells. CXCR3 is important for effector lymphocyte recruitment into inflamed tissue in various inflammatory and autoimmune diseases, such as chronically inflamed liver, Crohn's disease, rheumatoid arthritis, multiple sclerosis, and inflammatory skin diseases.

### Product Details

<b>Verified Reactivity</b>	Human, Cynomolgus, Rhesus
<b>Reported Reactivity</b>	African Green, Baboon
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	Human CXCR3 transfectants
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 $\mu$ l per million cells in 100 $\mu$ l staining volume or 5 $\mu$ l per 100 $\mu$ l of whole blood.
<b>Excitation Laser</b>	Red Laser (633 nm)
<b>Product Citations</b>	<ol style="list-style-type: none"> <li>1. Zhao Y, <i>et al.</i> 2021. Front Immunol. 12:665442. <a href="#">PubMed</a></li> <li>2. Zeng W, <i>et al.</i> 2017. Front Immunol. 0.806944444. <a href="#">PubMed</a></li> <li>3. Gupta R, <i>et al.</i> 2022. Front Immunol. 13:886442. <a href="#">PubMed</a></li> <li>4. Kollis PM, <i>et al.</i> 2022. Front Immunol. 13:850226. <a href="#">PubMed</a></li> <li>5. Sutton HJ, <i>et al.</i> 2021. Cell Reports. 34(6):108684. <a href="#">PubMed</a></li> <li>6. Golovkin A, <i>et al.</i> 2021. Viruses. 13:. <a href="#">PubMed</a></li> <li>7. Kudryavtsev IV, <i>et al.</i> 2021. Curr Issues Mol Biol. 44:194. <a href="#">PubMed</a></li> <li>8. Pascual-García M, <i>et al.</i> 2019. Nat Commun. 10:2416. <a href="#">PubMed</a></li> <li>9. Harb H, <i>et al.</i> 2021. Immunity. 54(6):1186-1199.e7. <a href="#">PubMed</a></li> <li>10. Arlehamn C, <i>et al.</i> 2015. Proc Natl Acad Sci U S A. 112:147. <a href="#">PubMed</a></li> <li>11. Ritacco C, <i>et al.</i> 2021. Bone Marrow Transplant. 56:1828. <a href="#">PubMed</a></li> <li>12. Zhang J, <i>et al.</i> 2020. Nat Microbiol. . <a href="#">PubMed</a></li> <li>13. Chulpanova DS, <i>et al.</i> 2021. Biology (Basel). 10:. <a href="#">PubMed</a></li> <li>14. Bonte S, <i>et al.</i> 2021. Oncoimmunology. 10:1954800. <a href="#">PubMed</a></li> <li>15. Kubin ME, <i>et al.</i> 2017. Acta Derm Venereol. 97:449. <a href="#">PubMed</a></li> </ol>

16. Yin S, *et al.* 2015. *Sci Rep.* 5: 14432. [PubMed](#)  
 17. Seo IH, *et al.* 2021. *Cell Reports.* 36(4):109438. [PubMed](#)

**RRID** AB\_10962949 (BioLegend Cat. No. 353707)  
 AB\_10983064 (BioLegend Cat. No. 353708)

## Antigen Details

<b>Structure</b>	CXC-chemokine receptor, G protein-coupled receptor, seven-pass transmembrane receptor
<b>Distribution</b>	T cell subset, NK cells, plasmacytoid dendritic cells, GM-CSF activated CD34 <sup>+</sup> hematopoietic progenitors, mast cells, alveolar macrophages, eosinophils, and airway epithelial cells
<b>Function</b>	Essential in T cell recruitment to sites of inflammation
<b>Ligand/Receptor</b>	CXCL9, CXCL10, and CXCL11
<b>Cell Type</b>	Dendritic cells, Eosinophils, Epithelial cells, Hematopoietic stem and progenitors, Macrophages, Mast cells, NK cells, T cells, Tregs
<b>Biology Area</b>	Cell Biology, Immunology, Neuroinflammation, Neuroscience
<b>Molecular Family</b>	CD Molecules, Cytokine/Chemokine Receptors, GPCR
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Loetscher M, <i>et al.</i> 1996. <i>J. Exp. Med.</i> 184:963.</li> <li>2. Cole KE, <i>et al.</i> 1998. <i>J. Exp. Med.</i> 187:2009.</li> <li>3. Aksoy MO, <i>et al.</i> 2006. <i>Am. J. Physiol. Lung Cell Mol. Physiol.</i> 290:L909.</li> <li>4. Curbishley SM, <i>et al.</i> 2005. <i>Am. J. Pathol.</i> 167:887.</li> <li>5. Turner JE, <i>et al.</i> 2007. <i>Mini. Rev. Med. Chem.</i> 7:1089.</li> <li>6. Wenzel J, <i>et al.</i> 2008. <i>J. Invest. Dermatol.</i> 128:67.</li> </ol>
<b>Gene ID</b>	<a href="#">2833</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)


## Other Formats

Purified anti-human CD183 (CXCR3), APC/Cyanine7 anti-human CD183 (CXCR3), FITC anti-human CD183 (CXCR3), PE anti-human CD183 (CXCR3), APC anti-human CD183 (CXCR3), Alexa Fluor® 488 anti-human CD183 (CXCR3), Alexa Fluor® 647 anti-human CD183 (CXCR3), PerCP/Cyanine5.5 anti-human CD183 (CXCR3), Brilliant Violet 421™ anti-human CD183 (CXCR3), PE/Cyanine7 anti-human CD183 (CXCR3), Pacific Blue™ anti-human CD183 (CXCR3), Brilliant Violet 510™ anti-human CD183 (CXCR3), Brilliant Violet 605™ anti-human CD183 (CXCR3), Brilliant Violet 650™ anti-human CD183 (CXCR3), Brilliant Violet 711™ anti-human CD183 (CXCR3), Purified anti-human CD183 (CXCR3) (Maxpar® Ready), PE/Dazzle™ 594 anti-human CD183 (CXCR3), PerCP anti-human CD183 (CXCR3), Brilliant Violet 785™ anti-human CD183 (CXCR3), Alexa Fluor® 700 anti-human CD183 (CXCR3), Biotin anti-human CD183 (CXCR3), TotalSeq™-A0140 anti-human CD183 (CXCR3), TotalSeq™-C0140 anti-human CD183 (CXCR3), Ultra-LEAF™ Purified anti-human CD183 (CXCR3), TotalSeq™-B0140 anti-human CD183 (CXCR3), APC/Fire™ 750 anti-human CD183 (CXCR3), TotalSeq™-D0140 anti-human CD183 (CXCR3), PE/Fire™ 810 anti-human CD183 (CXCR3) Antibody, APC/Fire™ 810 anti-human CD183 (CXCR3) Antibody, PE/Cyanine5 anti-human CD183 (CXCR3) Antibody, PE/Fire™ 640 anti-human CD183 (CXCR3)

## Product Data



Human peripheral lymphocytes were stained with CD3 FITC and CXCR3 (clone G025H7) APC (top) or mouse IgG1, κ APC isotype control (bottom).



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587