

APC anti-human CD79a (Igα) Antibody

Catalog# / Size	333505 / 25 tests 333506 / 100 tests
Clone	HM47
Regulatory Status	RUO
Workshop	V cB017
Other Names	Mb-1, Iga, CD79
Isotype	Mouse IgG1, κ
Description	CD79a is a 47 kD type I integral membrane protein, also known as mb-1 or Iga. It is a member of the Ig superfamily and disulphide-associated with CD79b (B29). The interaction of CD79a/CD79b heterodimer with B cell surface Ig forms B cell antigen complex. CD79a is expressed in B cells from early pre-B to plasma cell stage. It has been shown that CD79a is also weakly expressed in some precursors of T- and myeloid cells. CD79 mediates the transport of IgM to B cell surface and transduces signals initiated by BCR aggregation.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.
Excitation Laser	Red Laser (633 nm)
Product Citations	1. Lv L, <i>et al.</i> 2019. Exp Ther Med. 17:1360. PubMed
RRID	AB_1089071 (BioLegend Cat. No. 333505) AB_2075754 (BioLegend Cat. No. 333506)

Antigen Details

Structure	47 kD, type I integral protein, Ig superfamily, CD79a/CD79b heterodimer interacts with B cell membrane Ig.
Distribution	B cells (from early pre-B to plasma cell stage), some T and myeloid cell precursors
Function	In association with CD79b and B cell membrane Ig to form B cell antigen receptor complex
Cell Type	B cells, Hematopoietic stem and progenitors

Biology Area	Immunology
Molecular Family	CD Molecules
Antigen References	<ol style="list-style-type: none"> 1. Zola Heddy, <i>et al.</i> Eds. 2007. Leukocyte and Stromal Cell Molecules:The CD markers. WILEY-LISS 2. Astsaturov IA, <i>et al.</i> 1996. <i>Leukemia</i> 10:769 3. Mson DY, <i>et al.</i> 1995 <i>Blood</i> 86:1453 4. Hashimoto M, <i>et al.</i> 2002. <i>J. Pathol.</i> 197:341
Gene ID	973

Related Protocols


[Surface and Intracellular Cytokine Staining for Flow Cytometry - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD79a (Igα), PE anti-human CD79a (Igα), APC anti-human CD79a (Igα), PerCP/Cyanine5.5 anti-human CD79a (Igα), PE/Cyanine7 anti-human CD79a (Igα), FITC anti-human CD79a (Igα), Alexa Fluor® 594 anti-human CD79a (Igα), Alexa Fluor® 647 anti-human CD79a (Igα)

Product Data



Human peripheral blood lymphocytes surface stained with CD19 (HIB19) FITC and intracellular stained with HM47 APC

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
 Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587