

## APC/Fire™ 750 anti-human CD184 (CXCR4) Antibody

<b>Catalog# / Size</b>	306537 / 25 tests 306538 / 100 tests
<b>Clone</b>	12G5
<b>Regulatory Status</b>	RUO
<b>Workshop</b>	VII 70204
<b>Other Names</b>	CXCR4, Fusin
<b>Isotype</b>	Mouse IgG2a, κ
<b>Description</b>	CD184, also known as fusin or CXCR4, is a 45 kD seven transmembrane G-protein-linked CXC chemokine receptor. CD184 is widely expressed on blood and tissue cells, including B and T cells, monocytes, macrophages, dendritic cells, granulocytes, megakaryocytes/platelets, lymphoid, myeloid precursor cells, endothelial cells, epithelial cells, astrocytes, and neurons, among other tissue cells. CD184 is the receptor for CXC chemokine SDF-1, mediates blood cell migration, and is involved in B lymphopoiesis and myelopoiesis, cardiogenesis, blood vessel formation, and cerebellar development. CXCR4 is also a coreceptor of X4 HIV-1 and an alternative receptor for some isolates of HIV-2.

### Product Details

---

<b>Verified Reactivity</b>	Human, Cynomolgus, Rhesus
<b>Reported Reactivity</b>	African Green, Baboon, Chimpanzee, Sooty Mangabey
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	CP-MAC-infected Sup-T1 cells
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood.  * APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.
<b>Excitation Laser</b>	Red Laser (633 nm)
<b>Application Notes</b>	Additional reported applications (for the relevant formats) include: immunohistochemical staining of paraffin-embedded tissue sections <sup>11</sup> , immunocytochemistry <sup>3</sup> , immunofluorescence microscopy <sup>2,6</sup> , and blocking of CD4-independent infection by HIV-2 and CD4-dependent infection by some T cell-tropic isolates of HIV-1 <sup>4,5</sup> . Clone 12G5 may not be suitable for Western blotting. <sup>10</sup> The Ultra-LEAF™ purified antibody (Endotoxin <0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. Nos. 306539 & 306540).
<b>Application References</b>	1. McKnight A, <i>et al.</i> 1997. <i>J. Virol.</i> 71:1692. 2. Endres MJ, <i>et al.</i> 1996. <i>Cell</i> 87:745. (Immunogen, IF) 3. Volin MV, <i>et al.</i> 1998. <i>Biochem. Biophys. Res. Commun.</i> 242:46. (ICC)
<b>(PubMed link indicates BioLegend citation)</b>	

4. Berndt C, *et al.* 1998. *P. Natl. Acad. Sci. USA* 95:12556. (Block)
5. Ullrich CK, *et al.* 2000. *Blood* 96:1438. (Block)
6. Murga M, *et al.* 2005. *Blood* 105:1992. (IF)
7. Thompson BD. 2007. *J. Biol. Chem.* 282:9547. (FC) [PubMed](#)
8. Isnardi I, *et al.* 2010. *Blood* 115:5026. [PubMed](#)
9. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
10. Fischer T, *et al.* 2008. *PLoS One* 3:e4069.
11. Schmid BC, *et al.* 2004. *Breast Cancer Res. Treat.* 84:247. (IHC)

**RRID** AB\_2819935 (BioLegend Cat. No. 306537)  
 AB\_2819936 (BioLegend Cat. No. 306538)

## Antigen Details

---

<b>Structure</b>	Rhodopsin family, G-protein linked seven transmembrane glycoprotein, 45 kD
<b>Distribution</b>	T cells and B cells, dendritic cells, monocytes, granulocytes, hematopoietic progenitors, endothelial cells
<b>Function</b>	B lymphopoiesis and myeloopoiesis, cardiogenesis, blood vessel formation, cerebellar development
<b>Ligand/Receptor</b>	SDF-1 receptor, coreceptor for X4 HIV-1
<b>Cell Type</b>	B cells, Dendritic cells, Endothelial cells, Granulocytes, Hematopoietic stem and progenitors, Mesenchymal Stem Cells, Monocytes, Neural Stem Cells, T cells, Tregs
<b>Biology Area</b>	Cell Biology, Immunology, Innate Immunity, Neuroinflammation, Neuroscience, Neuroscience Cell Markers, Stem Cells
<b>Molecular Family</b>	CD Molecules, Cytokine/Chemokine Receptors, GPCR
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Berger E, <i>et al.</i> 1999. <i>Annu. Rev. Immunol.</i> 17:657.</li> <li>2. Loetscher P, <i>et al.</i> 2000. <i>Adv. Immunol.</i> 74:127.</li> <li>3. Murphy P, <i>et al.</i> 2000. <i>Pharmacol. Rev.</i> 52:145.</li> </ol>
<b>Gene ID</b>	<a href="#">7852</a>

## Related Protocols

---

[Cell Surface Flow Cytometry Staining Protocol](#)


## Other Formats

---

APC anti-human CD184 (CXCR4), Biotin anti-human CD184 (CXCR4), PE anti-human CD184 (CXCR4), PE/Cyanine5 anti-human CD184 (CXCR4), Purified anti-human CD184 (CXCR4), PerCP/Cyanine5.5 anti-human CD184 (CXCR4), PE/Cyanine7 anti-human CD184 (CXCR4), Brilliant Violet 421™ anti-human CD184 (CXCR4), Brilliant Violet 605™ anti-human CD184 (CXCR4), Purified anti-human CD184 (CXCR4) (Maxpar® Ready), PE/Dazzle™ 594 anti-human CD184 (CXCR4), APC/Cyanine7 anti-human CD184 (CXCR4), Brilliant Violet 785™ anti-human CD184 (CXCR4), TotalSeq™-A0366 anti-human CD184 (CXCR4), TotalSeq™-C0366 anti-human CD184 (CXCR4), Brilliant Violet 510™ anti-human CD184 (CXCR4), Ultra-LEAF™ Purified anti-human CD184 (CXCR4), APC/Fire™ 750 anti-human CD184 (CXCR4), TotalSeq™-B0366 anti-human CD184 (CXCR4)

## Product Data

---



Human peripheral blood lymphocytes were stained with CD3 FITC and CD184 (clone 12G5) APC/Fire™ 750 (left) or Mouse IgG2a, κ isotype control APC/Fire™ 750 (right).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
 Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587