

Biotin anti-human CD73 (Ecto-5'-nucleotidase) Antibody

Catalog# / Size	344017 / 50 µg
Clone	AD2
Regulatory Status	RUO
Workshop	V B-CD73.3
Other Names	Ecto-5'-nucleotidase, E.C3.1.3.5, L-VAP-2, NT5E, 5'-NT
Isotype	Mouse IgG1, κ
Description	CD73 is a 70 kD glycoposphatidylinositol (GPI)-linked 5'-nucleotidase, which is also known as ecto-5'-nucleotidase. It converts adenosine monophosphate (AMP) to adenosine. CD73 is expressed on subsets of T and B cells, mesenchymal stem cells, follicular dendritic cells, endothelial cells, and epithelial cells. It has been reported that CD73 costimulates T cell activation, and mediates adhesion of lymphocytes to follicular dendritic cells and endothelial cells.

Product Details

Verified Reactivity	Human
Reported Reactivity	African Green, Baboon
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with biotin under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes	Additional reported applications (for the relevant formats) include: immunofluorescence ³ . Clone AD2 has been noted to induce clustering and internalization of CD73 <i>in vivo</i> and inhibit metastasis in a murine breast cancer xenograft model ⁴ .
Application References	1. Nakamura T, <i>et al.</i> 1993. <i>J. Immunol.</i> 151:6933. 2. Liao J, <i>et al.</i> 2011. <i>J Endod.</i> 37:1217. PubMed 3. Touboul C, <i>et al.</i> 2013. <i>J. Transl. Med.</i> 11:28. (IF) 4. Terp MG, <i>et al.</i> 2013. <i>J Immunol.</i> 191: 4165-73 (Block)
(PubMed link indicates BioLegend citation)	
Product Citations	1. Priglinger E, <i>et al.</i> 2021. <i>J Extracell Vesicles.</i> 10:e12156. PubMed
RRID	AB_2564561 (BioLegend Cat. No. 344017)

Antigen Details

Structure	GPI-linked 5'-nucleotidase, 70 kD
Distribution	Subsets of T cells and B cells, mesenchymal stem cells, follicular dendritic cells, endothelial cells, and epithelial cells
Function	Catalyses dephosphorylation of adenosine monophosphate, costimulates T cell activation, mediates adhesion of lymphocytes to follicular dendritic cells and endothelial cells
Cell Type	B cells, Dendritic cells, Endothelial cells, Epithelial cells, Mesenchymal Stem Cells, T cells, Tregs
Biology Area	Costimulatory Molecules, Immunology, Stem Cells
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	<ol style="list-style-type: none"> 1. Zola H, <i>et al.</i> 2007. <i>Leukocyte and stromal Cell Molecules:the CD Markers</i>. A John Wiley & Sons Inc, Publication. 2. Airas L and Jalkanen S, <i>et al.</i> 1996. <i>Blood</i> 88:1755. 3. Gutensohn W, <i>et al.</i> 1995. <i>Cell Immunol.</i> 161:213. 4. Airas L, <i>et al.</i> 1995. <i>J. Exp. Med.</i> 182:1603.
Gene ID	4907


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

FITC anti-human CD73 (Ecto-5'-nucleotidase), Brilliant Violet 421™ anti-human CD73 (Ecto-5'-nucleotidase), Purified anti-human CD73 (Ecto-5'-nucleotidase), PE anti-human CD73 (Ecto-5'-nucleotidase), APC anti-human CD73 (Ecto-5'-nucleotidase), PE/Cyanine7 anti-human CD73 (Ecto-5'-nucleotidase), Pacific Blue™ anti-human CD73 (Ecto-5'-nucleotidase), PerCP/Cyanine5.5 anti-human CD73 (Ecto-5'-nucleotidase), Biotin anti-human CD73 (Ecto-5'-nucleotidase), PE/Dazzle™ 594 anti-human CD73 (Ecto-5'-nucleotidase), APC/Cyanine7 anti-human CD73 (Ecto-5'-nucleotidase), Brilliant Violet 605™ anti-human CD73 (Ecto-5'-nucleotidase), Brilliant Violet 711™ anti-human CD73 (Ecto-5'-nucleotidase), Brilliant Violet 785™ anti-human CD73 (Ecto-5'-nucleotidase), TotalSeq™-A0577 anti-human CD73 (Ecto-5'-nucleotidase), TotalSeq™-C0577 anti-human CD73 (Ecto-5'-nucleotidase), TotalSeq™-B0577 anti-human CD73 (Ecto-5'-nucleotidase), APC/Fire™ 750 anti-human CD73 (Ecto-5'-nucleotidase), TotalSeq™-D0577 anti-human CD73 (Ecto-5'-nucleotidase), Alexa Fluor® 700 anti-human CD73 (Ecto-5'-nucleotidase)

Product Data



Human peripheral blood lymphocytes were stained CD3 APC and biotinylated CD73 (clone AD2) (top), or biotinylated mouse IgG1, κ isotype control (bottom), followed by Streptavidin PE.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587