

Brilliant Violet 421™ anti-mouse CD4 Antibody

Catalog# / Size	100543 / 125 µL 100563 / 50 µg 100544 / 500 µL
Clone	RM4-5
Regulatory Status	RUO
Other Names	L3T4, T4
Isotype	Rat IgG2a, κ
Description	CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes and a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a co-receptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosine kinase lck.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	BALB/c mouse thymocytes
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
Preparation	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions.
Concentration	µg sizes: 0.2 mg/mL µL sizes: lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested ICC, IHC-F - Verified
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining using the µg size, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µL volume. For immunocytochemistry staining using µL sizes, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. For immunohistochemistry on frozen tissue sections using the µg size, a concentration range of 5.0 - 10 µg/mL is suggested. It is recommended that the reagent be titrated for optimal performance for each application.</p> <p>Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.</p> <p>Learn more about Brilliant Violet™.</p> <p>This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.</p>
Excitation Laser	Violet Laser (405 nm)
Application Notes	The RM4-5 antibody blocks the binding of GK1.5 antibody and H129.19 antibody to CD4 ⁺ T cells, but not RM4-4 antibody. Additional reported applications (for the relevant formats) include:

blocking of ligand binding, *in vivo* depletion of CD4⁺ cells¹, and immunohistochemistry of acetone-fixed frozen tissue sections^{2,3,11} and paraffin-embedded sections¹¹. Clone RM4-5 is not recommended for immunohistochemistry of formalin-fixed paraffin sections. Instead, acetone frozen or zinc-fixed paraffin sections are recommended. The Ultra-LEAF™ Purified antibody (Endotoxin < 0.01 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 100575 and 100576).

Application References

(PubMed link indicates BioLegend citation)

1. Kruisbeek AM. 1991. *In Curr. Protocols Immunol.* pp. 4.1.1-4.1.5. (Block, Deplete)
2. Nitta H, *et al.* 1997. *Cell Vision* 4:73. (IHC)
3. Fan WY, *et al.* 2001. *Exp. Biol. Med.* 226:1045.
4. Muraille E, *et al.* 2003. *Infect. Immun.* 71:2704. (IHC)
5. León-Ponte M, *et al.* 2007. *Blood* 109:3139. (FC)
6. Bourdeau A, *et al.* 2007. *Blood* doi:10.1182/blood-2006-08-044370. (FC)
7. Matsumoto M, *et al.* 2007. *J. Immunol.* 178:2499. [PubMed](#)
8. Shigeta A, *et al.* 2008. *Blood* 112:4915. [PubMed](#)
9. Zaborsky N, *et al.* 2010. *J. Immunol.* 184:725. [PubMed](#)
10. Rodrigues-Manzanet R, *et al.* 2010. *P. Natl Acad Sci USA* 107:8706. [PubMed](#)
11. Whiteland JL, *et al.* 1995. *J. Histochem. Cytochem.* 43:313. (IHC)

Product Citations

1. Bradley CP *et al.* 2017. *Cell host & microbe.* 22(5):697-704. [PubMed](#)
2. Flamar AL, *et al.* 2020. *Immunity.* 52(4):606-619.e6. [PubMed](#)
3. Yu Y, *et al.* 2022. *iScience.* 25:105004. [PubMed](#)
4. Lechuga-Vieco AV, *et al.* 2020. *Sci Adv.* 6:eaba5345. [PubMed](#)
5. Park S, *et al.* 2021. *Front Immunol.* 11:620166. [PubMed](#)
6. Lino AC *et al.* 2018. *Immunity.* 49(1):120-133. [PubMed](#)
7. Wang D, *et al.* 2018. *Immunity.* 48:659. [PubMed](#)
8. Baptista AP *et al.* 2019. *Immunity.* 50(5):1188-1201. [PubMed](#)
9. Gimblet C *et al.* 2017. *Cell host & microbe.* 22(1):13-24. [PubMed](#)
10. Prasad M, *et al.* 2020. *Cell Mol Immunol.* [PubMed](#)
11. Yang W, *et al.* 2020. *Nat Commun.* 3.553472222. [PubMed](#)
12. Stein RR, *et al.* 2018. *Elife.* 7:e30916. [PubMed](#)
13. Delacher M, *et al.* 2021. *Immunity.* 54(4):702-720.e17. [PubMed](#)
14. Acharya N, *et al.* 2020. *Immunity.* 53(3):658-671.e6. [PubMed](#)
15. LaFleur MW, *et al.* 2019. *Nat Immunol.* 20:1335. [PubMed](#)
16. Rodriguez-Ruiz M, *et al.* 2016. *Cancer Res.* 76: 5994 - 6005. [PubMed](#)
17. Köchl R, *et al.* 2020. *Elife.* 9:00. [PubMed](#)
18. Mamedov MR, *et al.* 2018. *Immunity.* 48:350. [PubMed](#)
19. LaFleur MW, *et al.* 2019. *Nat Commun.* 10:1668. [PubMed](#)
20. Parayath NN, *et al.* 2020. *Nat Commun.* 4.680555556. [PubMed](#)
21. Zhang W, *et al.* 2022. *Nat Methods.* 19:759. [PubMed](#)
22. Xiao S, *et al.* 2020. *Cell Reports.* 32(2):107892. [PubMed](#)
23. Ben-Yehuda H, *et al.* 2021. *Mol Neurodegener.* 16:39. [PubMed](#)
24. Hirata SI, *et al.* 2020. *Allergy.* 75:1939. [PubMed](#)
25. Chen EW, *et al.* 2018. *Sci Rep.* 8:10046. [PubMed](#)
26. Mairhofer D, *et al.* 2015. *J Invest Dermatol.* 135: 2785-93. [PubMed](#)
27. Lopes N, *et al.* 2022. *Elife.* 11: . [PubMed](#)
28. Kim D, *et al.* 2020. *Immunity.* 53(3):581-596.e5. [PubMed](#)
29. Shen H, *et al.* 2022. *Nat Commun.* 13:5013. [PubMed](#)
30. Liu X, *et al.* 2021. *Adv Sci (Weinh).* 8:e2100233. [PubMed](#)
31. Dubrot J, *et al.* 2021. *Immunity.* 54(3):571-585.e6. [PubMed](#)
32. Strait AA, *et al.* 2021. *Commun Biol.* 4:1005. [PubMed](#)
33. Stone S *et al.* 2018. *Glia.* 66(7):1331-1345. [PubMed](#)
34. Chen EW, *et al.* 2019. *Front Immunol.* 10:1718. [PubMed](#)
35. Lai NY, *et al.* 2020. *Cell.* 180:33:00. [PubMed](#)
36. Wang Y, *et al.* 2020. *Immunity.* 53(6):1168-1181.e7. [PubMed](#)

RRID

AB_10898318 (BioLegend Cat. No. 100543)
 AB_2563052 (BioLegend Cat. No. 100563)
 AB_11219790 (BioLegend Cat. No. 100544)

Antigen Details

Structure	Ig superfamily, 55 kD
Distribution	Majority of thymocytes, T cell subset
Function	TCR co-receptor, T cell activation
Ligand/Receptor	MHC class II molecule
Cell Type	Dendritic cells, T cells, Thymocytes, Tregs
Biology Area	Immunology
Molecular Family	CD Molecules

Antigen References

1. Barclay A, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
2. Bierer BE, *et al.* 1989. *Annu. Rev. Immunol.* 7:579.
3. Janeway CA. 1992. *Annu. Rev. Immunol.* 10:645.

Gene ID

[12504](#)

Related Protocols



[Immunohistochemistry Protocol for Frozen Sections](#)

[Cell Surface Flow Cytometry Staining Protocol](#)


Other Formats

APC anti-mouse CD4, Biotin anti-mouse CD4, FITC anti-mouse CD4, PE anti-mouse CD4, PE/Cyanine5 anti-mouse CD4, Purified anti-mouse CD4, PE/Cyanine7 anti-mouse CD4, APC/Cyanine7 anti-mouse CD4, Alexa Fluor® 647 anti-mouse CD4, Alexa Fluor® 488 anti-mouse CD4, Pacific Blue™ anti-mouse CD4, Alexa Fluor® 700 anti-mouse CD4, PerCP anti-mouse CD4, PerCP/Cyanine5.5 anti-mouse CD4, Brilliant Violet 421™ anti-mouse CD4, APC/Fire™ 750 anti-mouse CD4, Brilliant Violet 570™ anti-mouse CD4, Brilliant Violet 605™ anti-mouse CD4, Brilliant Violet 650™ anti-mouse CD4, Brilliant Violet 711™ anti-mouse CD4, Brilliant Violet 785™ anti-mouse CD4, Brilliant Violet 510™ anti-mouse CD4, Purified anti-mouse CD4 (Maxpar® Ready), PE/Dazzle™ 594 anti-mouse CD4, TotalSeq™-A0001 anti-mouse CD4, TotalSeq™-B0001 anti-mouse CD4, TotalSeq™-C0001 anti-mouse CD4, Ultra-LEAF™ Purified anti-mouse CD4, Spark Violet™ 423 anti-mouse CD4 (L3T4) Antibody


Product Data



C57BL/6 mouse splenocytes were stained with CD3 APC and CD4 (clone RM4-5) Brilliant Violet 421™ (top), or rat IgG2a, κ Brilliant Violet 421™ isotype control (bottom).



C57BL/6 mouse thymocytes were fixed with 1% paraformaldehyde (PFA), and then stained with 5 μg/ml of CD4 (clone RM4-5) Brilliant Violet 421™ (blue) for 30 minutes at room temperature. Nuclei were counterstained with Propidium iodide and are shown in red. The image was captured by 40X objective.



C57BL/6 mouse frozen spleen section was fixed with 4% paraformaldehyde (PFA) for 10 minutes at room temperature and blocked with 5% FBS for 30 minutes at room temperature. Then the section was stained with 10 μ g/ml of anti-mouse CD4 (clone RM4-5) Brilliant Violet 421™ (blue), anti-mouse B220 (clone RA3-6B2) Alexa Fluor® 488 (green), and anti-mouse CD8b (clone YTS156.7.7) Alexa Fluor® 647 (red) overnight at 4°C. The image was captured by 10X objective.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587