

PE anti-human IFN- γ Antibody

Catalog# / Size	502508 / 25 tests 502510 / 50 μ g 502509 / 100 tests
Clone	4S.B3
Regulatory Status	RUO
Other Names	Interferon- γ , Immune interferon, Type II interferon, T cell interferon, Macrophage-activating factor (MAF), IFN-g, IFN-gamma
Isotype	Mouse IgG1, κ
Description	Interferon- γ is a potent multifunctional cytokine which is secreted primarily by activated NK cells and T cells. Originally characterized based on anti-viral activities, IFN- γ also exerts anti-proliferative, immunoregulatory, and proinflammatory activities. IFN- γ can upregulate MHC class I and II antigen expression by antigen-presenting cells.

Product Details

Verified Reactivity	Human
Reported Reactivity	Chimpanzee, Baboon, Cynomolgus, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Partially purified, native human IFN- γ
Formulation	μ g size: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide. test size: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
Preparation	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions.
Concentration	μ g sizes: 0.2 mg/mL test sizes: lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	ELISA or ELISPOT Detection ⁵ : The biotinylated 4S.B3 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with purified NIB42 antibody (Cat. No. 502402/502404) or purified MD-1 antibody (Cat. No. 507502/507513) as the capture antibody. Flow Cytometry ^{3,4,6-8} : The fluorochrome-labeled 4S.B3 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IFN- γ -producing cells within mixed cell populations. Additional reported applications (for the relevant formats) include: neutralization ^{1,2} , Western blotting, immunohistochemical staining of paraformaldehyde-fixed, saponin-treated tissue sections, and immunocytochemistry. The 4S.B3 antibody can neutralize the bioactivity of natural or recombinant IFN- γ . Note: For testing human IFN- γ in serum or plasma, BioLegend's ELISA Max™ Sets (Cat. No. 430101 to 430106) are specially developed and recommended.

Application References

(PubMed link indicates BioLegend citation)

1. Meager A, *et al.* 1984. *J. Interferon Res.* 4:619. (Neut)
2. Meager A, 1987. *Lymphokines and Interferons: A Practical Approach.* IRL Press Ltd, Oxford, p. 105. (Neut)
3. Sester M, *et al.* 2002. *J. Virol.* 76:3748. (ICFC)
4. Infante-Duarte C, *et al.* 2000 *J. Immunol.* 165:6107. (ICFC)
5. Goodier M, *et al.* 2000. *J. Immunol.* 165:139. (ELISA)
6. Chen H, *et al.* 2005. *J. Immunol.* 175:591. (ICFC)
7. Smeltz RB, 2007. *J. Immunol.* 178:4786. (ICFC)
8. Iwamoto S, *et al.* 2007. *J. Immunol.* 179:1449. (ICFC) [PubMed](#)
9. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (ICFC)

Product Citations

1. Ishii H, *et al.* 2016. *Sci Rep.* 6:30153. [PubMed](#)
2. Kim MY, *et al.* 2021. *JCI Insight.* 6: [PubMed](#)
3. Cuadrado E, *et al.* 2018. *Immunity.* 48:1046. [PubMed](#)
4. Wang F, *et al.* 2018. *Oncogenesis.* 7:41. [PubMed](#)
5. Hermans L, *et al.* 2021. *Front Immunol.* 634402:12. [PubMed](#)
6. Chen L, *et al.* 2018. *Immunology.* 155:123. [PubMed](#)
7. Tang W, *et al.* 2019. *Oncoimmunology.* 8:e1593807. [PubMed](#)
8. Tian X, *et al.* 2021. *Annals of Translational Medicine.* 9(9):781. [PubMed](#)
9. Zebley CC, *et al.* 2021. *Cell Rep.* 37:109796. [PubMed](#)
10. Qiu XM, *et al.* 2020. *Reproduction.* 251:159. [PubMed](#)
11. Longhi M, *et al.* 2014. *PLoS One.* 9:87956. [PubMed](#)
12. Yin Q, *et al.* 2021. *Proc Natl Acad Sci U S A.* 118: [PubMed](#)
13. Chandran P, *et al.* 2014. *J Leukoc Biol.* 96:633. [PubMed](#)
14. Fan Z, *et al.* 2020. *EMBO Mol Med.* 12:e11571. [PubMed](#)
15. Bonifacius A, *et al.* 2021. *Immunity.* 54(2):340-354.e6. [PubMed](#)
16. Szentkereszty M, *et al.* 2020. *Pathol Oncol Res.* 26:1117. [PubMed](#)
17. Yan Y, *et al.* 2018. *JCI Insight.* 3. [PubMed](#)
18. Zhang H, *et al.* 2021. *Cell Rep.* 36:109708. [PubMed](#)
19. OConnor RA, *et al.* 2021. *Oncoimmunology.* 10(1):1940675. [PubMed](#)
20. Ishii H, *et al.* 2022. *Cell Rep Med.* 3:100520. [PubMed](#)
21. Hombrink P, *et al.* 2016. *Nat Immunol.* 17:1467-1478. [PubMed](#)
22. Almeida VG, *et al.* 2017. *Braz J Med Biol Res.* 50:e5163. [PubMed](#)
23. Wing A, *et al.* 2018. *Cancer Immunol Res.* 6:605. [PubMed](#)
24. Saraiva DP, *et al.* 2018. *Front Immunol.* 2:184027778. [PubMed](#)
25. Teni Nurlatifah HR, *et al.* 2021. *Med Arch.* 75:335. [PubMed](#)
26. Guimarães F, *et al.* 2017. *J Pediatr (Rio J).* 10.1016/j.jpmed.2017.08.005. [PubMed](#)
27. Roudko V, *et al.* 2020. *Cell.* 183(6):1634-1649.e17. [PubMed](#)
28. Katsuyama E, *et al.* 2020. *Cell Reports.* 30(1):112-123.e4. [PubMed](#)
29. Hr TN, *et al.* 2021. *Allergol Immunopathol (Madr).* 49:193. [PubMed](#)
30. Gussarow D, *et al.* 2021. *Front Med (Lausanne).* 8:770381. [PubMed](#)
31. Lu H, *et al.* 2020. *Oncoimmunology.* 9:1748991. [PubMed](#)
32. Xian S, *et al.* 2021. *EMBO Rep.* 22:e52509. [PubMed](#)
33. Zanetti S, *et al.* 2016. *J Biol Chem.* 291: 16541 - 16552. [PubMed](#)
34. Chen PM, *et al.* 2022. *Sci Adv.* 8:eabo4271. [PubMed](#)
35. Faissner S, *et al.* 2022. *Front Immunol.* 13:980526. [PubMed](#)
36. Feng Y, *et al.* 2022. *EClinicalMedicine.* 43:101226. [PubMed](#)

RRID

AB_315233 (BioLegend Cat. No. 502508)
AB_315235 (BioLegend Cat. No. 502510)
AB_315234 (BioLegend Cat. No. 502509)

Antigen Details

Structure	Cytokine; dimer; 20-25 kD (Mammalian)
Bioactivity	Antiviral/antiparasitic activities; inhibits proliferation; enhances MHC class I and II expression on APC
Cell Sources	CD8 ⁺ and CD4 ⁺ T cells, NK cells
Cell Targets	T cells, B cells, macrophages, NK cells, endothelial cells, fibroblasts
Receptors	IFN- γ R α (CDw119) dimerized with IFN- γ R β (AF-1)
Cell Type	Tregs
Biology Area	Cell Biology, Immunology, Neuroinflammation, Neuroscience
Molecular Family	Cytokines/Chemokines
Antigen References	<ol style="list-style-type: none">1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook.</i> Academic Press, San Diego.2. De Maeyer E, <i>et al.</i> 1992. <i>Curr. Opin. Immunol.</i> 4:321.3. Farrar M, <i>et al.</i> 1993. <i>Annu. Rev. Immunol.</i> 11:571.

4. Gray P, et al. 1987. *Lymphokines* 13:151.

Regulation

Upregulated by IL-2, FGF-basic, EGF; downregulated by vitamin D3 or DMN; labile at pH2

Gene ID

[3458](#)

Related Protocols


[Surface and Intracellular Cytokine Staining for Flow Cytometry - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

PE anti-human IFN- γ , APC anti-human IFN- γ , FITC anti-human IFN- γ , Biotin anti-human IFN- γ , Purified anti-human IFN- γ , Alexa Fluor® 488 anti-human IFN- γ , Alexa Fluor® 647 anti-human IFN- γ , Alexa Fluor® 700 anti-human IFN- γ , Pacific Blue™ anti-human IFN- γ , PerCP/Cyanine5.5 anti-human IFN- γ , APC/Cyanine7 anti-human IFN- γ , PE/Cyanine7 anti-human IFN- γ , Brilliant Violet 421™ anti-human IFN- γ , Brilliant Violet 570™ anti-human IFN- γ , Brilliant Violet 605™ anti-human IFN- γ , Brilliant Violet 650™ anti-human IFN- γ , Brilliant Violet 711™ anti-human IFN- γ , Brilliant Violet 785™ anti-human IFN- γ , Brilliant Violet 510™ anti-human IFN- γ , PE/Dazzle™ 594 anti-human IFN- γ , APC/Fire™ 750 anti-human IFN- γ , PerCP anti-human IFN- γ , Brilliant Violet 750™ anti-human IFN- γ , KIRAVIA Blue 520™ anti-human IFN- γ Antibody, Spark NIR™ 685 anti-human IFN- γ Antibody

Product Data



PMA/Ionomycin-stimulated human PBMCs were stained with CD3 PE/Cy5 and 4S.B3 PE

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587