

Purified anti-mouse CD127 (IL-7R α) (Maxpar[®] Ready) Antibody

Catalog# / Size	135029 / 100 μ g
Clone	A7R34
Regulatory Status	RUO
Other Names	IL-7 receptor α chain, IL-7R α
Isotype	Rat IgG2a, κ
Description	CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor α chain or IL-7R α . It forms a heterodimer with the common γ chain (γ c or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be a useful marker for identifying memory and effector T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cells proliferation and development.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	IL-7Ra-IgG1 fusion protein
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and EDTA.
Preparation	The antibody was purified by affinity chromatography.
Concentration	1.0 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C.
Application	FC - Quality tested CyTOF[®] - Verified
Recommended Usage	This product is suitable for use with the Maxpar[®] Metal Labeling Kits . For metal labeling using Maxpar [®] Ready antibodies, proceed directly to the step to Partially Reduce the Antibody by adding 100 μ l of Maxpar [®] Ready antibody to 100 μ l of 4 mM TCEP-R in a 50 kDa filter and continue with the protocol. Always refer to the latest version of Maxpar [®] User Guide when conjugating Maxpar [®] Ready antibodies.
Application Notes	A7R34 is able to block clone SB/199 binding to IL-7R.
Additional Product Notes	Maxpar [®] is a registered trademark of Standard BioTools Inc.
Application References	<ol style="list-style-type: none"> 1. Sudo T, <i>et al.</i> 1993. <i>P. Natl. Acad. Sci. USA</i> 90:9125. 2. Hashi H, <i>et al.</i> 2001. <i>J. Immunol.</i> 166:3702. 3. Taylor R, <i>et al.</i> 2007. <i>J. Immunol.</i> 178:5659. 4. Mazzon C, <i>et al.</i> 2011. <i>Blood.</i> 118:2733. PubMed 5. Jin J, <i>et al.</i> 2011. <i>J. Immunol.</i> doi:10.4049/jimmunol.1001238. PubMed
(PubMed link indicates BioLegend citation)	
Product Citations	<ol style="list-style-type: none"> 1. Wei SC <i>et al.</i> 2017. <i>Cell.</i> 170(6):1120-1133. PubMed 2. Wei SC, <i>et al.</i> 2019. <i>Immunity.</i> 50:1084. PubMed 3. McDonald B, <i>et al.</i> 2020. <i>Cell Host Microbe.</i> 28(5):660-668.e4. PubMed 4. Janela B, <i>et al.</i> 2019. <i>Immunity.</i> 50:1069. PubMed 5. Zhu YP <i>et al.</i> 2018. <i>Cell reports.</i> 24(9):2329-2341. PubMed
RRID	AB_2563716 (BioLegend Cat. No. 135029)

Antigen Details

Structure	Type I transmembrane glycoprotein, associate with CD132, 60-90 kD
Distribution	Immature B cells through early pre-B stage, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, bone marrow stromal cells
Function	T cell and immature B cell proliferation and development
Ligand/Receptor	IL-7
Cell Type	B cells, T cells, Thymocytes
Biology Area	Immunology
Molecular Family	CD Molecules, Cytokine/Chemokine Receptors
Antigen References	1. Sudo T, <i>et al.</i> 1993. <i>P. Natl. Acad. Sci. USA</i> 90:9125. 2. Okuno Y, <i>et al.</i> 2001. <i>P. Natl. Acad. Sci. USA</i> 99:6246. 3. Pillai M, <i>et al.</i> 2004. <i>Leukemia Lymphoma</i> 45:2403.
Gene ID	16197


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-mouse CD127 (IL-7R α), FITC anti-mouse CD127 (IL-7R α), PE anti-mouse CD127 (IL-7R α), APC anti-mouse CD127 (IL-7R α), PE/Cyanine7 anti-mouse CD127 (IL-7R α), PE/Cyanine5 anti-mouse CD127 (IL-7R α), Alexa Fluor[®] 488 anti-mouse CD127 (IL-7R α), Alexa Fluor[®] 647 anti-mouse CD127 (IL-7R α), PerCP/Cyanine5.5 anti-mouse CD127 (IL-7R α), Biotin anti-mouse CD127 (IL-7R α), Brilliant Violet 421[™] anti-mouse CD127 (IL-7R α), Brilliant Violet 605[™] anti-mouse CD127 (IL-7R α), Purified anti-mouse CD127 (IL-7R α) (Maxpar[®] Ready), PE/Dazzle[™] 594 anti-mouse CD127 (IL-7R α), Brilliant Violet 510[™] anti-mouse CD127 (IL-7R α), Brilliant Violet 711[™] anti-mouse CD127 (IL-7R α), Brilliant Violet 785[™] anti-mouse CD127 (IL-7R α), APC/Cyanine7 anti-mouse CD127 (IL-7R α), Brilliant Violet 650[™] anti-mouse CD127 (IL-7R α), TotalSeq[™]-A0198 anti-mouse CD127 (IL-7R α), TotalSeq[™]-C0198 anti-mouse CD127 (IL-7R α), Ultra-LEAF[™] Purified anti-mouse CD127 (IL-7R α), TotalSeq[™]-B0198 anti-mouse CD127 (IL-7R α)

Product Data



C57BL/6 mouse splenocytes stained with 169Tm-anti-TCRbeta (H57-597) and 175 Lu-anti-CD127 (A7R34). Lymphocytes are displayed in the analysis. Data provided by DVS Sciences.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587