

## Recombinant Mouse IL-13 (carrier-free)

<b>Catalog# / Size</b>	575902 / 10 µg 575904 / 25 µg 575906 / 100 µg 575908 / 500 µg
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	P600, IL-13, IL1
<b>Description</b>	Mouse IL-13 was initially cloned from cDNA libraries of activated T cells and was designated as P600. IL-13 is an immunoregulatory cytokine secreted predominantly by activated T(H)2 cells, and it is a key mediator in the pathogenesis of allergic inflammation. IL-13 shares many functional properties with IL-4, and they share a common receptor subunit, the alpha subunit of the IL-4 receptor (IL-4Ralpha). IL-13 mediates its effects by interacting with a complex receptor system comprised of IL-4Ralpha and two IL-13 binding proteins, IL-13Ralpha1 and IL-13Ralpha2. Ligation of the IL-13 receptor complex results in signaling via the insulin receptor substrate (IRS)-1 and 2 and STAT-6 pathways. Interleukin-13 (IL-13), like IL-4, is a cytokine produced by T(H)2 type helper T cells in response to signaling through the T cell antigen receptor and by mast cells and basophils upon cross-linkage of the high-affinity receptor for immunoglobulin E (IgE). IL-13 has been implicated in airway hypersensitivity and mucus hypersecretion, inflammatory bowel disease, and parasitic nematode expulsion.

### Product Details

---


<b>Source</b>	Mouse IL-13, amino acids Ser26-Phe131 (Accession # NM_008355), was expressed in <i>E. coli</i> .
<b>Molecular Mass</b>	The 106 amino acid recombinant protein has a predicted molecular mass of 11,677 Da. The DTT-reduced protein migrates at approximately 9kDa and the non-reduced protein migrates at approximately 8kDa by SDS-PAGE. The N-terminal amino acid is Serine.
<b>Purity</b>	Purity is >98%, as determined by Coomassie stained SDS-PAGE.
<b>Formulation</b>	0.22 µm filtered protein solution is in 10mM NaH <sub>2</sub> PO <sub>4</sub> , 150mM NaCl, pH 7.2.
<b>Endotoxin Level</b>	Endotoxin level is <0.1 EU/µg (<0.01ng/µg) protein as determined by the LAL method.
<b>Concentration</b>	10 and 25 µg sizes are bottled at 200 µg/mL. 100 µg size and larger sizes are lot-specific and bottled at the concentration indicated on the vial. To obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.
<b>Storage &amp; Handling</b>	Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to six months, or at -70°C or colder until the expiration date. For maximum results, quick spin vial prior to opening. The protein can be aliquoted and stored at -20°C or colder. Stock solutions can also be prepared at 50 - 100 µg/mL in appropriate sterile buffer, carrier protein such as 0.2 - 1% BSA or HSA can be added when preparing the stock solution. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. <b>Avoid repeated freeze/thaw cycles.</b>
<b>Activity</b>	Recombinant mouse IL-13 induces the proliferation of TF-1 cell in a dose-dependent manner. The ED <sub>50</sub> for this effect is 0.3 - 1.5 ng/mL.
<b>Application</b>	<a href="#">Bioassay</a>
<b>Application Notes</b>	BioLegend carrier-free recombinant proteins provided in liquid format are shipped on blue-ice. Our comparison testing data indicates that when handled and stored as recommended, the liquid format has equal or better stability and shelf-life compared to commercially available lyophilized proteins after reconstitution. Our liquid proteins are verified in-house to maintain activity after shipping on blue ice and are backed by our <a href="#">100% satisfaction guarantee</a> . If you have any concerns, contact us at <a href="mailto:tech@biolegend.com">tech@biolegend.com</a> .
<b>Product Citations</b>	<ol style="list-style-type: none"><li>1. Dalmás E <i>et al.</i> 2017. Immunity. 47(5):928-942 . <a href="#">PubMed</a></li><li>2. Kakiuchi N, <i>et al.</i> 2020. Nature. 260:577. <a href="#">PubMed</a></li><li>3. Mia MM, <i>et al.</i> 2020. PLoS Biol. 18:e3000941. <a href="#">PubMed</a></li><li>4. Boothby IC, <i>et al.</i> 2021. Nature. 599:667. <a href="#">PubMed</a></li></ol>

5. Noll J, *et al.* 2016. *J Hepatol.* 64:1147-1157. [PubMed](#)
6. Alban TJ, *et al.* 2020. *Front Immunol.* 11:285416667. [PubMed](#)
7. Hu Z, *et al.* 2021. *Science.* 374:eabe6723. [PubMed](#)
8. Shepardson K, *et al.* 2016. *MBio.* 7: e00506-16. [PubMed](#)
9. Hatchwell L, *et al.* 2015. *Thorax.* 70: 854-861. [PubMed](#)
10. Fujimoto M, *et al.* 2022. *Nat Commun.* 13:5408. [PubMed](#)
11. Knox T, *et al.* 2019. *Sci Rep.* 9:6136. [PubMed](#)
12. Gomez-Lopez N, *et al.* 2021. *JCI Insight.* 6: [PubMed](#)
13. Mantsounga CS, *et al.* 2022. *Cell Rep.* 38:110309. [PubMed](#)

## Antigen Details

<b>Structure</b>	Cytokine
<b>Distribution</b>	IL-13 is secreted by activated T cells, Th2 cells, basophils, mast cells, activated eosinophils and NK cells.
<b>Function</b>	IL-13 modulates human monocytes/macrophages and B cells. IL-13 induces an increase of MHC class II expression in human monocytes/macrophages, and direct inhibition of inflammatory cytokines such as TNF- $\alpha$ and IL-1 $\beta$ .
<b>Interaction</b>	IL-13 receptors are expressed on human B cells, basophils, eosinophils, mast cells, endothelial cells, fibroblasts, monocytes, macrophages, respiratory epithelial cells, and smooth muscle cells.
<b>Ligand/Receptor</b>	IL-4Ralpha, IL-13Ralpha1, and IL-13Ralpha2
<b>Biology Area</b>	Cell Biology, Immunology, Neuroinflammation, Neuroscience
<b>Molecular Family</b>	Cytokines/Chemokines
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Brown J, <i>et al.</i> <i>J. Immunol.</i> 142:679-687 1989.</li> <li>2. McKenzie ANJ, <i>et al.</i> <i>P. Natl. Acad. Sci. USA</i> 90:3735-3739 1993.</li> <li>3. Pappasavvas E, <i>et al.</i> <i>J. Immunol.</i> 175:5532-5540 2005.</li> <li>4. Kelly-Wellch, <i>et al.</i> <i>Science Signaling</i> 293:2005.</li> <li>5. Hershey GK <i>J Allergy Clin Immunol</i> 111:677-690 2003.</li> <li>6. Harris J, <i>et al.</i> <i>Immunity</i> 27:505-517 2007.</li> <li>7. LaPorte SL, <i>et al.</i> <i>Cell</i> 132:259-272 2008.</li> </ol>
<b>Gene ID</b>	<a href="#">16163</a>

## Product Data



Recombinant mouse IL-13 induces the proliferation of TF-1 cell in a dose-dependent manner. The ED<sub>50</sub> for this effect is 0.3 - 1.5 ng/mL.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587

