

Recombinant Mouse IL-3 (Animal-Free)

Catalog# / Size	714802 / 10 µg
Regulatory Status	RUO
Other Names	Interleukin-3, Burst promoting activity, Eosinophil colony stimulating factor (Eo-CSF), Hematopoietic cell growth factor (HCGF), Mast (MGF/MCGF), Multi-colony stimulating (Multi-CSF), P cell stimulating activity (PCSA), Thy1 inducing factor
Description	Interleukin-3 is a hematopoietic growth factor involved in the survival, proliferation, and differentiation of multipotent hemopoietic cells. IL-3 is the most potent growth factor for basophils, followed by granulocyte-macrophage colony-stimulating factor and IL-5. These cytokines also act on mature basophils through specific receptors, thereby mediating adhesion, migration, and releasability. IL-3 is highly expressed by mast cells, and large and rapidly released amounts of autocrine IL-3 production is responsible for mast cell survival by IgE in the absence of antigen. IL-3 has also been implicated in the pathogenesis of several chronic inflammatory diseases, including asthma, atherosclerosis, and neurodegenerative disorders, such as multiple sclerosis. IL-3 stimulates colony formation of megakaryocytes, neutrophils, and macrophages from bone marrow cultures. IL-3 plays a vital role in stimulating basophils and mast cell responses to parasite infections, and since basophils play an important role in Th2 immune responses, IL-3 is a critical regulator of allergic inflammation. IL-3 is expressed in the major embryonic vessels and regulates the survival and proliferation of hematopoietic stem cells in the early stages of embryonic development.

Product Details

Source	Mouse IL-3, amino acids Asp33-Cys166 (Accession# NM_010556), was expressed in <i>E. coli</i> .
Molecular Mass	The 135 amino acid N-terminal methionylated recombinant protein has a predicted molecular mass of 15.2 kD. The predicted N-terminal amino acid is Met.
Purity	>98%, as determined by Coomassie stained SDS-PAGE and HPLC analysis.
Formulation	Lyophilized, carrier-free.
Endotoxin Level	Less than 0.1 ng per µg of protein.
Storage & Handling	Unopened vial can be stored at -20°C or -70°C. For maximum results, quick spin vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Do not vortex. It is recommended to further dilute in a buffer, such as 5% Trehalose, and store working aliquots at -20°C to -80°C. Avoid repeated freeze/thaw cycles.
Activity	ED ₅₀ is ≤ 0.05 ng/ml, corresponding to a specific activity of ≥ 2.0 x 10 ⁷ units/mg as determined by the dose-dependent stimulation of the proliferation of mouse NFS-60 cells.
Application	Bioassay

Antigen Details

Structure	Cytokine
Distribution	Activated T cells, Th1, Th2, mast cells, eosinophils, keratinocytes, NK cells, and endothelial cells.
Function	IL-3 is a pleiotropic factor which promotes both self-renewal and differentiation of early multipotential progenitors and is involved in inducible hematopoiesis in response to infections.
Interaction	Erythroid cells, megakaryocytes, neutrophils, eosinophils, basophils, mast cells, monocytic lineages.
Ligand/Receptor	Heterodimer IL-3Rα (CD123) a common β subunit (hβc) that is shared with IL-5R/GM-CSFR.
Cell Type	Hematopoietic stem and progenitors
Biology Area	Cell Biology, Stem Cells

Molecular Family

Cytokines/Chemokines

Antigen References

1. Kitamura T, *et al.* 1989. *J. Cell. Physiol.* 140:323.
2. Kohno M, *et al.* 2005. *Blood* 105:2059.
3. Murphy JM and Young IG. 2006. *Vitam. Horm.* 74:1.
4. Kleemann R, *et al.* 2008. *Cardiovasc. Res.* 79:360.
5. Valent P, and Dahinden CA. 2010. *Curr. Opin. Hematol.* 17:60.
6. Mirza S, *et al.* 2010. *J. Biol. Chem.* 285:22370.

Gene ID[16187](#)

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587