

TotalSeq™-C0130 anti-mouse Ly-6A/E (Sca-1) Antibody

Catalog# / Size	108151 / 10 µg
Clone	D7
Regulatory Status	RUO
Other Names	Sca-1
Isotype	Rat IgG2a, κ
Barcode Sequence	TTCCTTTCCTACGCA
Description	Ly-6A/E, also known as Sca-1, is an 18 kD member of the Ly-6 multigene family. Ly6A/E is a glycosylphosphatidylinositol (GPI)-linked protein expressed on hematopoietic stem cells. In mice expressing the Ly-6.2 haplotype (e.g., AKR, C57BL, C57BR, DBA/2, SJL, SWR, and 129), Ly-6A/E is also expressed on peripheral B lymphocytes and thymic and peripheral T lymphocytes. Strains expressing the Ly-6.1 haplotype (e.g., BALB/c, CBA, C3H/He, DBA/1, and NZB) have low Ly-6A/E expression on resting peripheral lymphocytes. The expression of Ly-6A/E on lymphocytes is upregulated upon activation from both Ly6.1 and Ly6.2 haplotype mice. Ly-6A/E is thought to be involved in the regulation of both T and B cell responses.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	IL-2-dependent mouse T-cell line (CTL-L)
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA
Preparation	The antibody was purified by chromatography and conjugated with TotalSeq™-C oligomer under optimal conditions.
Concentration	0.5 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	PG - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis and the oligomer sequence is confirmed by sequencing. TotalSeq™-C antibodies are compatible with 10x Genomics Chromium Single Cell Immune Profiling Solution.</p> <p>To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.</p> <p>Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.</p>
Application Notes	<p>The D7 antibody has been reported to induce T cell activation and inhibit TCR-induced IL-2 production. Additional reported applications (for the relevant formats) include: Western blotting^{1,2}, immunoprecipitation¹, <i>in vitro</i> lymphocyte activation³⁻⁶, induction of redirected lysis⁷, induction of T cell inhibitory signalling⁸, immunofluorescence⁹, and immunohistochemical staining of acetone-fixed frozen sections¹³ and Bouin-fixed, paraffin-embedded samples⁹.</p> <p>The two Sca-1 recognizing clones D7 and E13-161.7 have been shown to bind distinct epitopes</p>

due to the inability of D7 to block the binding of E13-161.7.¹⁴

Additional Product Notes

TotalSeq™ reagents are designed to profile protein levels at a single cell level following an optimized protocol similar to the CITE-seq workflow. A compatible single cell device (e.g. [10x Genomics Chromium System and Reagents](#)) and sequencer (e.g. Illumina analyzers) are required. Please contact [technical support](#) for more information, or visit [biolegend.com/totalseq](#).

The barcode flanking sequences are CCGAGATGTGTATAAGAGACAGNNNNNNNNNN (PCR handle), and NNNNNNNNCCCATATAAGA*A*A (capture sequence). N represents either randomly selected A, C, G, or T, and * indicates a phosphorothioated bond, to prevent nuclease degradation.

View more applications data for this product in our [Scientific Poster Library](#).

Application References

(PubMed link indicates BioLegend citation)

1. Ortega G, *et al.* 1986. *J. Immunol.* 137:3240. (WB, IP)
2. Palfree RGE, *et al.* 1986. *Immunogenetics* 23:197. (WB)
3. Codias EK, *et al.* 1990. *J. Immunol.* 144:2197.
4. Malek TR, *et al.* 1986. *J. Exp. Med.* 164:709.
5. Codias EK, *et al.* 1990. *J. Immunol.* 145:1407.
6. Ivanov V, *et al.* 1994. *J. Immunol.* 153:2394.
7. Karlhofer FM, *et al.* 1991. *J. Immunol.* 146:3662.
8. Fleming T, *et al.* 1994. *J. Immunol.* 153:1955.
9. van Bragt MPA, *et al.* 2005. *Biol. Reprod.* 73:634. (IF, IHC)
10. Umland O, *et al.* 2007. *J. Immunol.* 178:4147.
11. Cridland SO, *et al.* 2009. *Blood Cell. Mol. Dis.* 45:149. (FC) [PubMed](#)
12. Pronk CJ, *et al.* 2011. *J. Exp Med.* [PubMed](#)
13. English A, *et al.* 2000. *J. Immunol.* 165:3763. (IHC)
14. Bamezai A and Rock KL. 1995. *Proc. Natl. Acad. Sci. USA* 92:4294.
15. Wiesner DL, *et al.* 2015. *PLoS Pathog.* 11:1004701. [PubMed](#)

RRID

AB_2832370 (BioLegend Cat. No. 108151)

Antigen Details

Structure	Ly-6 multigene family, 18 kD
Distribution	Hematopoietic stem cells, activated T cells and B cells, subset of resting B cells and T cells
Function	Regulates B and T cell responses
Cell Type	B cells, Hematopoietic stem and progenitors, Mesenchymal Stem Cells, T cells
Biology Area	Immunology, Stem Cells
Antigen References	<ol style="list-style-type: none">1. Rock KL, <i>et al.</i> 1989. <i>Immunol. Rev.</i> 111:195.2. Morrison SJ, <i>et al.</i> 1994. <i>Immunity</i> 1:661.3. Spangrude GJ, <i>et al.</i> 1988. <i>J. Immunol.</i> 141:3697.4. Malek T, <i>et al.</i> 1986. <i>J. Exp. Med.</i> 164:709.
Gene ID	110454

Related Protocols

[TotalSeq™-B or -C with 10x Feature Barcoding Technology](#)

Other Formats

APC anti-mouse Ly-6A/E (Sca-1), Biotin anti-mouse Ly-6A/E (Sca-1), FITC anti-mouse Ly-6A/E (Sca-1), PE anti-mouse Ly-6A/E (Sca-1), PE/Cyanine5 anti-mouse Ly-6A/E (Sca-1), Purified anti-mouse Ly-6A/E (Sca-1), PE/Cyanine7 anti-mouse Ly-6A/E (Sca-1), Alexa Fluor® 488 anti-mouse Ly-6A/E (Sca-1), Alexa Fluor® 647 anti-mouse Ly-6A/E (Sca-1), Pacific Blue™ anti-mouse Ly-6A/E (Sca-1), Brilliant Violet 421™ anti-mouse Ly-6A/E (Sca-1), PerCP anti-mouse Ly-6A/E (Sca-1), PerCP/Cyanine5.5 anti-mouse Ly-6A/E (Sca-1), APC/Cyanine7 anti-mouse Ly-6A/E (Sca-1), Brilliant Violet 510™ anti-mouse Ly-6A/E (Sca-1), Brilliant Violet 711™ anti-mouse Ly-6A/E (Sca-1), Brilliant Violet 605™ anti-mouse Ly-6A/E (Sca-1), Purified anti-mouse Ly-6A/E (Sca-1) (Maxpar® Ready), PE/Dazzle™ 594 anti-mouse Ly-6A/E (Sca-1), Brilliant Violet 785™ anti-mouse Ly-6A/E (Sca-1), Alexa Fluor® 700 anti-mouse Ly-6A/E (Sca-1), Brilliant Violet 650™ anti-mouse Ly-6A/E (Sca-1), APC/Fire™ 750 anti-mouse Ly-6A/E (Sca-1), TotalSeq™-A0130 anti-mouse Ly-6A/E (Sca-1), TotalSeq™-B0130 anti-mouse Ly-6A/E (Sca-1), TotalSeq™-C0130 anti-mouse Ly-6A/E (Sca-1)

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587